





70
60
50
40
30
20
10
0
Fossil-poor siliciclastic siltstones
Fossil-rich calcareous siltstones
gap due to non-exposure
Key
mudstone
grey siltstone
cementstone
fish bone or scale
broken shells
ostracod
plant fragments
strap-like leaves
wood
bivalve
bioturbation
intraclasts
convolute bedding
bed 7
bed 6
beds 1-4
cms
Cl
St
VF
F
M
Mst
Sst
Erosive-based limestone with higher proportion of siliciclastic grains and diverse ichnofauna. Intensely bioturbated bed originally deposited from a high-density current/debris flow.
Single clast-rich, clast-supported limestone bed with carbonate mud clasts and quartz grains. Deposited from a high-density turbidity current.
Dolomite-cemented fining-upward lime-stones. Deposited from multiple turbidity currents.



A
C
C
B
3
2
1
S
S
S
C
F
P
C
C
F
P
C
D
B
B
L
B

a
b
b1
b2
b3
b4
c
c1
c2
c3
c4
d
b1
b2
b3
b4
b5
e
e1
e2
e3
e4
e5
e6
f
f1
f2
f3
f4
g
g1
g2
g3
g4
h
h1
h2
h3
h4
i
i1
i2
i3
i4
i5
j
k


a
lingual
distal
symphyseal
labial
mesial
basal
occlusal


b
b1
b2
b3
b4
b5
b6


c
c1
c2
d
e
e1
e2
f
g


h
i
i1
i2
i3
j
j1
j2
j3
j4
j5
j6

a
b
c
c1
c2
d
d1
d2
e
e1
e2
f
f1
f2
g
h
i
i1
upper
i2
lower
l
l1
l4
symphyseal
mesial
j
k
k1
k2
k3
l2
l3
m
m1
m2
m3
m4
m5
m6
n
n1
n2
n3
n4
o
o1
o2
o3
p
p1
p2
q
q1
q2
q3
q4
q5
q6
r
s
s1
s2

a
a1
a2
a3
a4
a5
b
b1
b2
b3
b4
b5
c
d
d1
d2
d3
d4
d5
d6
d7
d8
d9
d10
e
e1
e2
e3
e4
e5
e6
e7
f
f1
f2
f3
f4
f5
f6
f7
f8
f9
f10
g
g1
g2
g3
g4
g5
g6
g5
g6
g7
g8
g9
g10

a

a1

a2

a3

a4

a5

a6

b

b1

b2

b3

b4

c

d

d1

d2

d3

d4