The Bajocian (Middle Jurassic): a key interval in the early Mesozoic phytoplankton radiation

Nickolas J. Wiggan [a.b*], James B. Riding [b], Robert A. Fensome [c], Emanuela Mattioli [d]

[a]Department of Earth Sciences, University of Cambridge, Downing Street, Cambridge, CB2 3EQ UK.

[b]British Geological Survey, Environmental Science Centre, Nicker Hill, Keyworth, Nottingham, NG12 5GG, UK.

[c]Natural Resources Canada, Geological Survey of Canada (Atlantic), Bedford Institute of Oceanography, Dartmouth, NS B2Y 4A2, Canada.

[d]Laboratoire CNRS UMR 5125, Université Claude Bernard Lyon 1, Campus de la Doua, F-69622 Villeurbanne Cedex, France

*Corresponding author email: njw56@cantab.net

ABSTRACT

Dinoflagellates and coccolithophores are two of the most important groups of phytoplankton in the modern oceans. These groups originated in the Triassic and radiated through the early Mesozoic, rising to ecological prominence. Within this long-term radiation, important short-term intervals of evolutionary and ecological change can be recognised. The Bajocian (Middle Jurassic, ~170–168 Ma) was characterised by an important ecological transition within the coccolithophores, and the radiation of one of the principal families of cyst-forming dinoflagellates, the Gonyaulacaceae. During the Early Bajocian, the coccolith genus *Watznaueria* diversified and expanded ecologically to dominate coccolith floras, a situation which continued for the remainder of the Mesozoic. This pattern was paralleled within dinoflagellate cyst floras by the ecological dominance of the genus *Dissiliodinium* in the mid-palaeolatitudes. These phenomena appear to be linked to a positive carbon isotope shift, and

an interval of enhanced productivity driven by a shift to a more humid climate, enhanced continental weathering and nutrient flux, or by changes in ocean circulation and upwelling. The latest Early Bajocian to earliest Bathonian was then characterised by the rapid increase in diversity of dinoflagellate cysts within the family Gonyaulacaceae. Through this interval, the Gonyaulacaceae transitioned from being a relatively minor component of dinoflagellate cyst floras, to becoming one of the prominent groups of cyst-forming dinoflagellates, which has persisted to the Holocene. In Europe, the pattern of this radiation was strongly influenced by sea level, with the increase in gonyaulacacean diversity reflecting a major second-order transgression. On a finer scale, the main pulses of first appearances correlate with third-order transgressive episodes. A rise in sea level, coupled with changes in the tectonic configuration of ocean gateways, appears to have controlled the pattern of plankton diversification in Europe. These palaeoceanographic changes may have enhanced water-mass transfer between Europe, the northwest Tethys Ocean and the Hispanic Corridor, which promoted the floral interchange of dinoflagellates. Whilst sea level rise and associated large-scale palaeoenvironmental shifts appear to have controlled the pattern of dinoflagellate cyst appearances in several regions outside Europe, there is no direct correlation between dinoflagellate cyst diversity and sea level rise on a global scale. Although the Bajocian was transgressive in several regions, widespread flooded continental area was also present throughout the preceding Aalenian, an interval of low gonyaulacacean diversity. Moreover, although the Middle Jurassic was an interval of major climatic cooling, there was a ~5 myr gap between the onset of cooling and the radiation of gonyaulacaceans during the Bajocian. The Bajocian was, however, marked by a major evolutionary radiation in the pelagic realm, including ammonites, giant suspension feeding fishes and planktonic foraminifera. These phenomena may indicate an underlying ecological driver to the radiation of dinoflagellates

during the Bajocian evolutionary explosion which could represent an extension of the Mesozoic Marine Revolution.

Keywords: dinoflagellates; coccolithophores; radiation; Mesozoic Marine Revolution

1. Introduction

Dinoflagellates and coccolithophores are two of the most important groups of primary producers in the modern oceans, forming a significant part of the base of ocean food webs. Moreover, the downward flux of phytoplankton to the interior of the oceans forms a significant component of the biological pump. Dinoflagellates and coccolithophores are part of the polyphyletic 'red lineage' of phytoplankton, as they utilize chlorophyll *c* as a primary accessory pigment (Falkowski et al., 2004a).

While it has been argued that on the basis of molecular and biogeochemical evidence, the coccolithophores and dinoflagellates have a Neoproterozoic origin, they are essentially a product of the post-Palaeozoic world, rising to ecological prominence through the Mesozoic (Fensome et al., 1996; Falkowski et al., 2004a, 2004b; Bown, 2005; Janouškovec et al., 2017). The early Mesozoic, defined here as the Middle Triassic to Middle Jurassic (~247–164 Ma), represents a critical interval in the evolution of the modern phytoplankton, when the coccolithophores and dinoflagellates emerged and underwent major evolutionary radiations (Fensome et al., 1996; Falkowski et al., 2004b; Bown et al., 2004). While many authors have reviewed the broad macroecological and macroevolutionary trajectories of the eukaryotic phytoplankton during the Mesozoic–Cenozoic (e.g. Bown et al., 2004, Falkowski et al., 2004a,b, Bown, 2005, Falkowski and Knoll 2007), many short-term diversifications within this broader radiation remain poorly documented.

In this contribution, we review the pattern of coccolithophore and dinoflagellate evolution through the early Mesozoic, primarily from the perspective of the fossil record, focussing on the Bajocian (Middle Jurassic, ~170–168 Ma [Ogg et al., 2016]). The driving mechanisms behind ecological and evolutionary change in phytoplankton during the Bajocian remain poorly understood, but it is clearly a critical interval in the evolutionary history of the two groups. During the Bajocian, coccolith floras became ecologically dominated by the genus *Watznaueria*, a condition that prevailed for the remainder of the Mesozoic (Giraud et. al., 2016; Aguado et al., 2017). Among the dinoflagellates, the Bajocian was characterised by the diversification of the family Gonyaulacaceae, which became one of the most prominent groups of encysting dinoflagellates ranging through to the present day (Fensome et al., 1996; Feist-Burkhardt and Götz, 2016; Wiggan et al., 2017). To set the Bajocian radiation in context, we provide an overview of the coccolith and dinoflagellate cyst fossil records, documenting the evolutionary history of the two groups from their emergence in the Triassic to their radiation through the Early–Middle Jurassic.

1.1. Dinoflagellates

Dinoflagellates are a diverse group of flagellated eukaryotes, represented by around 2,400 extant species, of which ~2000 are marine plankton (Fensome et al., 1996; de Vernal and Marret, 2007; Janouškovec et al., 2017). Dinoflagellates are characterised by the possession of a unique nucleus, the dinokaryon; and/or by having a life cycle stage with two dissimilar flagella, one transverse and one longitudinal, which allow the motile cell to propel itself through water (Fensome et al., 1996). The group can be broadly split into naked and thecate forms. Thecate (or armoured) dinoflagellates have a series of cellulosic plates held within vesicles (alveolae) in the cortex (amphiesma) of the cell. The arrangement of the thecal plates

forms a pattern, known as tabulation, tends to be consistent within taxa and is thus a critical feature in dinoflagellate classification (Dale, 1983; Evitt, 1985; Fensome et al., 1993, 1999; Janouškovec et al., 2017). It is usually reflected, at least partially, on the cysts that comprise the vast majority of fossil dinoflagellates. Thus, tabulation is used to classify both fossil and living dinoflagellates (Fensome et al., 1993).

Around half of all dinoflagellates are autotrophic or mixotrophic and, together with the diatoms and coccolithophores, are responsible for a significant proportion of primary production in the world's oceans (Delwiche, 2007). Heterotrophic dinoflagellates are an important component of the microzooplankton (Calbet, 2008). Around 13% of modern dinoflagellate species produce a zygotic resting cyst (Head, 1996). The wall of most preservable dinoflagellate cysts is composed of the complex macrobiomolecule dinosporin (Head, 1996; Bogus et al., 2014). The fossil record of dinoflagellates is based on their (refractory) resting cysts, adding a layer of complexity to the dinoflagellate fossil record with respect to other phytoplankton groups, as the dinoflagellate motile stage is virtually never preserved, as the theca is extremely labile (Evitt, 1985; Versteegh et al., 2004). With only around 13% of living motile species producing preservable cysts, the dinoflagellate cyst fossil record is inherently incomplete, and it is not known whether this percentage has changed through time (Head, 1996; Fensome et al., 1996; Riding and Lucas-Clark 2016).

Some earlier workers cast doubt on the ability of a highly selective fossil record to shed light on dinoflagellate evolution (Evitt, 1985). However, some higher taxa are far better represented than others in the record, notably the orders Gonyaulacales and Perdiniales (Fensome et al., 1999; de Vernal and Marret, 2007). Such groups have a much richer and more extensive fossil record than would be suggested by the low overall proportion of encysting dinoflagellates in the Recent (Fensome et al.,1999). Utilising patterns of dinoflagellate cyst occurrences in the fossil record (Fensome et al. 1996) along with quasi-

cladistic analyses of all dinoflagellates (i.e. the basis of the phylogenetic classification of Fensome et al. [1993]) provide significant evidence for the general evolutionary pattern. For example, that the appearance and expansion of dinoflagellates during the early Mesozoic is a true evolutionary radiation is compellingly shown by the extent of morphological 'experimentation', particularly in the Early–Middle Jurassic. Moreover, the mixing of features in individual taxa that would subsequently characterise separate higher-level taxa (e.g. *Nannoceratopsis* is part dinophysialean and part peridiniphycidean) and later morphological innovations becoming restricted to lower taxonomic ranks (Piel and Evitt, 1980; Fensome et al, 1996). The latest molecular data supports this general scenario, as tentatively does biogeochemical data (Janouškovec et al. 2016).

The radiation of dinoflagellates through the early Mesozoic was characterised by the appearance of stem forms with a gonyaulacoid-peridinoid tabulation, and ultimately by crown thecate clades, such as the Gonyaulacales and Peridiniales (Fensome et al., 1996; Janouškovec et al., 2017). The increase in dinoflagellate cyst diversity through the Mesozoic was accompanied by a synchronous increase in the abundance of dinosteranes in Mesozoic sedimentary rocks (Moldowan et al., 1996). Triaromatic dinosteranes are types of lipid biomarkers derived from dinosterols, and are primarily associated with dinoflagellates (Moldowan and Talyzina, 1998). However, recent molecular phylogenetic research has revealed that the acquisition of dinosterols within the dinoflagellate lineage occurred during the Mesozoic, shedding doubt on the reliability of dinosteranes as an unequivocal biomarker for dinoflagellates in pre-Mesozoic rocks (Janouškovec et al., 2017). Despite this, the concomitant increase in the abundance of dinosteranes and increase in dinoflagellate cyst diversity through the Jurassic strongly suggests that dinoflagellates rose to ecological prominence in the early Mesozoic. In summary, the appearance of high-level dinoflagellate taxa, which are supported by molecular data as distinct clades, and progressive innovations at

lower taxonomic ranks, along with transitionary forms such as *Nannoceratopsis,* and an increase in dinosterane abundance, strongly suggest that dinoflagellates underwent a true evolutionary radiation through the early Mesozoic (Fensome et al., 1996).

1.2. Coccolithophores

Coccolithophores are unicellular phytoplankton belonging to the division Haptophyta. They are defined by a cell-bounding series of calcareous plates, known as coccoliths, which interlock to form a coccosphere. Coccolithophores form part of a wider clade of noncalcifying/potentially calcifying/calficying haptophytes termed the Calcihaptophycidae (de Vargas et al., 2007). Coccolithophores are the most important calcifying organisms in the modern ocean and thus the downward flux of coccolithophores to the sea floor contributes significantly to the biological pump, making coccolithophores an important agent of oceanic biogeochemical cycles (Brownlee and Taylor, 2004; Rost and Riebesell, 2004).

For many species of coccolithophore, coccolith morphology varies with the haploid-diploid lifecycle. During the diploid phase, cells produce heterococcoliths, formed from a structurally complex arrangement of radial calcite crystals. In contrast, during the haploid phase, cells are naked (such as for the blooming species *Emiliania huxleyi*) or holococcoliths are produced, which are characterised by simple calcite rhombohedra.

The fossil record of coccolithophores is primarily based on coccoliths, as coccospheres are rarely preserved intact. Moreover, the record comprises primarily of heterococcoliths (de Vargas et al., 2007). Although fossil holococcoliths are known, the nature of the tiny calcite rhombohedra from which they are constructed makes them extremely susceptible to dissolution (Crudeli et al., 2006 and references therein). Hence, the dearth of fossil holococcoliths reflects a substantial taphonomic bias.

Within the early Mesozoic coccolith record, heterococcoliths are represented by two broad morphological groups, the muroliths and the placoliths (Bown, 1987; Bown et al., 2004). Muroliths represent the earliest coccoliths, and are characterised a simple structure with calcite elements extending vertically (Bown, 1987; Gardin et al., 2012). In the Early Jurassic, placolith coccoliths appeared and radiated to become the dominant coccolith group. The placolith structure is defined by two discs which lie on top of one another, and are connected by a central pillar with radial calcite elements (Bown, 1987).

Unlike dinoflagellates, there is no appreciable pre-Cenozoic biomarker record of the coccolithophores. Coccolithophores produce alkenones which are widely used in Cenozoic palaeoceanography as temperature and pCO_2 proxies. However, the oldest alkenones are only known from marine sediments of Early Cretaceous age (Knoll et al., 2007). Consequently, there is no alkenone record of coccolithophores through the Early Mesozoic, making the coccolith fossil record the only tool for directly tracking coccolithophore evolution through deep time.

2. Middle Triassic to Early Jurassic: emergence and radiation of dinoflagellates and coccolithophores

2.1. Emergence in the Middle to Late Triassic

The onset of the Mesozoic phytoplankton radiation in the fossil record is signalled by the appearance in Australia of the earliest unequivocal dinoflagellate cyst species, *Sahulidinium ottii* in the Middle Triassic (late Ladinian, ~238 Ma) (Riding et al., 2010a). Dinoflagellate cyst diversity increased through the Middle to Late Triassic with the appearances of the families Rhaetogonyaulacaceae and Suessiaceae (Figure 1; Fensome et al., 1996). The former represents the first appearance of the order Gonyaulacales, one of the the

two main orders represented by fossil dinoflagellate cysts (Fensome et al., 1999; Janouškovec et al., 2017). This later Triassic expansion of dinoflagellate cysts coincides with an increase in the abundance of dinosteranes (Moldowan et al., 1996), suggesting that dinoflagellates began their rise to ecological prominence through this interval (Fensome et al., 1996).

The earliest coccoliths appeared in the latest Norian (~209 Ma), with the oldest identifiable species, *Crucirhabdus minutus*, appearing in the earliest Rhaetian (~207 Ma) of the Austrian Alps (Gardin et al., 2012). These early coccoliths are extremely small (2–3 μm), and have a murolith structure (Bown, 1987; Bown and Young, 1998; Bown et al., 2004). *Crucirhabdus* is also known from the Rhaetian of Australia, thus indicating a global distribution for coccoliths during the Late Triassic (Bralower et al., 1992; Gardin et al., 2012). However, diversity remained low during this interval, with a maximum of five species (Bown et al., 2004). Although there is no biomarker record of coccolithophores through the early Mesozoic, molecular divergence times are broadly in agreement with the coccolith fossil record. Molecular clocks based on SSU and LSU rDNA give ~270–240 and ~ 200 Ma respectively as the earliest possible dates for the origin of the coccolithophores (de Vargas et al. 2007; Medlin et al., 2008).

Falkowski et al. (2004a, b) argued that the emergence of the 'red lineage' of phytoplankton during the Triassic was driven by changes in ocean chemistry in the aftermath of the Permo-Triassic mass extinction. These authors postulated that prolonged periods of oceanic anoxia during the Early Triassic forced heterotrophic cells to acquire and retain plastids. In this hypothesis, prolonged euxinic conditions may have induced widespread denitrification, generating a paucity of fixed nitrogen, one of the most important macronutrients in the Triassic seas. Although photoautotrophs would have been affected by the limited availability of fixed nitrogen, a reduction in primary production and decline in upward transfer of biomass would have severely affected consumers. Acquisition of plastids

by heterotrophic cells, via endosymbiosis of a red alga, would have provided a mixotrophic 'duel fuel' strategy. This would have been an advantageous ecological strategy in unstable and stressed intervals of environmental change during the Early Triassic (Falkowski et al., 2004a).

2.2. The End-Triassic Mass Extinction

The end-Triassic mass extinction (~201 Ma) significantly affected the diversity of phytoplankton. Only four dinoflagellate cyst genera and one coccolith species survived this major environmental perturbation (Bown et al., 2004; van de Schootbrugge et al., 2007). The temporal coincidence between the diversity collapse at the end of the Triassic and the emplacement of the Central Atlantic Magmatic Province (CAMP) has led many workers to suggest a causal relationship between these phenomena (e.g. Schoene et al., 2010; Blackburn et al., 2013; Percival et al., 2017). Volcanic outgassing may have driven an increase in pCO_2, resulting in acidification of epicontinental seas and prolonged euxinia (Ruhl et al., 2011; Schaller et al., 2011; Hönisch et al., 2012; Richoz et al., 2012). As such, prolonged anaerobic/euxinic conditions, and ocean acidification may have driven the decline in dinoflagellate cyst and coccolith diversity (van de Schootbrugge et al., 2007; 2013).

In many areas through Europe, the collapse in dinoflagellate cyst and coccolith diversity through the Triassic–Jurassic transition is concomitant with an increase in the abundance of the phycoma (resting cysts) of prasinophytes (chlorophyte green algae) (Kuerschner et al., 2007; van de Schootbrugge et al., 2007; Bonis et al., 2009). Moreover, the earliest Hettangian (~200 Ma, earliest Jurassic) is marked by an increase in the abundance of diaryl isoprenoid molecules, derived from green sulphur bacteria (Richoz et al., 2012; van de Schootbrugge et al., 2013). The turnover in organic-walled phytoplankton communities, and

presence of green sulphur bacteria, indicates that the photic zone became euxinic (Richoz et al., 2012). Green sulphur bacteria require both light and free hydrogen sulphide (H_2S) to photosynthesise, whereas prasinophytes appear to have thrived during intervals of palaeoenvironmental stress during the Mesozoic (van de Schootbrugge et al., 2013).

2.2. Early to Middle Jurassic: recovery and radiation

Both dinoflagellates and coccolithophores recovered during the earliest Jurassic and radiated through the late Early and Middle Jurassic, with the appearances of numerous families (Figure 1; Fensome et al., 1996; Bown et al., 2004). Dinoflagellate cyst are generally sparse in Hettangian strata, with low-diversity assemblages (Wall, 1965; van de Schootbrugge et al., 2013). The succeeding Sinemurian was marked by the sudden, and short-lived, appearance of *Liasidium variabile* (see Brittain et al. 2010). The appearance of *Liasidium variable* is significant as it marks the first appearance in the fossil record of the order Peridiniales, and although its precise taxonomic affinity is unclear due to lack of a fully reflected tabulation, *Liasidium variable* may be the earliest representative of the family Peridiniaceae (Feist-Burkhardt, 2009). It is also possible that *Liasidium variabile* was heterotrophic, as it has a pigmented wall structure (Feist-Burkhardt 2009; Riding et al., 2013). The sudden appearance of this species in the Late Sinemurian has been related to the opening of the Hispanic Corridor, which allowed *Liasidium variabile* to invade the Tethys Ocean from the Panthalassa Ocean (van de Schootbrugge et al., 2005). Recent work has linked the acme of *Liasidium variabile* to a negative carbon isotope excursion, and a brief interval of global warming (Riding et al., 2013).

A significant increase in dinoflagellate cyst diversity occurred during the Late Pliensbachian, with the radiation of numerous families of gonyaulacaleans and peridinialeans

(Figure 1; Morgenroth, 1970; Feist-Burkhardt and Wille, 1992). Another important first appearance in the Late Pliensbachian was that of *Nannoceratopsis*, a genus confined to the Jurassic and the only representative of the family Nannoceratopsiaceae. *Nannoceratopsis* has a tabulation that is a combination of dinophysioid and gonyaulacoid-peridinoid, and as such appears to be an evolutionary link between the Peridiniphycidae and the Dinophysiales. The latter has virtually no fossil record (Janouškovec et al., 2017).

The most significant radiation of the coccolithophores during the Mesozoic occurred in the Early Jurassic. During that interval, 10 of the known 16 coccolith families appeared. The Hettangian to Sinemurian was characterised by the recovery and radiation of murolith forms, which had survived the end-Triassic mass extinction (Bown et al. 2004). The family Biscutaceae appeared around the Sinemurian–Pliensbachian boundary and radiated rapidly through the Pliensbachian, becoming an important component of coccolith floras (Mattioli et al., 2004). The Biscutaceae represent the emergence of the placolith morphology, making the Pliensbachian one of the most important intervals in coccolithophore evolution (de Kaenel and Bergen, 1993; Bown and Cooper, 1998; Mattioli and Erba, 1999; Mattioli et al., 2013). The emergence of the placolith morphology was extremely important because it allowed coccoliths to physically lock together to form a coccosphere for the first time. Prior to the development of placoliths, elements in the muroliths were presumed to have been bound together by organic material (Bown, 1987). The initial development of placoliths occurred in two steps. In the Early Pliensbachian, radiating placoliths first appeared; these have a simple structure comprising proximal and distal shields (Bown, 1987). In the Late Pliensbachian, more complex imbricating coccoliths with a bicyclic distal shield appeared, and this morphology dominated coccolith floras for the remainder of the Jurassic (Bown, 1987; Mattioli and Erba, 1999; Mattioli et al., 2013).

The Pliensbachian was clearly an important interval in the evolution of the dinoflagellates and coccolithophores. It was also an interval of major palaeoclimatic and palaeoceanographic change. The Late Pliensbachian was marked by cooling palaeotemperatures, with a pronounced fall in seawater temperature of ~5 ℃ (Dera et al., 2011). This cooling may have allowed continental ice sheets to develop, and the Late Pliensbachian regression may have reflected glacioeustasty (Suan et al., 2010; Bougeault et al., 2017). van de Schootbrugge et al. (2005) argued that cooling temperatures drove the radiation of dinoflagellates during the Late Pliensbachian. These authors argued that cooling increased ocean turbulence and enhanced vertical mixing, which resulted in well-ventilated bottom waters, creating conditions ideal for cyst-forming dinoflagellates. Moreover, an increase in upper ocean turbulence may have driven, larger, more frequent pulses of nutrients to the photic zone, which in turn may have favoured bloom-forming dinoflagellates (Falkowski et al., 2004b). In contrast to the dinoflagellates, the drivers behind the emergence of placolith coccoliths are unclear. The Sinemurian–Pliensbachian transition was marked by a negative carbon isotope shift, and the burial of organic-rich marine sediments (Boussaha et al., 2014; Price et al., 2016; Ruhl et al., 2016; Bougeault et al., 2017). The negative shift in carbon isotopes has been linked to volcanic outgassing from an interval of CAMP volcanism (Ruhl et al., 2016). Moreover, volcanic outgassing may driven a switch to more humid climatic conditions in Europe, with enhanced weathering and run-off (Cohen et al., 2004; Brazier et al., 2015; Ruhl et al., 2016; Bougeault et al., 2017). As such, it is likely that nutrient flux was elevated at this time. The temporal coincidence between these phenomena and the appearance of placolith coccoliths may indicate that elevated nutrient levels were a driver of coccolith evolution during the Early Jurassic. Indeed, elevated nutrient levels have been linked to evolutionary changes in coccolithophores during the Middle Jurassic (section 3; Suchéras-Marx et al., 2015). Moreover, the Early Pliensbachian was marked by sea level

rise (Hallam, 2001), which would have increased shallow-sea ecospace for neritic phytoplankton. An alternative scenario for the sudden onset of placolith coccoliths within the western Tethys may be related to the opening of the Hispanic Corridor during the Late Sinemurian–Early Pliensbachian (Aberhan, 2001; van de Schootbrugge et al., 2005). The circulation pattern through the Hispanic Corridor was most likely eastward, from the Panthalassic Ocean, which may have allowed placoliths to colonize the Tethys Ocean from the Panthalassic Ocean (Plancq et al., 2016). Alternatively, changes in ocean circulation related to the opening of the Hispanic Corridor could have stimulated the emergence of placolith coccoliths in the Tethys Ocean (Plancq et al., 2016)

The Early Toarcian was marked by the Toarcian Oceanic Anoxic Event (T-OAE) (~180 Ma), a relatively brief (~900 kyr) phase of widespread oceanic anoxia/euxinia and climate change. It was characterised by extensive, coeval organic-rich shales and a negative carbon isotope excursion, which affected the entire carbon cycle (Hesselbo et al., 2000; Kemp et al., 2005; van de Schootbrugge et al., 2005; Suan et al., 2008, 2011; Xu et al., 2017).

Dinoflagellate cyst diversity was temporarily affected by the T-OAE, as diversity declined sharply (Bucefalo Palliani et al., 2002; van de Schootbrugge et al., 2005; Correia et al., 2017a, b). Both the diversity and abundance of coccoliths declined through the T-OAE, and this was accompanied by a trend to decreasing coccolith size through the T-OAE, which has been linked to enhanced pCO_2 driving a biocalcification crisis (Mattioli et al., 2004; Tremolada et al., 2005; Suan et al., 2010) or, alternatively, the effects of pCO_2 increase were indirect, acting on sea-surface temperatures and marine stratification, giving rise to stressed conditions in surface waters (Mattioli et al., 2009; Fraguas et al., 2012). Therefore, as in the latest Triassic, organic-walled phytoplankton communities became dominated by prasinophytes during the T-OAE (van de Schootbrugge et al., 2005). Similarly, green sulphur

bacteria appear to have thrived at this time, indicating photic zone euxinia (van de Schootbrugge et al., 2013).

After the T-OAE, normal oceanographic conditions, with fully aerobic water columns, were re-established during the *Hildoceras bifrons* ammonite biozone. The majority of the dinoflagellate cyst taxa that had characterised the Late Pliensbachian reappeared. The Late Toarcian was characterised by an important evolutionary event. This was the appearance of the first Gonyaulacaceae, heralded by the emergence of *Sentusidinium* in the Late Toarcian of Yorkshire, northern England and northern Germany (Riding, 1984; Prauss, 1989). Dinoflagellate cysts belonging to the family Gonyaulaceae are defined by several characters, including the presence a particular configuration of plates on the hypocyst, termed a sexiform tabulation (Evitt, 1985). Other key characters include a mid-ventral sulcus, and a symmetrical posterior outline. Although *Sentusidinium* lacks a fully reflected tabulation, its overall morphology is strongly gonyaulacacean (Wood et al., 2016). The Gonyaulacaceae would go on to become the most important and diverse family of encysting dinoflagellates in the fossil record (section 3).

The coccolithophores recovered gradually from the T-OAE; the re-appearance of deep-dwelling coccoliths such as *Crepidolithus crassus* indicated the end of anoxic/dysoxic conditions within the deep photic zone. Subsequently, the entire calcareous nannoplankton flora recovered (Mattioli et al., 2008). The post-T-OAE recovery interval was characterised by the appearance of the genus *Watznaueria* (see Mattioli and Erba, 1999), which is significant because *Watznaueria* became ecologically and numerically the dominant Middle Jurassic to Cretaceous coccolith genus (Suchéras-Marx et al., 2015; Giraud et al., 2016). The *H. bifrons* zone was also marked by a significant increase in size of all the dominant *Lotharingius* morphospecies (Ferreira et al., 2017). This size change is partly related to a species-specific replacement within the *Lotharingius*, with small-sized morphospecies being

abundant in the Early Toarcian and large-sized morphospecies in the Middle–Late Toarcian. However, this pattern was also accompanied by a size increase of all the *Lotharingius* morphospecies across the middle Toarcian.

3. The Bajocian: a critical interval of dinoflagellate and coccolithophore evolution and ecology

3.1. Late Aalenian to Early Bajocian: expansion of Watznaueria *and the ecological dominance of* Dissiliodinium

The Bajocian was an important interval of dinoflagellate and coccolithophore ecological development and evolution. The latest Aalenian to Early Bajocian was marked by the diversification and ecological expansion of the coccolith genus *Watznaueria*, marking the beginning of 100 myr of ecological dominance by this genus within coccolithophore communities (Giraud et al., 2016). Through the Aalenian–Bajocian transition, *Watznaueria* replaced *Lotharingus* as the dominant coccolith genus (Suchéras-Marx et al., 2012, 2015; Giraud et al., 2016; Aguado et al., 2017; Ferreira et al., 2017). Suchéras-Marx et al. (2015) examined the diversification and expansion of *Watznaueria* from two areas within the European Epicontinental Seaway, namely the French Subalpine Basin and the Lusitanian Basin of Portugal. These authors recorded an increase in both the absolute and relative abundance of *Watznaueria* through the Early Bajocian, and noted that the absolute abundance of coccoliths as a whole increased through that interval, driven by the increase in the abundance of *Watznaueria* (Figure 2). The majority of other coccolith taxa did not decrease in absolute abundance, but decreased in relative abundance during the same interval, implying that *Watznaueria* did not outcompete other genera but integrated into the coccolith community. Further, the expansion of *Watznaueria* appears to have had two critical steps.

First, in the earliest Bajocian (*Hyperlioceras discites* ammonite biozone), coccolith floras were dominated by the species *Watznaueria contracta* and *Watznaueria colacicchii*. Second, towards the end of the Early Bajocian, coccolith abundances stabilised as communities became dominated by *Watznaueria britannica* and *Watznaueria* aff. *manivitiae.* Giraud et al. (2016) compiled data on coccolith records from the Late Aalenian to Early Bajocian of northwest Europe. These authors noted a similar pattern, with the increase in abundance of *Watznaueria*, beginning around the Aalenian–Bajocian transition. This initial increase in the abundance of *Watznaueria* was produced by the rise to dominance of a group of *Watznaueria* species characterised by a central cross area (Figure 3). This group comprises *Watznaueria colacichii*, *Watznaueria contracta* and *Watznaueria* aff. *Watznaueria contracta*. The next step in this transition, in the Early Bajocian, was characterised by the increase in the abundance of species lacking a central cross, and those with a central bar, including *Watznaueria britannica* and *Watznaueria* aff. *Watznaueria manivitiae* (Figure 3). Furthermore, Aguado et al. (2017) documented the coccolith record through the Late Aalenian to Early Bajocian of the southern Iberian palaeomargin. These authors reported a similar pattern, with an increase in the relative and absolute abundance of *Watznaueria* through the latest Aalenian–earliest Bajocian, particularly that of *Watznaueria contracta*. Coccolith assemblages around the middle of the Lower Bajocian (*Witchellia laeviuscula* ammonite biozone) are marked by an increase in *Watznaueria britannica* and *Watznaueria manivitiae*, an increase that persisted throughout the remainder of the Early Bajocian. Significantly, Aguado et al. (2017) noted that the Early Bajocian was marked by a decline in *Crepidolithus crassus,* a coccolith interpreted as deep-photic zone dwelling (Figure 4).

Within the dinoflagellate cysts, the Early Bajocian was characterised by the ecological dominance of the gonyaulacacean genus *Dissiliodinium* (Wiggan et al., 2017). *Dissiliodinium* first appeared in the Middle Aalenian as the species *Dissiliodinium lichenoides* (see Feist-

Burkhardt and Pross, 2010). During the latest Aalenian and Early Bajocian, *Dissiliodinium* diversified with the appearances of *Dissiliodinium caddaense, Dissiliodinium giganteum* and *Dissiliodinium psilatum* (see Helby et al., 1987; Prauss, 1989; Feist-Burkhardt, 1990; Wiggan et al., 2017). During this interval, *Dissiliodinium* was also ecologically dominant in several mid-palaeolatitude regions. *Dissiliodinium giganteum* dominated dinoflagellate cyst floras in the southern German Basin, Poland/northern Slovakia, and Switzerland (Figures 5, 6; Gedl, 2008; Hostettler et al., 2017; Wiggan et al., 2017). Outside Europe, dinoflagellate cyst floras from the Early Bajocian of offshore Western Australia were dominated by *Dissiliodinium caddaense* (Figure 6; Riding et al, 2010b). The *Dissiliodinium caddaense* dinoflagellate cyst biozone has been correlated by ammonites and strontium isotopes to the Early Bajocian *Witchellia laeviuscula* ammonite biozone of Europe (Riding et al., 2010b). Consequently, the acme of *Dissiliodinium caddaense* is synchronous with the acme of *Dissiliodinium giganteum* in Europe (Figure 6). Furthermore, *Dissiliodinium* was also abundant in the Early Bajocian of the Neuquén Basin in Argentina (Stukins et al., 2013). In this region, dinoflagellate cyst assemblages were dominated by *Dissiliodinium psilatum* (Figure 6).

3.2. Latest Early Bajocian to Early Bathonian: stabilisation of coccolithophore communities and the rapid diversification of gonyaulacacean dinoflagellates

Through the latest Early Bajocian to Early Bathonian, coccolith floras were dominated by morphotypes of *Watznaueria* defined by a central bridge, or lacking a central area (Figure 3; Giraud et al., 2016). The Early Bathonian was marked by the appearance of *Watznaeuria barnesiae*, a form belonging to the morphological group of taxa defined by lacking a central area. This group steadily increased in abundance through the Bathonian as taxa defined by a central bridge declined (Figure 3; Giraud et al., 2016). *Watznaueria barnesiae* became a

prominent component of coccolith communities from its appearance in the Early Bathonian, to the end of Cretaceous, and underwent little change in morphology over this 100 myr interval (Lees et al., 2005; Bornemann and Mutterlose, 2006; Tiraboschi and Erba, 2010).

Whereas the Late Aalenian to Early Bajocian interval marks the transition to long-term ecological dominance of *Watznaueria* within coccolithophore communities, the acme of *Dissiliodinium* within the dinoflagellate cysts was comparatively short-lived. *Dissiliodinium* declined in abundance towards the end of the Early Bajocian in Europe, and formed a relatively minor component of dinoflagellate cyst floras through the Late Bajocian (Figure 5, Wiggan et al. 2017). In Australia, *Dissiliodinium caddaense* became extinct around the Early–Late Bajocian transition, and the genus formed a more moderate component of dinoflagellate cyst floras in the Late Bajocian to Early Bathonian (Riding et al., 2010a; Mantle and Riding, 2012). As such, the acme of *Dissiliodinium* represents a small component of the wider diversification and ecological expansion of gonyaulacacean dinoflagellates, and this genus may have been out-competed by newly emerging taxa. In particular, ~50 species appeared between the latest Early Bajocian and the Early Bathonian (Feist-Burkhardt and Monteil, 1997; Wiggan et al., 2017). Wiggan et al. (2017) demonstrated that, in Europe, there was a continuous pattern of appearances through the Late Aalenian to Early Bathonian, but the highest number of first appearances were in the *Stephanoceras humphriesianum* (latest Early Bajocian) and *Parkinsonia parkinsoni* ammonite biozones (latest Bajocian) (Figures 7, 8). Moreover, from the *Stephanoceras humphriesianum* ammonite biozone onwards, appearances were dominated by gonyaulacacean taxa. Similarly, in terms of diversity, species richness steadily increased through the Late Aalenian to Early Bajocian, and there was a particularly pronounced increase in richness in the *Stephanoceras humphriesianum* and *Parkinsonia parkinsoni* ammonite biozones, as result of the large number of first appearances in these zones (Figure 9). During the Late Aalenian, gonyaulacacean taxa comprised around

25% of dinoflagellate cyst taxa, but by the earliest Bathonian, *Zigzagiceras zigzag* ammonite biozone, this family represented over 60% of all dinoflagellate cyst taxa (Figure 9). Thus, the Bajocian represents a critical interval in the evolutionary history of the Gonyaulacaceae.

The diversification of gonyaulacaceans during the Bajocian–Bathonian is observed from the low to mid palaeolatitudes of the northern and southern hemispheres, indicating a global increase in diversity. There was an increase in the diversity of gonyaulacacean dinoflagellate cysts through the Middle Jurassic in offshore eastern Canada, but in this region the majority of appearances are recorded from the Bathonian, rather than the Bajocian (Bujak and Williams, 1977). Outside of the mid-northerly palaeolatitudes, the radiation of gonyaulacaceans is observed in the Middle Jurassic of Australia, which was then located on the southern edge of the Tethys Ocean, around 30°S. The pattern of dinoflagellate cyst appearances in the Northwest Shelf of Australia is comparable to that of Europe, with a significant increase in the diversity of gonyaulacaceans from the latest Early Bajocian to Early Bathonian (Mantle and Riding, 2012). Gonyaulacaceans also diversified through the low palaeolatitude regions; there is a significant increase in the appearances and diversity of gonyaulacacean dinoflagellate cysts through the Bajocian–Bathonian of Egypt, Israel and Qatar (Conway, 1990; Ibrahim et al., 2002, 2003). Gonyaulacaceans do not appear to have radiated through the Bajocian–Bathonian in the high palaeolatitudes, as this group comprises only a minor component of dinoflagellate cyst floras from the Middle Jurassic of Arctic Canada (Figure 10; Davies, 1983). As such, the Gonyaulacaceae appear to have been taxonomically and ecologically dominant in the low to mid-palaeolatitudes.

The radiation of gonyaulacaceans was accompanied by a significant increase in the diversity of form, or disparity of dinoflagellate cysts, and innovations in the archaeopyle (the excystment aperture). Prior to the Bajocian, gonyaulacaceans were represented by *Batiacasphaera*, *Dissiliodinium, Kallosphaeridum* and *Sentusidinium* (see Feist-Burkhardt

and Wille 1992; Feist-Burkhardt and Pross, 2010). These genera are characterised by simple, mostly non-tabulate cysts (Figure 11). *Batiacasphaera*, *Kallosphaeridium* and *Sentusidinium* have apical archaeopyles, whereas *Dissiliodinium* has a multi-plate precingular archaeopyle (Figure 11). Through the Bajocian 19 gonyaulacacean genera appeared, which exhibit a wide array of morphologies (Figure 11). The Early Bajocian was marked by the diversifications of genera with multi-plate precingular archaeopyles such as *Dissiliodinium* and *Durotrigia*, and included the first gonyaulacaceans with more-or-less fully reflected tabulation, such as *Durotrigia daveyi* (Figure 11). Morphological features such as cavation (the separation of wall layers) appeared within the Gonyaulacacae, as represented by the appearances of the genera *Cavatodissiliodinium* and *Endoscrinium* (Feist-Burkhardt and Götz, 2016). Moreover, gonyaulacaceans developed ornamental features such as distally complex processes and prominent sutural crests (Figure 11). These morphological features were accompanied by an increase in the number of archaeopyle types. Over the course of the Bajocian, genera with epicystal, one-plate precingular and two-plate precingular archaeopyles appeared. While both fossil and recent dinoflagellate cyst taxa can exhibit intraspecific variability within archaeopyle formation (e.g. Evitt, 1985; Harding, 1986; Ellegaard et al., 2002), the one-plate precingular archaeopyle appears to have been developed with the Gonyaulacacae during the Bajocian, and has remained a common gonyaualacean excystment strategy to the present day. Therefore, the increase in taxonomic diversity of gonyaulacacean dinoflagellate cysts through the Bajocian was accompanied by morphological innovations in archaeopyle formation, and a rapid increase in morphological disparity.

3.3. Drivers behind the ecological and evolutionary change in dinoflagellates and coccolithophores during the Bajocian

3.3.1. The diversification and expansion of Watznaueria*, and the ecological dominance of* Dissiliodinium

The diversification and expansion of *Watznaueria* during the Early Bajocian was coincident with a Europe-wide positive carbon isotope excursion (CIE). This excursion has been reported from central Italy (Bartolini et al., 1996, 1999; Bartolini and Cecca, 1999), the Isle of Skye, Scotland, (Jenkyns et al., 2002; Korte et al., 2015), Yorkshire, northern England (Hesselbo et al., 2003), southern Spain (O'Dogherty et al., 2006), northern France (Brigaud et al., 2009), Portugal (Suchéras-Marx et al., 2012), south-eastern France (Suchéras-Marx et al., 2013; Giraud et al., 2016), and most recently, Morocco (Bodin et al., 2017). The positive shift in $\delta^{13}C$ was coincident with an increase in biosiliceous sedimentation through the western Tethys; which may reflect a eutrophication of surface waters throughout the Western Tethys during the Early Bajocian (Figures 3, 4; Bartolini and Cecca, 1999; Bartolini et al., 1999; O'Dogherty et al., 2006; Suchéras-Marx et al., 2015; Giraud et al., 2016; Aguado et al., 2017).

This Early Bajocian phase of elevated nutrient levels has been invoked as the causal mechanism behind the ecological expansion of *Watznaueria* (see Suchéras-Marx et al., 2015; Giraud et al., 2016; Aguado et al., 2017). Suchéras-Marx et al. (2015) suggested that the increase in the abundance of *Watznaueria,* and coccoliths as a whole may reflect an increase in ecosystem carrying capacity driven by enhanced nutrient levels. Moreover, the two-step pattern of the diversification of *Watznaueria* may represent an increase in nutrient levels, with the initiation of the expansion of *Watznaueria* triggered by the increase in nutrient levels. The second step reflects the acquisition of a bloom-forming ecological strategy by

Watznaueria britanica and *Watznaueria* aff. *Watznaueria manivitiae* (see Suchéras-Marx et al., 2015; Giraud et al., 2016).

Two possible scenarios have been invoked as the drivers behind an increase in nutrient levels through the Late Aalenian to Early Bajocian (Suchéras-Marx et al., 2015;Giraud et al., 2016; Aguado et al., 2017). The first hypothesis is a climatically driven increase in continental weathering and nutrient flux during the Early Bajocian. The Middle Jurassic is increasingly being recognised as a 'cool mode' of the Jurassic climate (Korte et al., 2015). In Europe, oxygen isotope data from belemnites, bivalves and brachiopods, indicate that there was a cooling of seawater temperature of ~10°C across the Early–Middle Jurassic transition, and relatively cool temperatures persisted until the Bathonian (Figure 12; Korte et al., 2015).The initiation of cooling through the Early–Middle Jurassic transition is thought to have been driven by the magmatic doming in the North Sea, which blocked water-mass exchange through the Viking Corridor, and resulted in the thermal isolation of the Boreal Sea (Korte et al., 2015). However, a possible short-term interval of warming may have occurred during the latest Early Bajocian (*S. humphriesianum* ammonite biozone) in Europe (Dera et al., 2011; Korte et al., 2015; Giraud et al., 2016). This has been argued to reflect an increase in latitudinal temperature gradient between the mid and high palaeolatitudes (Suchéras-Marx et al., 2015). A more humid climate in Europe during the Early Bajocian is evidenced by the switch from charcoal to coal as the dominant preservational mode of fossil wood through the Upper Aalenian to Lower Bajocian of the Cleveland Basin of Yorkshire (Hesselbo et al., 2003). Changes in clay mineral assemblages may also be indicative of a warm, humid climate. Brigaud et al. (2009) noted that clay mineral assemblages from the *Stephanoceras humphrieisianum* ammonite biozone of the Paris Basin contain around 20% kaolinite. In modern environments, kaolinite forms in humid tropical climates from intense chemical weathering, which led Brigaud et al. (2009) to argue that there was a change to a

559warmer, more humid, climate during the Early Bajocian. Rausik et al. (2001) and Raucsik
560and Varga (2008) documented clay mineral assemblages dominated by mixed-layer
561illite/smectite from the Lower Bajocian of the Mecsek Mountains of Hungary. Mixed-layer
562illite/smectite clays form in modern environments of warm, arid conditions punctuated by
563short, intense wet seasons; therefore, Raucsik and Varga (2008) argued the climate of the
564Early Bajocian in Europe was monsoon-like. Taken together, Sucheras-Marx et al. (2015)
565and Aguado et al. (2017) argued that these lines of geochemical, mineralogical and
566palaeontological evidence are indicative of a warm, humid climate during the Early Bajocian,
567which enhanced chemical weathering and increased the flux of biolimiting nutrients into the
568European Epicontinental Seaway. Moreover, Philippe et al. (2017) reported the northward
569spread of the low-latitude fossil-wood genus *Brachyoxylon* during the Early Bajocian, which
570may be indicative of a warming and/or an increase in humidity in the mid-latitudes.

571 The alternative scenario is an increase in surface water productivity driven by changes
572in enhanced upwelling and ocean circulation during the Early Bajocian. The Aalenian–
573Bathonian was characterised by significant changes in the tectonic configuration of ocean
574gateways, particularly the gateways which connected the European Epicontinental Seaway
575(EES) to the Tethyan and Panthalassic oceans, and the Boreal Sea. In the north, the Viking
576Corridor, which at times provided a marine connection from the EES to the Boreal Sea
577(Figure 6), was largely blocked during the Aalenian due to magmatic doming in the North
578Sea area (Korte et al., 2015). The Hispanic Corridor to the west, which opened as a marine
579connection between Gondwana and Laurentia during the Early Jurassic (Figure 6), was
580affected by changes in relative plate motion and spreading rate between Africa and North
581America (Labails et al., 2010). These changes drove a widening and deepening of the
582Hispanic Corridor during the Bajocian (Aberhan, 2001; Labails et al., 2010; Dera et al.,
5832014). This was accompanied by increased seafloor spreading in the Alpine Tethys (Bill et

al., 2001). Dera et al. (2014) reported a marked positive shift in neodymium isotopes during the Aalenian–Bathonian.These authors suggested this may have been forced by a stronger influence of radiogenic Tethyan waters in the European Epicontinental Seaway driven by the widening of the Hispanic Corridor and the restriction of the Viking Corridor, which may have promoted enhanced upwelling along the Tethyan margin.

Giraud et al. (2016) argued these changes in ocean circulation drove the turnover in *Watznaueria* through ocean fertilisation. These authors noted that cool seawater temperatures in Europe prevailed during the earliest Bajocian (*H. discites* and *W. laeviuscula* ammonite biozones), consistent with cool, upwelling water masses. Recently, however, Aguado et al. (2017) argued against the upwelling hypothesis. These authors demonstrated that the expansion of *Watznaueria* through the Early Bajocian was concomitant with a decline in the coccolith *Crepidolithus crassus* (Figure 4). This species is characterised by thick, robust coccoliths, which has been interpreted as reflecting a deep-dwelling ecological strategy, as the thick coccoliths act to ballast the coccosphere, while increasing the refractive index with water, and thus allowing a greater amount of light to pass into the cell (Bour et al., 2010; Reggiani et al., 2010; Aguado et al., 2017). As such, Aguado et al. (2017) argued that the decline in the abundance of this deep-dwelling species is inconsistent with nutrients being supplied by upwelling deepwaters, as upwelling would fertilise the photic zone from the bottom up. Instead, Aguado et al. (2017) argued the decline of *C. crassus* is more consistent with an increased supply of continentally derived nutrients which drove a shallowing of the nutricline.

The Early Bajocian eutrophication event has not previously been considered in the context of dinoflagellate ecology and evolution. However, given the diversification and ecological dominance of *Dissiliodinium* in several regions, it is possible that this pattern also reflects a response to elevated nutrient levels, particularly as the acme of this genus in the

Early Bajocian is synchronous with the positive carbon isotope excursion (Figure 13). The palaeoenvironmental distribution and palaeoecological preferences of *Dissiliodinium* may indicate that there was an increase in the supply of nutrients to proximal depositional areas in the mid-palaeolatitudes during the Early Bajocian. In Europe, *Dissiliodinium giganteum* appears to have thrived in palaeoenvironments which were in close proximity to terrestrial discharge (Wiggan et al., 2017). Moreover, this species is typified by a large size (~100 μm). As such, *Dissiliodinium giganteum* may have proliferated under high-nutrient conditions (Wiggan et al. 2017), as large phytoplankton cells require high nutrient levels due to the diffusion limitation of nutrient uptake (Marañón, 2015). Moreover, *Dissiliodinium caddaense* is also typified by a large size (~150 μm; Helby et al., 1987) and is recorded in abundance from the Athol and Cadda formations of north-western and western Australia, which were deposited in nearshore depositional settings (Riding et al., 2010b). These data suggest that, like *Dissiliodinium giganteum*, *Dissiliodinium caddaense* was adapted to high nutrient levels, and could tolerate lowered salinities. Similarly, *Dissiliodinium psilatum* appears to show similar morphological and ecological adaptations. In the Early Bajocian of Argentina, *Dissiliodinium psilatum* dominated assemblages in the marginal/deltaic palaeoenvironments of the Neuquén Basin (Stukins et al., 2013). *Dissiliodinium psilatum* is of comparable morphology to *Dissiliodinium giganteum* and, although slightly smaller, is still relatively large with an average width of 80 μm (Prauss, 1989). The abundance of this species in nearshore and deltaic deposits indicates it could have tolerated reduced salinities, and the proximity to terrigenous discharge and its relatively large size indicate an adaption to high nutrient levels. The proliferation of *Dissiliodinium* in palaeoenvironments close to sources of terrestrial discharge through the Early Bajocian could have been driven by enhanced runoff from increased continental weathering. However, the Early Bajocian was marked by sea level rise in Europe, South America, and Australia (Hallam, 2001; Riding et al., 2010a). Rising sea

level can trap nutrients in marginal marine regions, yet can also introduce nutrient-rich deepwaters into more proximal areas (Arthur and Sageman, 2005). The latter phenomenon seems incompatible with the shallowing nutricline hypothesis of Aguado et al. (2017). As such, further documentation from other Lower Bajocian successions is required to fully resolve of this interval of Early Bajocian palaeoenvironmental change.

3.3.2. The Bajocian radiation of dinoflagellates

To date, few works have attempted to link the Bajocian dinoflagellate radiation to wider Middle Jurassic climatic or oceanographic changes. Feist-Burkhardt and Götz, (2016) suggested that the Bajocian radiation may have been related to opening of the Atlantic Ocean, and changes in ocean circulation. Recently, Wiggan et al. (2017) argued that the pattern of dinoflagellate cyst first appearances in northwest Europe was controlled by transgression. These authors noted that, in the Swabian Basin of southern Germany, the main influx of first appearances was associated with a palaeoenvironmental trend to more distal, offshore conditions. In turn, this palaeoenvironmental trend reflects a major episode of sea level rise during the Bajocian. In Europe, this took the form of the T7 second-order transgression, which lasted from the Late Aalenian to the Early Bathonian (Jacquin et al., 1998). This second-order transgression was comprised of a series of third-order cycles. In the Boreal realm, this transgression had two pulses, the first lasting from the Late Aalenian to Early Bajocian and the second from the latest Early Bajocian to Early Bathonian (Figure 14). In the Tethyan realm, this transgression comprised one pulse, which lasted from the Late Aalenian to Early Bathonian (Figure 14). Maximum transgression of the T7 cycle as a whole corresponds to the maximum flooding surface above Bj5 of the Early Bathonian (*Zigzagiceras zigzag* ammonite biozone) (Jacquin et al., 1998). Hallam (2001) argued that

this transgression was eustatic, as there is evidence of sea level rise in Europe, South America and the Himalayas.

We have compared dinoflagellate cyst stratigraphic data from Europe to sequence stratigraphic records. These data demonstrate a correlation between the increase in dinoflagellate cyst diversity and sea level rise through the Bajocian in Europe (Figure 14). At a zonal level, the large number of appearances recorded from the *Stephanoceras humphriesianum* and *Parkinsonia parkinsoni* ammonite biozones correspond to pronounced third-order transgressive pulses (Figure 14). Appearances of taxa through the Late Aalenian to Early Bathonian could therefore have been brought into the European Epicontinental Seaway by transgressive waters, as indicated by the correlation between large numbers of appearances and transgressive pulses around the *Stephanoceras humphriesianum* and *Parkinsonia parkinsoni* ammonite biozones. Further, in addition to sea-level rise, changes in ocean circulation related to the widening of the Hispanic Corridor and thermal doming in the Viking Corridor may have influenced the pattern of dinoflagellate cyst appearances. Due to the restriction of the Viking Corridor, the main path for the passive dispersal of dinoflagellates would have been to the south, from the northwest Tethys Ocean, and the Hispanic Corridor. The widening of the Hispanic Corridor, and rise in sea level may have driven a rapid influx of gonyaulacacean taxa into Europe during the Bajocian, as intervals of higher sea level can promote the interconnectivity and floral interchange between ocean basins (van de Schootbrugge et al., 2005). As such, enhanced water-mass transfer may have increased floral interchange between the European Epicontinental Seaway, the northwestern Tethys and the Panthalassa Ocean via the Hispanic Corridor. This idea is supported by the occurrence of dinoflagellate cyst taxa, such as *Dissiliodinium psilatum* in both Europe and Argentina (Stukins et al., 2013). Additionally, climatic factors may have also influenced the pattern of appearances in Europe. A brief interval of warming occurred in the *Stephanoceras*

683*humphriesianum* ammonite biozone in Europe. As this zone also corresponds to a major
684influx of gonyaulacacean taxa, warmer seawater temperatures, coupled with sea-level rise,
685could have promoted the spread of taxa from lower palaeolatitude Tethyan regions into the
686European Epicontinental Seaway. Further, the temporal coincidence between climatic
687warming and transgression in the *Stephanoceras humphriesianum* ammonite biozone may
688indicate a common cause via a glacioeustatic control on third-order sea level oscillations.

689 The correlation between the increase in dinoflagellate cyst diversity and sea level rise
690suggests that an increase in the area of epicontinental seas may have directly driven the
691radiation of dinoflagellates, via an increase in available ecospace. Indeed, cyst-forming
692dinoflagellates predominantly inhabit shallow seas as cyst formation is most viable in water
693depth of <200 m (Fensome et al., 1996). However, while the preceding Aalenian was a
694largely regressive interval in Europe, sea level was relatively high globally (Hallam, 2001),
695yet gonyaulacaceans were low in diversity through the Aalenian (Feist-Burkhardt and Pross,
6962010). As such, it does not appear that sea level directly drove the diversification of
697gonyaulacacean dinoflagellates. Furthermore, the Late Pliensbachian was a critical interval of
698dinoflagellate evolution even though it was a time of widespread regression (van de
699Schootbrugge et al., 2005). Similarly, the earliest Jurassic was marked by a widespread
700marine transgression during the Hettangian, yet dinoflagellate cysts were extremely low in
701diversity through this interval (Figure 1; Hallam, 2001; van de Schootbrugge et al., 2013).

702 Dinoflagellates appear to have diversified markedly during cool intervals of the
703Jurassic (Riding and Michoux, 2013). The major pulse of dinoflagellate diversification during
704the Late Pliensbachian was linked to a phase of climatic cooling, which may have increased
705upper-ocean turbulence and enhanced vertical mixing, with an increase in bottom-water
706ventilation, creating conditions ideal for cyst-forming dinoflagellates (van de Schootbrugge et
707al., 2005). Although there may have been a switch to a warmer, more humid climate in during

the Early Bajocian, overall the Aalenian–Bathonian was a time of predominantly cool temperatures in Europe (Korte et al., 2015). This given, it is possible that climatic cooling may have driven dinoflagellate evolution in the Bajocian as it appears to have done during the Late Pliensbachian., However, the Late Pliensbachian was marked by the incursion of high palaeolatitude Boreal taxa into Europe (Bucefalo Palliani and Riding, 1997). In contrast, in the Bajocian, gonyaulacaceans appear to have formed a comparatively minor component of dinoflagellate cyst floras in the high palaeolatitudes (Figure 10; Davies, 1983). Furthermore, there is a ~5 myr gap between the cooling across the Early–Middle Jurassic transition and the diversification of gonyaulacaceans in the Bajocian. Thus, there is no obvious temporal link between cooling climate and the diversification of gonyaulacaceans which suggests that climate did not drive the diversification of gonyaulacacean dinoflagellates during the Middle Jurassic.

Palaeoclimatic and palaeoceanographic changes appear to have controlled the stratigraphic pattern of the Bajocian radiation, as well as influencing palaeoecological patterns seen within dinoflagellate cyst floras. However, these factors do not appear to form the underlying driver of diversification. It is notable, however, that the diversity of dinoflagellates and coccolithophores through the Early–earliest Middle Jurassic broadly mirrors that of benthic and pelagic metazoans, suggesting a coupling between phytoplankton and metazoan evolution (Figure 15). The Mesozoic oceans were characterised by the escalatory evolution of predators and their prey, encapsulated in the concept of the Mesozoic Marine Revolution (Vermeij, 1977; 1987; 2008). Although the concept of the Mesozoic Marine Revolution was originally applied to the Cretaceous, it has since been realised that many of the escalation-related adaptions appeared through the Late Triassic to Early Jurassic (Harper et al., 1998; Vermeij, 2008). Notably, during the Middle Jurassic, important ecological and evolutionary changes occurred in planktivorous metazoans. The trophic guild

of giant suspension feeding fishes first appeared in the Middle Jurassic, and the earliest fossil evidence of this group comes from the Bajocian (Friedman et al., 2010). Within other nekton, the Late Aalenian to Early Bajocian was characterised by the extinction of ammonite families that dominated Early Jurassic ammonite faunas, and their replacement by families that dominated through the Middle Jurassic (Figure 15; Sandoval et al., 2001). Moreover, the Bajocian was marked by the first appearance of heteromorph ammonites (O'Dogherty et al., 2006). Evidence from buccal morphology and fossilised food remains indicates that diet of both planispiral and heteromorph ammonites included mesozooplankton (Jäger and Fraaye, 1997; Kruta et al., 2011). Given the turnover within ammonites, and the appearance of giant suspension feeding fishes, these patterns may indicate that important evolutionary and ecological changes occurred within mesozooplankton communities. Although the fossil record of mesozooplankton is extremely sparse (Selden et al., 2010), it is notable that many of the gonyaulacacean dinoflagellate cyst taxa that appeared through the Bajocian were characterised by thick walls and/or an ornament of sutural crests/spines e.g. *Acanthaulax*, *Aldorfia*, *Ctenidodinium, Gonyaulacysta* and *Meiourogonyaulax* (Figure 11); perhaps such morphological developments served as defensive adaptations (Sarjeant et al., 1987). Additionally, the presence of dinoflagellate cysts in zooplankton fecal pellets suggests that their predation, although, little-considered, may be an important factor in dinoflagellate ecology (Persson, 2000; Montresor et al., 2003). Given the temporal coincidence between phytoplankton diversification and groups of planktivorous cephalopods and fishes, these phenomena may reflect an underlying ecological driver to the Bajocian radiation, with connecting links provided by the mesozooplankton. Viewed in this light, the Middle Jurassic dinoflagellate radiation might represent a largely-overlooked component of the Mesozoic Marine Revolution.

4. Conclusions

Both dinoflagellates and coccolithophores underwent a major radiation through the Middle Triassic to Middle Jurassic. Within this longer-term radiation, important short-term intervals of ecological and evolutionary change can be recognised. The Bajocian was characterised by a major ecological transition in coccoliths and the rapid diversification of gonyaulacacean dinoflagellate cysts. The rise to ecological dominance of the coccolith genus *Watznaueria* during the Early Bajocian appears to have been driven by an increase in nutrient levels. These changes in coccolith floras were paralleled among dinoflagellate cysts by the ecological dominance of the genus *Dissiliodinium* in several mid-palaeolatitude epicratonic basins. An increase in nutrient levels may have been driven by enhanced continental weathering and run off, or changes in ocean circulation. The proliferation of *Dissiliodinium* in proximal palaeoenvironments in the mid-palaeolatitudes during the Early Bajocian may indicate an increase in the supply of continentally derived nutrients, However, the interval of Early Bajocian palaeoenvironmental changes has so far only been documented from Europe and North Africa, which during the Jurassic were located on the northwestern border of the Tethys Ocean. Thus, the global extent and driving mechanisms of these phenomena remains unclear. Future work could focus on documenting the Bajocian carbon isotope record from areas that lay outside of the northwestern Tethys region to accurately assess whether there was a global shift in the carbon cycle during the Early Bajocian. Moreover, the generation of high resolution osmium and strontium isotope records for the Early Bajocian would provide insight into the extent of weathering increase through this interval, in much the same way that these tracers have been employed for examining hydrological cycling during the Toarcian OAE.

The Bajocian was marked by a significant increase in the diversity of gonyaulacacean dinoflagellate cysts, marking the transition to ecological dominance of the Gonyaulacaceae

within cyst-forming dinoflagellates. Thus, the Bajocian represents an important ecological transition for dinoflagellates as well as for coccolithophores. However, this radiation remains poorly documented outside of Europe and Australia. Within Europe, the stratigraphic pattern of the radiation of dinoflagellate cysts appears to have been controlled by sea level, with rising sea level and changes in ocean circulation bringing newly emerging taxa into the European Epicontinental Seaway, by a combination of enhanced water-mass transfer and the spread of marine palaeoenvironments. However, whereas oceanographic changes may have controlled the pattern of dinoflagellate cyst appearances in Europe, they may not have been the underlying driver of diversification. The preceding Aalenian was marked by large volumes of flooded continental area, yet gonyaulacacean diversity was low. Moreover, while gonyaulacaceans appear to have diversified during a 'cool mode' of Jurassic climate, there is a ~5 my gap between rapid cooling through the Early–Middle Jurassic transition and the radiation of gonyaulacaceans during the Bajocian. Furthermore, the Gonyaulacaceae do not appear to have formed a large component of dinoflagellate cyst floras at high palaeolatitudes during the Bajocian. While climatic and oceanographic factors do not appear to have driven the diversification of gonyaulacaceans, the temporal coincidence between the radiation of dinoflagellates, coccolithophores and multiple trophic levels of benthic and pelagic metazoans may point towards an underlying ecological driver. Knoll and Follows (2016) suggested that the radiation of mixotrophic dinoflagellates through the Mesozoic may have provided a bottom-up mechanism to the Mesozoic Marine Revolution, by enhancing primary production. However, many of the morphological features to appear in dinoflagellate cysts through the Bajocian could be interpreted as defensive mechanisms, which might reflect a co-evolutionary escalation between the primary producers and their immediate consumers, the micro- and mesozooplankton. As such, the radiation of dinoflagellates during the Bajocian might represent an important, but largely overlooked component of the Mesozoic Marine

Revolution. Although mesozooplankton represent the most important trophic link between the phytoplankton and higher trophic levels, they have an extremely sparse fossil record. However, ‘zooclasts’, fragments of metazoans, particularly arthropods, are relatively common in Phanerozoic palynomorph assemblages (e.g. Tyson, 1995). Whereas these small carbonaceous fossils have provided a wealth of palaeobiological insight into the Proterozoic–Phanerozoic transition (e.g. Butterfield and Harvey, 2012), including the first record of the copepods (Harvey et al., 2012; Harvey and Pedder, 2013), they are a relatively unexplored source of micropaleontological data for the Mesozoic and Cenozoic. In part, this is because traditional palynological processing methods may destroy these comparatively delicate organic microfossils (Butterfield and Harvey, 2012). Future work may more closely compare the fossil record of phytoplankton with that of pelagic metazoans such as ammonites and fish, and examine Mesozoic rocks for small carbonaceous fossils in order to link the phytoplankton record to that of the zooplankton, which in turn could be linked to higher trophic levels.

Regardless of driving mechanisms, the Bajocian was a critical interval of ecological and evolutionary changes within the dinoflagellates and coccolithophores, and represents an important step within the wider early Mesozoic phytoplankton radiation, which may in turn form part of the wider Mesozoic Marine Revolution.

Acknowledgements

This work has arisen from the PhD project of Nickolas J. Wiggan which was supported by NERC BGS DTG award reference BUFI S246, entitled *The mid Jurassic plankton explosion*. This was funded jointly between the British Geological Survey and the University of Cambridge. James B. Riding publishes with the approval of the Executive Director, British

Geological Survey (NERC). We thank the Review Papers Coordinator, Tim Horscroft, for inviting this contribution. Nick Butterfield (Cambridge) is thanked for discussions and suggestions during NJW's PhD project. We also thank Daniel Mantle and Fabienne Giraud, whose insightful reviews greatly improved the quality of this manuscript.

References

Aberhan, M., 2001. Bivalve palaeobiogeography and the Hispanic Corridor: time of opening and effectiveness of a proto-Atlantic seaway. Palaeogeogr. Palaeoclimatol. Palaeoecol. 165, 375–394.

Aguado, R., O'Dogherty, L., Sandoval, J., 2017. Calcareous nannofossil assemblage turnover in response to the Early Bajocian (Middle Jurassic) palaeoenvironmental changes in the Subbetic Basin. Palaeogeogr. Palaeoclimatol. Palaeoecol. 472, 128–145.

Arthur, M.A., Sageman, B.B., 2005. Sea-level control on source-rock development: perspectives from the Holocene Black Sea, the mid-Cretaceous Western Interior Basin of North America, and the Late Devonian Appalachian Basin.

Bartolini, A., Cecca, F., 1999. 20 My hiatus in the Jurassic of Umbria-Marche Apennines (Italy): carbonate crisis due to eutrophication. Comptes Rendus Académie Sci. - Ser. IIA - Earth Planet. Sci. 329, 587–595.

Bartolini, A., Baumgartner, P., Hunziker, J., 1996. Middle and Late Jurassic carbon stable-isotope stratigraphy and radiolarite sedimentation of the Umbria-Marche Basin (Central Italy). Eclogae Geol. Helvetiae 89, 811–844.

Bartolini, A., Baumgartner, P.O., Guex, J., 1999. Middle and Late Jurassic radiolarian palaeoecology versus carbon-isotope stratigraphy. Palaeogeogr. Palaeoclimatol. Palaeoecol. 145, 43–60.

Bill, M., O'Dogherty, L., Guex, J., Baumgartner, P.O., Masson, H., 2001. Radiolarite ages in Alpine-Mediterranean ophiolites: Constraints on the oceanic spreading and the Tethys-Atlantic connection. Geol. Soc. Am. Bull. 113, 129–143.

Blackburn, T.J., Olsen, P.E., Bowring, S.A., McLean, N.M., Kent, D.V., Puffer, J., McHone, G., Rasbury, E.T., Et-Touhami, M., 2013. Zircon U-Pb geochronology links the end-Triassic extinction with the Central Atlantic Magmatic Province. Science 340, 941–945.

Bodin, S., Hönig, M.R., Krencker, F.-N., Danisch, J., Kabiri, L., 2017. Neritic carbonate crisis during the Early Bajocian: Divergent responses to a global environmental perturbation. Palaeogeogr. Palaeoclimatol. Palaeoecol. 468, 184–199.

Bogus, K., Mertens, K.N., Lauwaert, J., Harding, I.C., Vrielinck, H., Zonneveld, K.A., Versteegh, G.J., 2014. Differences in the chemical composition of organic-walled dinoflagellate resting cysts from phototrophic and heterotrophic dinoflagellates. J. Phycol. 50, 254–266.

Bonis, N., Kürschner, W., Krystyn, L., 2009. A detailed palynological study of the Triassic–Jurassic transition in key sections of the Eiberg Basin (Northern Calcareous Alps, Austria). Rev. Palaeobot. Palynol. 156, 376–400.

Bougeault, C., Pellenard, P., Deconinck, J.-F., Hesselbo, S.P., Dommergues, J.-L., Bruneau, L., Cocquerez, T., Laffont, R., Huret, E., Thibault, N., 2017. Climatic and palaeoceanographic changes during the Pliensbachian (Early Jurassic) inferred from clay mineralogy and stable isotope (C-O) geochemistry (NW Europe). Glob. Planet. Change 149, 139–152.

Bornemann, A., Mutterlose, J., 2006. Size analyses of the coccolith species Biscutum constans and Watznaueria barnesiae from the Late Albian "Niveau Breistroffer"(SE France): taxonomic and palaeoecological implications. Geobios 39, 599–615.

Bour, I., Mattioli, E., Pittet, B., 2007. Nannofacies analysis as a tool to reconstruct paleoenvironmental changes during the Early Toarcian anoxic event. Palaeogeogr. Palaeoclimatol. Palaeoecol. 249, 58–79.

Boussaha, M., Pittet, B., Mattioli, E., Duarte, L., 2014. Spatial characterization of the late Sinemurian (Early Jurassic) palaeoenvironments in the Lusitanian Basin. Palaeogeogr. Palaeoclimatol. Palaeoecol. 409, 320–339.

Bown, P.R., 1987. The structural development of early Mesozoic coccoliths and its evolutionary and taxonomic significance. Abh Geol B-A 39, 33–49.

Bown, P.R., 2005. Calcareous nannoplankton evolution: a tale of two oceans. Micropaleontology 51, 299–308.

Bown, P. R., Young, J. R. 1998. Introduction. In: Bown, P. R. (eds.). Calcareous Nannofossil Biostratigraphy. British Micropalaeontological Society Series. Chapman and Hall/Kluwer Academic Publishers, London, 1–15.

Bown, P. R., Cooper, M. K. E. 1998. Jurassic. *In*: Bown, P. R. (eds). Calcareous Nannofossil Biostratigraphy, British Micropalaeontological Society Series. Chapman and Hall/Kluwer Academic Publishers, London, 86–131.

Bown, P.R., Lees, J.A., Young, J.R., 2004. Calcareous nannoplankton evolution and diversity through time, in: Coccolithophores. Springer, Berlin, Heidelberg, pp. 481–508.

Bralower, T.J., Bown, P.R., Siesser, W.G., 1992. 25. Upper Triassic calcareous nannoplankton biostratigraphy, Wombat Plateau, northwest Australia**.** Proceedings of the Ocean Drilling Program, Scientific Results, 122, 437–451.

Brazier, J.M., Suan, G., Tacail, T., Simon, L., Mattioli, E., Martin, J.E., Balter, V., 2015. Calcium isotope evidence for dramatic increase of continental weathering during the

902Toarcian Oceanic Anoxic Event (Early Jurassic). Earth and Planetary Science Letters 411,
903164-176.

904Brigaud, B., Durlet, C., Deconinck, J.-F., Vincent, B., Pucéat, E., Thierry, J., Trouiller, A.,
9052009. Facies and climate/environmental changes recorded on a carbonate ramp: A
906sedimentological and geochemical approach on Middle Jurassic carbonates (Paris Basin,
907France). Sediment. Geol. 222, 181–206.

908Brittain, J.M., Higgs, K.T. and Riding, J.B. 2010. The palynology of the Pabay Shale
909Formation (Lower Jurassic) of SW Raasay, northern Scotland. Scottish Journal of Geology
91046(1), 67–75.

911Brownlee, C., Taylor, A., 2004. Calcification in coccolithophores: A cellular perspective, in:
912Coccolithophores. Springer, Berlin, Heidelberg, pp. 31–49.

913Bucefalo Palliani, R., Riding, J., 1997. The influence of palaeoenvironmental change on
914dinoflagellate cyst distribution-An example from the Lower and Middle Jurassic of Quercy,
915southwest France. Bull. Cent. Rech. Explor.-Prod. Elf Aquitaine 21, 107–123.

916Bucefalo Palliani, R., Mattioli, E., Riding, J.B., 2002. The response of marine phytoplankton
917and sedimentary organic matter to the early Toarcian (Lower Jurassic) oceanic anoxic event
918in northern England. Mar. Micropaleontol. 46, 223–245.

919Bujak, J.P., Williams, G.L., 1977. Jurassic palynostratigraphy of offshore eastern Canada.
920Dev. Palaeontol. Stratigr. 6, 321–339.

921Butterfield, N., Harvey, T., 2012. Small carbonaceous fossils (SCFs): A new measure of early
922Paleozoic paleobiology. Geology 40, 71–74.

923Calbet, A., 2008. The trophic roles of microzooplankton in marine systems. ICES J. Mar. Sci.
924J. Cons. 65, 325–331.

925Cohen, A.S., Coe, A.L., Harding, S.M., Schwark, L., 2004. Osmium isotope evidence for the
926regulation of atmospheric CO2 by continental weathering. Geology 32, 157–160.

927Conway, B.H., 1990. Palynostratigraphy of the Jurassic succession in the subsurface of Israel.
928Geological Survey of Israel Bulletin 82, 1–39 pp.

929Correia, V.F., Riding, J.B., Fernandes, P., Duarte, L.V., Pereira, Z., 2017a. The palynology of
930the lower and middle Toarcian (Lower Jurassic) in the northern Lusitanian Basin, western
931Portugal. Rev. Palaeobot. Palynol. 237, 75–95.

932Correia, V.F., Riding, J.B., Fernandes, P., Duarte, L.V. and Pereira, Z. 2017b. The
933palynological response to the Toarcian Oceanic Anoxic Event (Early Jurassic) at Peniche,
934Lusitanian Basin, western Portugal. Marine Micropalaeontology, in press.

935Crudeli, D., Young, J.R., Erba, E., Geisen, M., Ziveri, P., de Lange, G.J., Slomp, C.P., 2006.
936Fossil record of holococcoliths and selected hetero-holococcolith associations from the
937Mediterranean (Holocene–late Pleistocene): Evaluation of carbonate diagenesis and
938palaeoecological–palaeocenographic implications. Palaeogeogr. Palaeoclimatol. Palaeoecol.
939237, 191–212.

940Dale, B., 1983. Dinoflagellate resting cysts:'benthic plankton'. In: Surviv. Strateg. Algae.
941Phycological Society of America.

942Davies, E.H., 1983. The dinoflagellate oppel-zonation of the Jurassic-Lower Cretaceous
943sequence in the Sverdrup Basin, arctic Canada. Geological Survey of Canada Bulletin.

944De Kaenel, E., Bergen, J., 1993. New Early and Middle Jurassic coccolith taxa and
945biostratigraphy from the eastern proto-Atlantic (Morocco, Portugal and DSDP Site 547 B).
946Eclogae Geol. Helvetiae 86, 861–907.

de Vargas, C., Aubry, M., Probert, I., Young, J., 2007. Origin and evolution of coccolithophores: from coastal hunters to oceanic farmers. Evol. Prim. Prod. Sea 12, 251–285.

de Vernal, A., Marret, F., 2007. Chapter Nine Organic-Walled Dinoflagellate Cysts: Tracers of Sea-Surface Conditions, in: Vernal, C.H. and A.D. (eds.), Developments in Marine Geology, Proxies in Late Cenozoic Paleoceanography. Elsevier, pp. 371–408.

Delwiche, C., F., 2007. CHAPTER 10 - The Origin and Evolution of Dinoflagellates In: Falkowski, P. G.,Knoll, A.H. (eds.), Evolution of Primary Producers in the Sea. Academic Press, Burlington, pp. 191–205.

Dera, G., Brigaud, B., Monna, F., Laffont, R., Pucéat, E., Deconinck, J.-F., Pellenard, P., Joachimski, M.M., Durlet, C., 2011. Climatic ups and downs in a disturbed Jurassic world. Geology 39, 215–218.

Dera, G., Prunier, J., Smith, P.L., Haggart, J.W., Popov, E., Guzhov, A., Rogov, M., Delsate, D., Thies, D., Cuny, G., Pucéat, E., Charbonnier, G., Bayon, G., 2014. Nd isotope constraints on ocean circulation, paleoclimate, and continental drainage during the Jurassic breakup of Pangea. Gondwana Res. 27, 1599–1615.

Dera, G., Toumoulin, A., De Baets, K., 2016. Diversity and morphological evolution of Jurassic belemnites from South Germany. Palaeogeogr. Palaeoclimatol. Palaeoecol. 457, 80–97.

Ellegaard, M., Lewis, J., Harding, I., 2002. Cyst–Theca Relationship, Life Cycle, and Effects of Temperature and Salinity on the Cyst Morphology of Gonyaulax Baltica Sp. Nov. (dinophyceae) from the Baltic Sea Area1. J. Phycol. 38, 775–789.

969Evitt, W.R., 1985. Sporopollenin Dinoflagellate Cysts: Their Morphology and Interpretation.
970Amer Assn of Stratigraphic, Dallas, Tex.

971

972Falkowski, P.G., Knoll, A.H. (eds). 2007. The evolution of primary producers in the sea.
973Elsevier Academic Press.

974Falkowski, P.G., Katz, M.E., Knoll, A.H., Quigg, A., Raven, J.A., Schofield, O., Taylor,
975F.J.R., 2004a. The Evolution of Modern Eukaryotic Phytoplankton. Science 305,

976Falkowski, P.G., Schofield, O., Katz, M.E., Van de Schootbrugge, B., Knoll, A.H., 2004b.
977Why is the land green and the ocean red?, in: Coccolithophores. Springer, pp. 429–453.

978Feist-Burkhardt, S., 1990. Dinoflagellate cyst assemblages of the Hausen coreholes (Aalenian
979to early Bajocian), southwest Germany. Bull. Cent. Rech. Explor.-Prod. Elf-Aquitaine 14,
980611–633.

981Feist-Burkhardt, S., 2009. Palynology of the Sinemurian/Pliensbachian boundary (Lower
982Jurassic) in the Wutach area, SW Germany: dinoflagellate cyst systematics, biostratigraphy
983and heterotrophic character of Liasidium variabile. Neues Jahrb. Für Geol. Paläontol.-Abh.
984254, 293–313.

985Feist-Burkhardt, S., Götz, A.E., 2016. Ultra-High-Resolution Palynostratigraphy of the Early
986Bajocian Sauzei and Humphriesianum Zones (Middle Jurassic) from Outcrop Sections in the
987Upper Rhine Area, Southwest Germany, in: Stratigraphy & Timescales. Elsevier, pp. 325–
988392.

989Feist-Burkhardt, S., Wille, W., 1992. Jurassic palynology in southwest Germany — state of
990the art. Cah. Micropaléontologie Nouv. Sér. 7, 141–156.

Feist-Burkhardt, S., Monteil, E., 1997. Dinoflagellate cysts from the Bajocian stratotype (Calvados, Normandy, western France). Bull. Cent. Rech. Explor.-Prod. Elf-Aquitaine 21, 31–105.

Feist-Burkhardt, S., Pross, J., 2010. Dinoflagellate cyst biostratigraphy of the Opalinuston Formation (Middle Jurassic) in the Aalenian type area in southwest Germany and north Switzerland. Lethaia 43, 10–31. doi:10.1111/j.1502-3931.2009.00170.x

Fensome, R.A., MacRae, R.A., Moldowan, J.M., Taylor, F.J.R., Williams, G.L., 1996. The early Mesozoic radiation of dinoflagellates. Paleobiology 22, 329–338.

Fensome, R.A., Saldarriaga, J.F., Taylor, "Max" F. J. R., 1999. Dinoflagellate phylogeny revisited: reconciling morphological and molecular based phylogenies. Grana 38, 66–80.

Fensome, R.A., Taylor, F., Norris, G., Sarjeant, W., Wharton, D., Williams, G., 1993. A classification of living and fossil dinoflagellates.

Ferreira, J., Mattioli, E., Van de Schootbrugge, B., 2017. Palaeoenvironmental vs. evolutionary control on size variation of coccoliths across the Lower-Middle Jurassic. Palaeogeogr. Palaeoclimatol. Palaeoecol. 465, 177–192.

Fraguas, Á., Comas-Rengifo, M.J., Gómez, J.J., Goy, A., 2012. The calcareous nannofossil crisis in Northern Spain (Asturias province) linked to the Early Toarcian warming-driven mass extinction. Mar. Micropaleontol. 94, 58–71.

Friedman, M., Shimada, K., Martin, L.D., Everhart, M.J., Liston, J., Maltese, A., Triebold, M., 2010. 100-Million-Year Dynasty of Giant Planktivorous Bony Fishes in the Mesozoic Seas. Science 327, 990–993.

Gardin, S., Krystyn, L., Richoz, S., Bartolini, A., Galbrun, B., 2012. Where and when the earliest coccolithophores? Lethaia 45, 507–523.

Giraud, F., Mattioli, E., López-Otálvaro, G.E., Lécuyer, C., Suchéras-Marx, B., Alméras, Y., Martineau, F., Arnaud-Godet, F., de Kænel, E., 2016. Deciphering processes controlling mid-Jurassic coccolith turnover. Mar. Micropaleontol. 125, 36–50.

Gedl, P., 2008. Organic-walled dinoflagellate cyst stratigraphy of dark Middle Jurassic marine deposits of the Pieniny Klippen Belt, West Carpathians. Stud. Geol. Pol. 131, 7–227.

Guinot, G., Cavin, L., 2015. 'Fish' (Actinopterygii and Elasmobranchii) diversification patterns through deep time: 'Fish' diversification patterns through deep time. Biol. Rev.

Hallam, A., 1976. Stratigraphic distribution and ecology of European Jurassic bivalves. Lethaia 9, 245–259.

Hallam, A., 2001. A review of the broad pattern of Jurassic sea-level changes and their possible causes in the light of current knowledge. Palaeogeogr. Palaeoclimatol. Palaeoecol. 167, 23–37.

Hardenbol, J., Thierry, J., Farley, M.B., Jacquin, T., De Graciansky, P.-C., Vail, P.R. (eds) 1998. Mesozoic and Cenozoic sequence chronostratigraphic framework of European basins. SEPM Special publication no. 60.Harding, I.C., 1986. Archaeopyle variability in Early Cretaceous dinocysts of the partiform gonyaulacoid genus *Druggidium* Habib. J. Micropalaeontology 5, 17–26.

Harper, E.M., Forsythe, G.T., Palmer, T., 1998. Taphonomy and the Mesozoic marine revolution; preservation state masks the importance of boring predators. Palaios 13, 352–360.

Harvey, T.H., Pedder, B.E., 2013. Copepod mandible palynomorphs from the Nolichucky Shale (Cambrian, Tennessee): Implications for the taphonomy and recovery of small carbonaceous fossils. Palaios 28, 278–284.

Harvey, T.H., Vélez, M.I., Butterfield, N.J., 2012. Exceptionally preserved crustaceans from western Canada reveal a cryptic Cambrian radiation. Proc. Natl. Acad. Sci. 109, 1589–1594.

Head,M., 1996. Modern dinoflagellate cysts and their biological affinities. In: Jansonius, J., McGregor, D.C. (Eds.), Palynology: Principles and Applications 3. American Association of Stratigraphic Palynologists, Dallas, Texas, pp. 1197–1248.

Helby, R., Morgan, R., Partridge, A., 1987. A palynological zonation of the Australian Mesozoic. Mem. Assoc. Australas. Palaeontol. 4, 1–94.

Hesselbo, S.P., Gröcke, D.R., Jenkyns, H.C., Bjerrum, C.J., Farrimond, P., Bell, H.S.M., Green, O.R., 2000. Massive dissociation of gas hydrate during a Jurassic oceanic anoxic event. Nature 406, 392–395.

Hesselbo, S.P., Morgans-Bell, H.S., McElwain, J.C., Rees, P.M., Robinson, S.A., Ross, C.E., 2003. Carbon-cycle perturbation in the Middle Jurassic and accompanying changes in the terrestrial paleoenvironment. J. Geol. 111, 259–276.

Hönisch, B., Ridgwell, A., Schmidt, D.N., Thomas, E., Gibbs, S.J., Sluijs, A., Zeebe, R., Kump, L., Martindale, R.C., Greene, S.E., Kiessling, W., Ries, J., Zachos, J.C., Royer, D.L., Barker, S., Marchitto, T.M., Moyer, R., Pelejero, C., Ziveri, P., Foster, G.L., Williams, B., 2012. The Geological Record of Ocean Acidification. Science 335, 1058–1063. doi:10.1126/science.1208277

Hostettler, B., Reisdorf, A.G., Jaeggi, D., Deplazes, G., Bläsi, H., Morard, A., Feist-Burkhardt, S., Waltschew, A., Dietze, V., Menkveld-Gfeller, U., 2017. Litho- and biostratigraphy of the Opalinus Clay and bounding formations in the Mont Terri rock laboratory (Switzerland). Swiss J. Geosci. 110, 23–37.

Ibrahim, M.I., Aboul Ela, N.M., Kholeif, S.E., 2002. Dinoflagellate cyst biostratigraphy of Jurassic-Lower Cretaceous formations of the North Eastern Desert, Egypt. Neues Jahrb. Für Geol. Paläontol.-Abh. 255–319.

Ibrahim, M., Kholeif, S., Al-Saad, H., 2003. Dinoflagellate cyst biostratigraphy and paleoenvironment of the Lower-Middle Jurassic succession of Qatar, Arabian Gulf. Rev. Esp. Micropaleontol. 35, 171–194.

Jacquin, T., Dardeau, G., Durlet, C., de Graciansky, P.-C., Hantzpergue, P., 1998. The North Sea cycle: an overview of 2nd-order transgressive/regressive facies cycles in western Europe. In: de Graciansky, P.C., Hardenbol, J., Jacquin, T., Vail, P.R. (Eds.), Mesozoic and Cenozoic Sequence Stratigraphy of European Basins. SEPM Special Publication. 60, pp. 445–466.

Jäger, M., Fraaye, R., 1997. The diet of the Early Toarcian ammonite Harpoceras falciferum. Palaeontology 40, 557–574.

Janouškovec, J., Gavelis, G.S., Burki, F., Dinh, D., Bachvaroff, T.R., Gornik, S.G., Bright, K.J., Imanian, B., Strom, S.L., Delwiche, C.F., Waller, R.F., Fensome, R.A., Leander, B.S., Rohwer, F.L., Saldarriaga, J.F., 2017. Major transitions in dinoflagellate evolution unveiled by phylotranscriptomics. Proc. Natl. Acad. Sci. 114, 171–180.

Jenkyns, H.C., Jones, C.E., GrÖcke, D.R., Hesselbo, S.P., Parkinson, D.N., 2002. Chemostratigraphy of the Jurassic System: applications, limitations and implications for palaeoceanography. J. Geol. Soc. 159, 351–378.

Kemp, D.B., Coe, A.L., Cohen, A.S., Schwark, L., 2005. Astronomical pacing of methane release in the Early Jurassic period. Nature 437, 396.

Knoll, A.H., Follows, M.J., 2016. A bottom-up perspective on ecosystem change in Mesozoic oceans. Proc. R. Soc. B Biol. Sci. 283, 20161755.

Knoll, A.H., Summons, R.E., Waldbauer, J.R., Zumberge, J.E., 2007. The geological succession of primary producers in the oceans. In: Falkowski, P.G., Knoll, A.H. (eds) Evol. Prim. Prod. Sea 133–163.

Korte, C., Hesselbo, S.P., Ullmann, C.V., Dietl, G., Ruhl, M., Schweigert, G., Thibault, N., 2015. Jurassic climate mode governed by ocean gateway. Nat. Commun. 6, 10015. doi:10.1038/ncomms10015

Kruta, I., Landman, N., Rouget, I., Cecca, F., Tafforeau, P., 2011. The role of ammonites in the Mesozoic marine food web revealed by jaw preservation. Science 331, 70–72.

Kuerschner, W.M., Bonis, N.R., Krystyn, L., 2007. Carbon-isotope stratigraphy and palynostratigraphy of the Triassic–Jurassic transition in the Tiefengraben section—Northern Calcareous Alps (Austria). Palaeogeogr. Palaeoclimatol. Palaeoecol. 244, 257–280.

Labails, C., Olivet, J.-L., Aslanian, D., Roest, W.R., 2010. An alternative early opening scenario for the Central Atlantic Ocean. Earth Planet. Sci. Lett. 297, 355–368.

Lees, J.A., Bown, P.R., Mattioli, E., 2005. Problems with proxies? Cautionary tales of calcareous nannofossil paleoenvironmental indicators. Micropaleontology 51, 333–343.

MacRae, R.A., Fensome, R.A., Williams, G.L., 1996. Fossil dinoflagellate diversity, originations, and extinctions and their significance. Can. J. Bot. 74, 1687–1694.

Mantle, D.J., Riding, J.B., 2012. Palynology of the Middle Jurassic (Bajocian–Bathonian) Wanaea verrucosa dinoflagellate cyst zone of the North West Shelf of Australia. Rev. Palaeobot. Palynol. 180, 41–78.

Marañón, E., 2015. Cell Size as a Key Determinant of Phytoplankton Metabolism and Community Structure. Annu. Rev. Mar. Sci. 7, 241–264. doi:10.1146/annurev-marine-010814-015955

Mattioli, E., Erba, E., 1999. Biostratigraphic syntesis of calcareous nannofossil events in the Tethyan Jurassic. Riv Ital Paleontol Strat. 105, 343–376.

Mattioli, E., Pittet, B., Young, J.R., Bown, P.R., 2004. Biometric analysis of Pliensbachian-Toarcian (Lower Jurassic) coccoliths of the family Biscutaceae: intra- and interspecific variability versus palaeoenvironmental influence. Mar. Micropaleontol., Calcareous nannofossil palaeoecology and palaeocenographic reconstructions 52, 5–27.

Mattioli, E., Pittet, B., Suan, G., Mailliot, S., 2008. Calcareous nannoplankton changes across the early Toarcian oceanic anoxic event in the western Tethys. Paleoceanography 23.

Mattioli, E., Plancq, J., Boussaha, M., Duarte, L., Pittet, B., 2013. Calcareous nannofossil biostratigraphy: new data from the Lower Jurassic of the Lusitanian Basin. Comun. Geológicas.

Mattioli, E., Pittet, B., Petitpierre, L., Mailliot, S., 2009. Dramatic decrease of pelagic carbonate production by nannoplankton across the Early Toarcian anoxic event (T-OAE). Glob. Planet. Change 65, 134–145.

Medlin, L., Sáez, A., Young, J., 2008. A molecular clock for coccolithophores and implications for selectivity of phytoplankton extinctions across the K/T boundary. Mar. Micropaleontol. 67, 69–86.

Moldowan, J.M., Dahl, J., Jacobson, S.R., Huizinga, B.J., Fago, F.J., Shetty, R., Watt, D.S., Peters, K.E., 1996. Chemostratigraphic reconstruction of biofacies: Molecular evidence linking cyst-forming dinoflagellates with pre-Triassic ancestors. Geology 24, 159–162.

Moldowan, J.M., Talyzina, N.M., 1998. Biogeochemical evidence for dinoflagellate ancestors in the Early Cambrian. Science 281, 1168–1170.

Montresor, M., Nuzzo, L., Mazzocchi, M.G., 2003. Viability of dinoflagellate cysts after the passage through the copepod gut. J. Exp. Mar. Biol. Ecol. 287, 209–221.

Morgenroth, P., 1970. Dinoflagellate cysts from the Lias delta of Lühnde/Germany. Neues Jahrb. Für Geol. Paläontol. Abh. 136, 345–359.

O'Dogherty, L., Sandoval, J., Bartolini, A., Bruchez, S., Bill, M., Guex, J., 2006. Carbon–isotope stratigraphy and ammonite faunal turnover for the Middle Jurassic in the Southern Iberian palaeomargin. Palaeogeogr. Palaeoclimatol. Palaeoecol. 239, 311–333.

Ogg, J.G., Ogg, G., Gradstein, F.M., 2016. A Concise Geologic Time Scale: 2016. Elsevier.

Percival, L.M., Ruhl, M., Hesselbo, S.P., Jenkyns, H.C., Mather, T.A., Whiteside, J.H., 2017. Mercury evidence for pulsed volcanism during the end-Triassic mass extinction. Proc. Natl. Acad. Sci. 201705378.

Persson, A., 2000. Possible predation of cysts—a gap in the knowledge of dinoflagellate ecology? J. Plankton Res. 22, 803–809.

Philippe, M., Puijalon, S., Suan, G., Mousset, S., Thévenard, F., Mattioli, E., 2017. The palaeolatitudinal distribution of fossil wood genera as a proxy for European Jurassic terrestrial climate. Palaeogeogr. Palaeoclimatol. Palaeoecol. 466, 373–381.

Piel KM, Evitt WR. 1980. Paratabulation in the Jurassic dinoflagellate genus *Nannoceratopsis* and a comparison with modern taxa. Palynology 4:79–104.

Plancq, J., Mattioli, E., Pittet, B., Baudin, F., Duarte, L.V., Boussaha, M., Grossi, V., 2016. A calcareous nannofossil and organic geochemical study of marine palaeoenvironmental changes across the Sinemurian/Pliensbachian (early Jurassic, ~191Ma) in Portugal. Palaeogeogr. Palaeoclimatol. Palaeoecol. 449, 1–12.

Price, G.D., Baker, S.J., VanDeVelde, J., Clémence, M.-E., 2016. High-resolution carbon cycle and seawater temperature evolution during the Early Jurassic (Sinemurian-Early Pliensbachian). Geochem. Geophys. Geosystems 17, 3917–3928.

Prauss, M., 1989. Dinozysten-Stratigraphie und Palynofazies im Oberen Lias und Dogger von NW-Deutschland. Palaeontogr. Abt. B 1–124.

Raucsik, B., Varga, A., 2008. Climato-environmental controls on clay mineralogy of the Hettangian–Bajocian successions of the Mecsek Mountains, Hungary: An evidence for extreme continental weathering during the early Toarcian oceanic anoxic event. Palaeogeogr. Palaeoclimatol. Palaeoecol. 265, 1–13.

Raucsik, B., A. Demény, I. Borbély Kiss, and G. Szabó. 2001. Monsoon-like climate during the Bajocian. Clay mineralogical and geochemical study on a limestone/marl alternation (Komló Calcareous Marl Formation, Mecsek Mountains, Southern Hungary). Hantkeniana 3, 149–176.

Reggiani, L., Mattioli, E., Pittet, B., Duarte, L., De Oliveira, L.V., Comas-Rengifo, M., 2010. Pliensbachian (Early Jurassic) calcareous nannofossils from the Peniche section (Lusitanian Basin, Portugal): A clue for palaeoenvironmental reconstructions. Mar. Micropaleontol. 75, 1–16.

Richoz, S., van de Schootbrugge, B., Pross, J., Püttmann, W., Quan, T.M., Lindström, S., Heunisch, C., Fiebig, J., Maquil, R., Schouten, S., Hauzenberger, C.A., Wignall, P.B., 2012. Hydrogen sulphide poisoning of shallow seas following the end-Triassic extinction. Nat. Geosci. 5, 662–667.

Riding, J.B., 1984. A palynological investigation of Toarcian to early Aalenian strata from the Blea Wyke area, Ravenscar, North Yorkshire. Proc. Yorks. Geol. Soc. 45, 109–122.

1171Riding, J.B., Thomas, J.E., 1992. Dinoflagellate cysts of the Jurassic System, in: In: Powell,
1172A.J. A Stratigraphic Index of Dinoflagellate Cysts. Chapman and Hall, London, pp. 7–97.

1173Riding, J.B. and Michoux, D. 2013. Further observations on the Jurassic dinoflagellate cyst
1174*Gonyaulacysta dentata* (Raynaud 1978) Lentin & Vozzhennikova 1990 emended Riding
11752012. *Review of Palaeobotany and Palynology* 196, 51–56.

1176Riding, J.B. and Lucas-Clark, J. 2016. The life and scientific work of William R. Evitt
1177(1923–2009). Palynology 40, Supplement 1, 2–131.

1178Riding, J.B., Mantle, D.J., Backhouse, J., 2010a. A review of the chronostratigraphical ages
1179of Middle Triassic to Late Jurassic dinoflagellate cyst biozones of the North West Shelf of
1180Australia. Rev. Palaeobot. Palynol. 162, 543–575.

1181Riding, J.B., Westermann, G.E.G., Darbyshire, D.P.F., 2010b. New evidence for the age of
1182the Athol Formation (Middle Jurassic; Bajocian) in the Tusk-1 and Tusk-2 wells, offshore
1183Carnarvon Basin, Western Australia. Alcheringa Australas. J. Palaeontol. 34, 21–35.

1184Riding, J.B., Leng, M.J., Kender, S., Hesselbo, S.P., Feist-Burkhardt, S., 2013. Isotopic and
1185palynological evidence for a new Early Jurassic environmental perturbation. Palaeogeogr.
1186Palaeoclimatol. Palaeoecol. 374, 16–27.

1187Rost, B., Riebesell, U., 2004. Coccolithophores and the biological pump: responses to
1188environmental changes. Coccolithophores—From Mol. Process. Glob. Impact 76–99.

1189Ruhl, M., Bonis, N.R., Reichart, G.-J., Damsté, J.S.S., Kürschner, W.M., 2011. Atmospheric
1190Carbon Injection Linked to End-Triassic Mass Extinction. Science 333, 430–434.

1191Ruhl, M., Hesselbo, S.P., Hinnov, L., Jenkyns, H.C., Xu, W., Riding, J.B., Storm, M.,
1192Minisini, D., Ullmann, C.V., Leng, M.J., 2016. Astronomical constraints on the duration of

the Early Jurassic Pliensbachian Stage and global climatic fluctuations. Earth Planet. Sci. Lett. 455, 149–165.

Sandoval, J., O'Dogherty, L., Guex, J., 2001. Evolutionary Rates of Jurassic Ammonites in Relation to Sea-Level Fluctuations. PALAIOS 16, 311.

Sarjeant, W.A.S., Lacalli, T., Gaines, G., 1987. The Cysts and Skeletal Elements of Dinoflagellates: Speculations on the Ecological Causes for Their Morphology and Development. Micropaleontology 33, 1–36.

Scotese, C.R., 2014. Atlas of Jurassic Paleogeographic Maps, PALEOMAP Atlas for ArcGIS, volume 4, The Jurassic and Triassic, Maps 32-42, Mollweide Projection, PALEOMAP Project, Evanston, IL.

Schaller, M.F., Wright, J.D., Kent, D.V., 2011. Atmospheric Pco2 Perturbations Associated with the Central Atlantic Magmatic Province. Science 331, 1404–1409.

Schoene, B., Guex, J., Bartolini, A., Schaltegger, U., Blackburn, T.J., 2010. Correlating the end-Triassic mass extinction and flood basalt volcanism at the 100 ka level. Geology 38, 387–390.

Selden, P.A., Huys, R., Stephenson, M.H., Heward, A.P., Taylor, P.N., 2010. Crustaceans from bitumen clast in Carboniferous glacial diamictite extend fossil record of copepods. Nat. Commun. 1, 50.

Simmons, M., BouDagher-Fadel, M., Banner, F., Whittaker, J., 1997. The Jurassic Favusellacea, the earliest Globigerinina, in: The Early Evolutionary History of Planktonic Foraminifera. Springer, pp. 17–51.

Stukins, S., Jolley, D.W., McIlroy, D., Hartley, A.J., 2013. Middle Jurassic vegetation dynamics from allochthonous palynological assemblages: An example from a marginal

marine depositional setting; Lajas Formation, Neuquén Basin, Argentina. Palaeogeogr. Palaeoclimatol. Palaeoecol. 392, 117–127.

Suan, G., Pittet, B., Bour, I., Mattioli, E., Duarte, L.V., Mailliot, S., 2008. Duration of the Early Toarcian carbon isotope excursion deduced from spectral analysis: consequence for its possible causes. Earth Planet. Sci. Lett. 267, 666–679.

Suan, G., Mattioli, E., Pittet, B., Lécuyer, C., Suchéras-Marx, B., Duarte, L.V., Philippe, M., Reggiani, L., Martineau, F., 2010. Secular environmental precursors to Early Toarcian (Jurassic) extreme climate changes. Earth Planet. Sci. Lett. 290, 448–458.

Suan, G., Nikitenko, B.L., Rogov, M.A., Baudin, F., Spangenberg, J.E., Knyazev, V.G., Glinskikh, L.A., Goryacheva, A.A., Adatte, T., Riding, J.B., Föllmi, K.B., Pittet, B., Mattioli, E., Lécuyer, C., 2011. Polar record of Early Jurassic massive carbon injection. Earth Planet. Sci. Lett. 312, 102–113.

Suchéras-Marx, B., Guihou, A., Giraud, F., Lécuyer, C., Allemand, P., Pittet, B., Mattioli, E., 2012. Impact of the Middle Jurassic diversification of Watznaueria (coccolith-bearing algae) on the carbon cycle and $\delta^{13}C$ of bulk marine carbonates. Glob. Planet. Change 86–87, 92–100.

Suchéras-Marx, B., Giraud, F., Fernandez, V., Pittet, B., Lecuyer, C., Olivero, D., Mattioli, E., 2013. Duration of the Early Bajocian and the associated δ 13 C positive excursion based on cyclostratigraphy. J. Geol. Soc. 170, 107–118.

Suchéras-Marx, B., Mattioli, E., Giraud, F., Escarguel, G., 2015. Paleoenvironmental and paleobiological origins of coccolithophorid genus Watznaueria emergence during the late Aalenian–early Bajocian. Paleobiology 41, 415–435.

Taylor, P.D., Ernst, A., 2008. Bryozoans in transition: The depauperate and patchy Jurassic biota. Palaeogeogr. Palaeoclimatol. Palaeoecol., Jurassic Marine Palaeobiology 263, 9–23.

Tiraboschi, D., Erba, E., 2010. Calcareous nannofossil biostratigraphy (Upper Bajocian–Lower Bathonian) of the Ravin du Bès section (Bas Auran, Subalpine Basin, SE France): evolutionary trends of Watznaueria barnesiae and new findings of "Rucinolithus" morphotypes. Geobios 43, 59–76.

Tremolada, F., Van de Schootbrugge, B., Erba, E., 2005. Early Jurassic schizosphaerellid crisis in Cantabria, Spain: implications for calcification rates and phytoplankton evolution across the Toarcian oceanic anoxic event. Paleoceanography 20.

Tyson, R.V., 1995. Abundance of organic matter in sediments: TOC, hydrodynamic equivalence, dilution and flux effects, in: Sedimentary Organic Matter. Springer, pp. 81–118.

van de Schootbrugge, B., Bailey, T.R., Rosenthal, Y., Katz, M.E., Wright, J.D., Miller, K.G., Feist-Burkhardt, S., Falkowski, P.G., 2005. Early Jurassic climate change and the radiation of organic-walled phytoplankton in the Tethys Ocean. Paleobiology 31, 73–97.

van de Schootbrugge, B., Tremolada, F., Rosenthal, Y., Bailey, T.R., Feist-Burkhardt, S., Brinkhuis, H., Pross, J., Kent, D.V., Falkowski, P.G., 2007. End-Triassic calcification crisis and blooms of organic-walled 'disaster species.' Palaeogeogr. Palaeoclimatol. Palaeoecol., Triassic-Jurassic Boundary events: problems, progress, possibilities 244, 126–141.

van de Schootbrugge, B., Bachan, A., Suan, G., Richoz, S., Payne, J.L., 2013. Microbes, mud and methane: cause and consequence of recurrent Early Jurassic anoxia following the end-Triassic mass extinction. Palaeontology 56, 685–709. doi:10.1111/pala.12034

Vermeij, G.J., 1977. The Mesozoic marine revolution: evidence from snails, predators and grazers. Paleobiology 3, 245–258.

Vermeij, G.J., 1987. Evolution and escalation: an ecological history of life. Princeton University Press.

Vermeij, G.J., 2008. Escalation and its role in Jurassic biotic history. Palaeogeogr. Palaeoclimatol. Palaeoecol., Jurassic Marine Palaeobiology 263, 3–8.

Versteegh, G.J.M., Blokker, P., Wood, G.D., Collinson, M.E., Sinninghe Damsté, J.S., de Leeuw, J.W., 2004. An example of oxidative polymerization of unsaturated fatty acids as a preservation pathway for dinoflagellate organic matter. Org. Geochem., Selected papers from the sixteenth international symposium on environmental biogeochemistry 35, 1129–1139.

Wall, D., 1965. Microplankton, Pollen, and Spores from the Lower Jurassic in Britain. Micropaleontology 11, 151–190. doi:10.2307/1484516

Wiggan, N.J., Riding, J.B., Franz, M., 2017. Resolving the Middle Jurassic dinoflagellate radiation: The palynology of the Bajocian of Swabia, southwest Germany. Rev. Palaeobot. Palynol. 238, 55–87.

Wood, S.E.L., Riding, J.B., Fensome, R.A. and Williams, G.L. 2016. A review of the *Sentusidinium* complex of dinoflagellate cysts. Review of Palaeobotany and Palynology 234, 61–93.

Xu, W., Ruhl, M., Jenkyns, H.C., Hesselbo, S.P., Riding, J.B., Selby, D., Naafs, B.D.A., Weijers, J.W., Pancost, R.D., Tegelaar, E.W., 2017. Carbon sequestration in an expanded lake system during the Toarcian oceanic anoxic event. Nat. Geosci. 10, 129–134.

Figure Captions:

Figure 1. Dinoflagellate cyst and coccolith species richness throughout the Middle Triassic to Jurassic modified from Wiggan et al. (2017, fig. 1). The timescale is based on Gradstein et al. (2012). The family-level spindle plot for dinoflagellate cysts clearly shows the significant increase in diversity of the family Gonyaulaceae during the Bajocian. Key bioevents for coccolithophores are indicated, notably the evolution of placolith coccoliths in the Pliensbachian, and the ecological transition to the dominance of *Watznaueria* in the Bajocian. The dinoflagellate cyst species richness is taken from MacRae et al. (1996), the spindle plot data is after Fensome et al. (1996) and the coccolith species richness is after Bown et al. (2004).

Figure 2. Coccolith abundances and $\delta^{13}C$ for the Late Aalenian to Early Bajocian marls of the Subalpine Basin, France. Note that the increase in the absolute and relative abundances of *Watznaueria* through the Early Bajocian parallels the increase in $\delta^{13}C$. There is also a shift in the absolute abundance of coccoliths of around an order of magnitude, from 10^7–10^8 coccoliths per g in the latest Aalenian to earliest Bajocian, to 10^8–10^9 coccoliths per g in the latest Early Bajocian. Modified from Suchéras-Marx et al. (2015, fig. 4).

Figure 3. Synthetic relative abundances of three important morphogroups of the coccolithophore genus *Watznaueria* through the Late Aalenian to Early Bathonian from Giraud et al. (2016). Note that at the Aalenian–Bajocian transition, there is an increase in taxa defined by a central cross area. This is then followed by an increase in the abundances of taxa without a central area, and those with a central bar. This increase in the abundance of

Watznaueria is concomitant with a long-term decline in the abundance of *Lotharingius*. Abundances after Giraud et al. (2016, fig. 8) and coccolith morphologies after Giraud et al. (2016, fig. 2). Ammonite biozone abbrevations: L.m. = *Ludwigia murchisonae*; G.c = *Graphoceras concavum*; H.d. = *Hyperlioceras discites*; W.l. = *Witchellia laeviuscula*; S.p. = *Sonninia propinquans*; S.h. = *Stephanoceras humphriesianum*; S.n. = *Strenoceras niortense*; G.g. = *Garantiana garantiana*; P.p. = *Parkinsonia parkinsoni*; Z.z. = *Zigzagiceras zigzag*; P.a. = *Procerites aurigerus*; P.p. = *Procerites progracilis*, T.s. = *Tulites subcontractus*; M.m. = *Morrisiceras morrisi*; H.r. = *Hecticoceras retrocostatum*; C.d. = *Clydoniceras discus*.

Figure 4. Coccolith abundances and $\delta^{13}C$ for the Late Aalenian to Early Bajocian of the southern Iberian palaeomargin, Spain modified after Aguado et al. (2017, fig. 2). Note the increase in the absolute abundance of *Watznaueria* which parallels an increase in $\delta^{13}C$. The increase in the abundance of *Watznaueria* is concomitant with the decline in the abundance of *Crepidolithus crassus*, a coccolith which has been interpreted as inhabiting the deep photic zone.

Figure 5. Absolute abundances of *Dissiliodinium giganteum* through the Early Bajocian of borehole B404-2 in southern Germany based on raw data from Wiggan et al. (2017). *Dissiliodinium giganteum* forms a significant proportion (>80%) of the dinoflagellate cyst assemblages throughout the *Witchellia laeviuscula* ammonite biozone, but declined in abundance towards the latest Early Bajocian (*Stephanoceras humphriesianum* ammonite biozone). Two specimens of *Dissiliodinium giganteum* are shown to the right. The upper specimen is in apical view; the loss of multiple precingular plates during archaeopyle formation can be observed. The lower image is in dorsoventral view.

Figure 6. Bajocian palaeogeography and the abundance distribution of the dinoflagellate cyst genus *Dissiliodinium* during the Early Bajocian. *Dissiliodinium giganteum* was abundant in basins which bordered the northwest Tethys Ocean on the edge of the European Epicontinental Seaway (EES) (i.e. southern Germany, Poland/northern Slovakia and Switzerland). *Dissiliodinium caddaense* was abundant in the Early Bajocian of Australia, and *Dissiliodinium psilatum* was prominent in the Early Bajocian of the Neuquén Basin, Argentina. The abundances of *Dissiliodinium giganteum* are based on raw data in Wiggan et al. (2017) and unpublished information. The abundances of *Dissiliodinium caddaense* are based on raw data from Riding et al. (2010b). The abundances of *Dissiliodinium psilatum* are based on raw data from Stukins et al. (2013). The palaeogeography is modified after Scotese (2014).

Figure 7. The stratigraphic ranges of dinoflagellate cysts from the latest Aalenian to the earliest Bathonian of northwest Europe, based on data from Wiggan et al. (2017). Note that there is a relatively continuous pattern of first appearances, but there are numerous first appearances in the *Stephanoceras humphriesianum* ammonite biozone (*S.h.*). The grey fill denotes dinoflagellate cyst taxa which belong to the family Gonyaulacaceae. The ammonite biozone abbreviations are as per Figure 3.

Figure 8. The stratigraphic ranges of dinoflagellate cysts from the latest Early Bajocian to the earliest Bathonian of northwest Europe based on data from Wiggan et al. (2017). Note the large number of first appearances in the *Stephanoceras humphriesianum* (*S.h.*) and *Parkinsonia parkinsoni* (*P.p.*) ammonite biozones. First appearances from the *Stephanoceras*

humphriesianum zone onwards are dominated by gonyaulacacean taxa (grey fill). The ammonite biozone abbreviations are as per Figure 3. F-B & M (1997) = Feist-Burkhardt and Monteil (1997).

Figure 9. A dinoflagellate cyst species richness curve from the latest Aalenian to the earliest Bathonian based on stratigraphic data from Wiggan et al. (2017). Note how the increase in diversity is primarily driven by a rise in the number of gonyaulacacean taxa (grey). There are also pronounced increases in species richness in the *Stephanoceras humphriesianum* and *Parkinsonia parkinsoni* ammonite biozones.

Figure 10. Percentage of gonyaulacacean dinoflagellate cyst taxa for the Bajocian and Bathonian low-, mid-, and high-palaeolatitude regions. It is clear that gonyaulacaceans formed a larger proportion of the dinoflagellate cyst floras in the low- to mid-palaeolatitudes than the high-palaeolatitudes during the Middle Jurassic. The graph is based on raw data from: Bujak and Williams (1977) (eastern Canada); Davies (1983) (Arctic Canada); Conway (1990) (Israel); Ibrahim et al. (2003) (Egypt); Mantle and Riding (2012) (Australia); Stukins et al. (2013) (Argentina); and Wiggan et al. (2017) (northwest Europe).

Figure 11. The diversity and disparity of gonyaulacacean dinoflagellate cyst taxa from the Aalenian and Bajocian. Note how, in the Aalenian, the Gonyaulaceae were represented by four genera, all of relatively simple morphology and largely nontabulate (i.e. not exhibiting tabulation). *Batiacasphaera, Kallosphaeridium* and *Sentusidinium* all have apical archaeopyles, whilst *Dissiliodinium* has a multiplate precingular archaeopyle. In the Bajocian, the earliest fully tabulate sexiform gonyaulacaceans appeared, represented by

Durotrigia daveyi. Other Bajocian developments included thick, ornamented cyst walls (e.g. *Acanthaulax, Aldorfia*), and prominent sutural crests (e.g. *Ctendidodinium, Meiourogonyaulax*). Moreover, the single-plate precingular archaeopyle appeared in genera such as *Acanthaulax, Aldorfia* and *Gonyaulacysta*, which has remained a prominent excystment strategy in gonyaulacaceans to the present day. Similarly, the epicystal archaeopyle type appeared in genera such *Ctendodinium, Korystocysta* and *Wanaea*. Note that *Meiourogonyaulax* and *Rhynchodiniopsis* are each represented by two photographs of the same specimen taken at different focal depths. The left image of *Meiourogonyaulax* was taken at the mid-focus, and the right taken at low focus, the latter shows the apical archaeopyle. The left image of *Rhynchodiniopsis* was taken at the high focus and right at low focus, the latter shows the one-plate precingular archaeopyle.

Figure 12. Seawater palaeotemperatures based on the $\delta^{18}O$ of brachiopod and mollusc carbonate for the Early and Middle Jurassic (Pliensbachian–Bajocian) of Europe, simplified and redrawn after Korte et al. (2015, fig. 2). It is clear that a shift to cooler palaeotemperatures of ~10°C occurred through the Early–Middle Jurassic transition. The pronounced warming pulse in the latest Early Bajocian corresponds to the *Stephanoceras humphriesianum* ammonite biozone.

Figure 13. Carbon isotope and abundances of *Dissiliodinium giganteum* for the Late Aalenian to Early Bathonian.The central column depicts a composite $\delta^{13}C$ curve for bulk carbonate from the Late Aalenian to Early Bathonian of Europe based on raw data from O'Dogherty et al. (2006) and Suchéras-Marx et al. (2013). Note the ~1.5 ‰ shift to more positive $\delta^{13}C$ values in the Early Bajocian. The right hand panel illustrates the abundance of the dinoflagellate cyst *Dissiliodinium giganteum* in the Swabian Basin of southern Germany

taken from Wiggan et al. (2017) and supplemented by unpublished data. Note that the acme of *Dissiliodinium giganteum* is coeval with the positive shift in $\delta^{13}C$.

Figure 14. A comparison of sequence stratigraphy, sea level and dinoflagellate cyst species richness and appearances per zone through the Late Aalenian to Early Bathonian. The Bajocian was characterised by the T7 second-order transgression, which lasted from the latest Aalenian *Graphoceras concavum* ammonite biozone to the earliest Bathonian *Zigzagiceras zigzag* ammonite biozone. Maximum transgression is represented by the maximum flooding surface above third-order boundary Bj5 (the dotted line), which corresponds to a major flooding event through many European basins (Jacquin et al., 1998; Hallam, 2001). Note how the increase in species richness roughly parallels the rise in sea level. The large number of first appearances from the *Stephanoceras humphriesianum* (*S.h.*) and *Parkinsonia parkinsoni* (*P.p.*) ammonite biozones correspond to the pronounced third-order transgressive pulses Bj2 and Bj5.The ammonite biozone abbreviations are as per Figure 3. The sea level cycles are as per Figure 12. The stratigraphy and sea level were generated using Timescale Creator v.6.4.2; the mean sea level curve is based on midpoint of coastal onlap charts of Hardenbol et al. (1998).

Figure 15. Species richness curves and the appearances of key marine fossil groups for the Early and Middle Jurassic. It is clear that the species richness curves of coccoliths and dinoflagellate cysts closely parallel those from higher trophic levels. All groups are plotted per stage/substage, except for the ammonites which are plotted per ammonite biozone. Coccolith and dinoflagellate cyst species richness data were taken from Figure 1. The planktonic foraminifera data are from Simmons et al. (1997). The bivalve data are those of

1428Hallam (1976). Data from bryozoans are from Taylor and Ernst (2008). The elasmobranch
1429diversity is after Guinot and Cavin (2015). The belemnite information is after Dera et al.
1430(2016), and the ammonite data are from Sandoval et al. (2001) and O'Dogherty et al. (2006).
1431The range of giant suspension feeders is after Friedman et al. (2010).
1432