

1 **Oxygen and placental development; parallels and differences with tumour biology**

2

3 <sup>1</sup>Graham J Burton, <sup>2</sup>Eric Jauniaux and <sup>1</sup>Andrew J Murray

4

5 <sup>1</sup>Centre for Trophoblast Research, Department of Physiology, Development and

6 Neuroscience, University of Cambridge, Cambridge, UK

7 <sup>2</sup>Department of Obstetrics and Gynaecology, University College London Hospitals and

8 UCL Institute for Women's Health, University College London (UCL), London, UK.

9

10 Short title: Oxygen and placental development

11 Key words: oxygen, oxidative stress, metabolism, glycolysis

12

13

14

15 Address for correspondence:

16 Professor G.J. Burton,

17 Physiological Laboratory,

18 Downing Street,

19 Cambridge, CB2 3EG

20 UK

21 Email: [gjb2@cam.ac.uk](mailto:gjb2@cam.ac.uk)

22 Tel: +44 1223333856

23

24 **Abstract**

25 Human placentation involves the invasion of the conceptus into the wall of the uterus,  
26 and establishment of a blood supply from the maternal spiral arteries. The placenta has  
27 therefore been likened to a malignant tumour, albeit a highly regulated one. Oxygen  
28 plays an important role in controlling both placental development and tumour  
29 behaviour. In the placenta, early development takes place in a physiological low oxygen  
30 environment, which undergoes a transition with onset of the full maternal arterial  
31 circulation towards the end of the first trimester. By comparison, in tumours there is  
32 often a progressive hypoxia as the mass outgrows its blood supply. Both early placental  
33 tissues and tumour cells show high rates of proliferation, and the energy required to  
34 support these comes principally from glycolysis. Glycolysis is maintained in placental  
35 tissues by reoxidation of pyridine nucleotides through the polyol pathways, whereas in  
36 tumours there is fermentation to lactate, Warburg metabolism. In both cases, the  
37 reliance on glycolysis rather than oxidative phosphorylation preserves carbon skeletons  
38 that can be utilised in the synthesis of nucleotides, cell membranes and organelles, and  
39 that would otherwise be excreted as carbon dioxide. In the placenta, this reliance may  
40 also protect the embryo from free radical-mediated teratogenesis. Local oxygen  
41 gradients within both sets of tissues may influence the cell behaviour. In particular, they  
42 may induce an epithelial-mesenchymal transition, promoting extravillous trophoblast  
43 invasion in the placenta and metastasis in a tumour. Further investigations into the two  
44 scenarios may provide new insights of benefit to these contrasting, but similar, fields of  
45 cellular biology.

46

47



49

## 50 **Introduction**

51 Placental development displays many of the same growth characteristics as are seen in  
52 malignant tumour progression, such as a high proliferative rate, invasion into the host  
53 tissue, and immunological modulation (1). There are parallels too in terms of  
54 oxygenation and tissue metabolism, but also significant divergences. Here, we review  
55 the major similarities and differences.

56

57

## 58 **The first trimester placental environment**

59 Fertilization and early development of the conceptus occur in the Fallopian tube,  
60 supported by the oviductal secretions. In the human, measurements performed during  
61 the non-pregnant cycle indicate an oxygen tension of 15-19 mmHg (2, 3), and it is likely  
62 that similar conditions prevail during early pregnancy. Data from the mouse show that  
63 oxygen consumption by the early conceptus is low, at approximately 4  $\mu\text{l}/\text{mg}$  dry weight  
64 per hour, prior to implantation, although it peaks transiently at the time of blastocyst  
65 formation due to the higher energy demands associated with ionic pumping and protein  
66 synthesis (4). This low level of oxygen consumption has been coined 'quiet metabolism'  
67 (5), and is considered to be beneficial as it limits the production of potentially harmful  
68 reactive oxygen species. These species, and their non-radical intermediates, may cause  
69 damage to diverse biomolecules, including lipid peroxidation, protein carbonylation and  
70 DNA strand breaks. Indeed, more active 'noisy' metabolism is associated with higher  
71 levels of DNA damage, and with poorer outcomes in assisted reproductive technologies  
72 (6).

73

74 Early placental development can be seen as a continuation of this 'quiet metabolism', for  
75 the oxygen concentration within the intervillous space and the embryonic  
76 compartments remains at approximately 20 mmHg during most of the first trimester (7,  
77 8). Following implantation, the conceptus lies within the superficial endometrium, and  
78 as the trophoblast mantle expands it erodes into neighbouring capillaries and into the  
79 endometrial glands. Maternal arterial inflow into the placenta only occurs towards the  
80 end of the first trimester, as initially the endovascular trophoblast invasion that occurs  
81 as part of remodelling of the spiral arteries is sufficiently voluminous to occlude the  
82 mouths of most of the vessels (9, 10). A network of narrow intercellular spaces exists  
83 between the endovascular trophoblast cells, however, enabling maternal plasma to pass  
84 into the placenta at a slow rate. Consequently, there is a continual supply of oxygen,  
85 albeit at a low partial pressure and content as it is carried principally in solution in the  
86 absence of maternal erythrocytes.

87

88 The distribution of this oxygen to the deeper placental and fetal tissues must initially  
89 occur by simple diffusion, for the fetal heart does not start beating until the 5th week of  
90 pregnancy, and an effective circulation through the placental villi is only achieved  
91 towards the end of the first trimester. Diffusion is facilitated by the large surface area  
92 provided by the villous morphology of the placenta, and presence of fluid-filled stromal  
93 channels within the villi that communicate with the extra-embryonic coelom (11). The  
94 oxygen within the exocoelomic fluid is able reach the deeper tissues within the embryo  
95 as the intra- and extra-embryonic coeloms are in free communication before the  
96 anterior body folds fuse at around 6 weeks post-fertilisation. When the fetal-placental  
97 circulation is established, oxygen transport is achieved during the first three months of  
98 pregnancy by high-affinity embryonic haemoglobin (Hb) located inside red cells which

99 are mainly nucleated. The oxygen binding characteristics of embryonic Hb and the high  
100 viscosity of circulating blood containing a high proportion of nucleated red cells  
101 contribute to limiting oxygen transfer to the fetal tissues (12-14).

102

103 By comparison, although the oxygen tension in many tumours is low, and indeed lower  
104 than that inside the early placenta, the situation has a very different ontology. The  
105 pattern in tumour masses is one of increasing hypoxia, compared to the steady state  
106 seen within the early placenta. In tumours, the initiating growth normally occurs at  
107 ordinary tissue oxygen levels, but with expansion the tumour gradually outstrips its  
108 blood supply. In solid tumours the opportunity for diffusion is limited, and as a result  
109 the central core becomes increasingly hypoxic (15). Although angiogenesis is stimulated  
110 through the release of VEGF, the degree of hypoxia may be sufficient to induce necrosis  
111 in the core, an event never seen in first trimester placental tissues or in fetal  
112 development. In fact, variation in blood flow distribution to the periphery of the early  
113 placenta leads to a high level of oxygen exposure inducing apoptosis and degeneration  
114 of two-thirds of the original placental mass, a process which is pivotal for the formation  
115 of the membranes.

116

### 117 **Early placental metabolism**

118 Early placental tissues display a high proliferative rate, as do tumour cells, although the  
119 drivers are different. In the placenta, proliferation is thought to be stimulated  
120 exogenously by mitogens secreted by the endometrial glands (16). Both epidermal  
121 growth factor and the insulin-like growth factors promote proliferation of the  
122 cytotrophoblast cells when applied to first trimester villous explants (17, 18). These  
123 mitogens are presumably transported through the syncytiotrophoblast by the same

124 endocytotic/exocytotic pathways that lead to the accumulation of other gland products,  
125 such as glycodeilin, in the amniotic fluid (19). By contrast, in tumour cells the drive for  
126 proliferation arises as the result of endogenous mutations within growth promoting  
127 pathways. However, unlike in a tumour, the placental tissues display no evidence of  
128 hypoxic stress. Hypoxia cannot be defined simply by the prevailing partial pressure that  
129 cells are exposed to, but rather by whether the oxygen supply is sufficient to meet the  
130 metabolic requirements of the cells. Hence, it is notable that the ATP/ADP ratio in  
131 placental tissues is the same during the first trimester as it is later in the second  
132 trimester and at term (20). Furthermore, there is no stabilisation of either hypoxia  
133 inducible factors (HIF-1 and HIF-2) in villi removed by a chorionic villous sampling  
134 technique, which avoids any confounding stress induced by exposure to maternal blood  
135 as occurs during curettage (20). These differences with the tumour situation most likely  
136 reflect the replenishment of oxygen through the perfusion of the intervillous chamber  
137 with maternal plasma, and also the different ontological progressions. In addition, the  
138 placental tissues are provided with a rich source of glucose for glycolysis by the  
139 endometrial glands, along with lipid and proteinaceous substrates (21).

140

141 The exocoelomic fluid is in free communication with the placenta tissues, and so its  
142 metabolic profile predominantly reflects placental metabolism. Analysis of the fluid  
143 indicates evidence of limited anaerobic metabolism, in that the pH of the fluid at 7-10  
144 weeks of gestation is approximately 7.17, with a base excess of -8.9 mmol/l (22). The  
145 concentration of lactate is, however, not excessively high (0.6 mmol/l). In part, this may  
146 be due to metabolism of lactate by the fetus, but it also reflects the reliance of the  
147 placenta on phylogenetically old carbohydrate metabolic pathways involving the  
148 formation of polyols (23).

149

150 **The importance of glycolysis**

151 One of the most striking similarities between the early placenta and tumours is their  
152 reliance on glycolysis for energy production, although the pathways involved in enabling  
153 this are quite different. In the case of the placenta, glycolysis is closely interlinked with  
154 the polyol and pentose-phosphate pathways. Conversion of glucose to pyruvate  
155 generates two molecules of ATP, and requires a supply of NAD<sup>+</sup>. Under full aerobic  
156 conditions that NAD<sup>+</sup> is normally regenerated via the tricarboxylic acid (TCA) cycle,  
157 whereas in adult tissues under anaerobic conditions NAD<sup>+</sup> is regenerated by  
158 fermentation of pyruvate to lactate (Figure 1). The polyol pathways provide an  
159 alternative mechanism for maintaining the oxidation-reduction balance of pyridine  
160 nucleotides. Conversion of ribose 5-phosphate created from glucose in the pentose-  
161 phosphate pathway to ribitol regenerates NAD<sup>+</sup>. Similarly, formation of erythritol and  
162 sorbitol regenerates NADP<sup>+</sup>. The concentrations of these polyols are much higher in the  
163 coelomic fluid than in maternal serum during early pregnancy (23).

164

165 By contrast, in tumours fermentation to lactate appears to be the principal method for  
166 regeneration of NAD<sup>+</sup>, even under conditions of adequate oxygenation. This process is  
167 therefore referred to as aerobic glycolysis, or eponymously as the Warburg effect. In  
168 hypoxic cells and tissues, such as the tumour, glycolysis is directly stimulated following  
169 HIF-1 stabilisation, with the upregulation of most, if not all, glycolytic enzymes (24).  
170 Notably, aerobic glycolysis is also specifically promoted, and mitochondrial pyruvate  
171 oxidation bypassed, via inhibition of pyruvate dehydrogenase (PDH) activity (Figure 1).  
172 HIF-1 dependent upregulation of PDH kinase 1 (PDK-1) (25, 26) leads to the  
173 phosphorylation of the E1 subunit of PDH, and thus its inhibition. Under such conditions,



174 pyruvate is therefore not converted into acetyl-CoA, and the TCA cycle cannot be fuelled,  
175 leading to a fall in mitochondrial oxygen consumption (26), which promotes survival in  
176 the face of hypoxia. Hypoxic cells instead accumulate pyruvate, some of which is  
177 converted to lactate under the action of lactate dehydrogenase (LDH), another HIF-1  
178 regulated enzyme (24), and lactate is in turn transported out of the cell (27). The build-  
179 up of pyruvate also favours transformation of fructose-6-phosphate to D-ribose-5-  
180 phosphate, promoting the synthesis of nucleic acids to support cell proliferation (Figure  
181 1). At present it is unclear whether glycolysis is promoted by mitochondrial inhibition  
182 in a similar fashion in the case of the placenta, though in the apparent absence of HIF-1  
183 stabilisation this would seem unlikely. Instead, placental glycolysis may possibly be  
184 promoted early in pregnancy as a necessary means of supporting ATP-synthesis in the  
185 absence of significant mitochondrial activity. The time-course of changes in placental  
186 mitochondrial density has not yet been established; however, the initiation of significant  
187 mitochondrial biogenesis may only coincide with the rise in oxygenation towards the  
188 end of the first trimester.

189

190 It might be supposed that in both situations metabolism is relatively inefficient, and  
191 does not take advantage of the higher yield of ATP that can be gained through oxidative  
192 phosphorylation. However, unlike differentiated cells in adult tissues the rapidly  
193 proliferating cells of the placenta and a tumour have additional requirements. There is a  
194 need for carbon skeletons that can be incorporated into nucleotides, amino acids and  
195 sterols that support synthesis of DNA, cell and organelle membranes, and proteins.  
196 Instead of breaking glucose down completely and excreting the carbon atoms as carbon  
197 dioxide, maintaining the carbon skeletons as lactate or through the pentose-phosphate  
198 pathways allows them to be incorporated into the biomass (28).

199

200 There are other potential benefits for the fetal-placental unit derived through reliance  
201 on the polyol pathways. Firstly, conversion of glucose to ribose 5-phosphate produces  
202 two molecules of NADPH. NADPH is required for the regeneration of reduced  
203 glutathione from its oxidised form, and hence is key to the antioxidant defences of a cell.  
204 Developing systems are highly prone to perturbation by oxidative stress, which can lead  
205 to severe congenital abnormalities (29, 30). Adequate antioxidant defences are  
206 therefore crucial. Secondly, polyols such as sorbitol are incapable of crossing cell  
207 membranes and so act as powerful osmolytes. Sorbitol is produced from glucose by the  
208 action of aldose reductase, one of the first enzymes to be expressed in the sheep  
209 conceptus. In this species there is a rapid expansion of the embryonic sac into a thread-  
210 like structure, and sorbitol may assist in driving this process by drawing water across  
211 the trophoblast epithelium. In the human there is a similar, though less extensive, need  
212 to expand the extra-embryonic coelom.

213

#### 214 **The benefits of a low oxygen environment for fetal-placental development**

215 Although at first sight the reliance on glycolysis for energy production in the early  
216 placenta and tumours may appear to be inefficient, there is no reason to assume that it  
217 cannot meet the cells' requirements as long as there is a sufficient supply of glucose  
218 (31). In the case of the placenta there is a plentiful supply in the secretions derived from  
219 the endometrial glands, and accumulation of glycogen within the syncytioplasm is a  
220 conspicuous feature during early pregnancy (21, 32).

221

222 Thus, these metabolic pathways enable a high rate of proliferation to be maintained  
223 under a relatively low oxygen concentration. The rise in oxygen concentration within

224 the placenta and the embryonic compartments at the end of the first trimester notably  
225 coincides with the completion of organogenesis. At this stage of development the risk of  
226 teratogenesis falls sharply, as differentiation of the major organ systems is completed.  
227 The risks from oxygen free radicals therefore falls somewhat, and so the metabolic  
228 balance may tip in favour of oxidative phosphorylation. Evidence for such a shift comes  
229 from the rapid fall in placental glycogen content at the end of the first trimester (33),  
230 and it may explain the rise in growth rate of the embryo seen at this stage (34).

231

232 Increasing evidence from the field of stem cell biology indicates that adult stem cell  
233 niches are located in low oxygen environments, roughly equivalent to the intraplacental  
234 oxygen concentration during the first trimester (35). Consistent with this, studies have  
235 revealed that culture of primary cytotrophoblast cells under low oxygen conditions  
236 favours proliferation, whereas higher concentrations promote differentiation and  
237 invasion (36, 37). With respect to this finding, it is notable that levels of CDX2 and ELF5,  
238 two transcription factors that act as gate-keepers of the trophoblast lineage, drop  
239 sharply at the end of the first trimester (38). This suggests a reduction in the  
240 proliferative potential of the placenta, but whether this is due to the three-fold rise in  
241 intra-placental oxygen concentration that occurs at the start of the second trimester (8),  
242 or the loss of growth factors from the endometrial glands with the switch from  
243 histotrophic to haemotrophic nutrition has not yet been clarified.

244

#### 245 **Oxygen and cell differentiation**

246 Within the placenta, a sub-population of trophoblast cells, the extravillous trophoblast,  
247 undergo a partial epithelial-mesenchymal transition and migrate from the outer surface  
248 of the cytotrophoblastic shell into the endometrium (39). In doing so they adopt a

249 pleiotrophic phenotype and move into an area of higher oxygen concentration, for the  
250 decidua is always better oxygenated than the placenta (8). In many ways this resembles  
251 the process of metastasis, albeit a highly regulated one, but the influence of oxygen on  
252 the transition is still unclear. Experimental studies of first trimester explant cultures  
253 have demonstrated that oxygen may be a significant factor, for culture under low oxygen  
254 conditions (3% v 21%) inhibits invasion. This effect is mediated through the HIF-1 and  
255 transforming growth factor beta (TGF $\beta$ ) pathways, and is associated with changes in  
256 matrix metalloproteinase activity (40).

257

258 Local oxygen concentrations also appear to play a role in remodelling of the early  
259 placenta into its definitive form. Villi initially form over the entire surface of the  
260 chorionic sac, but later regress to leave the discoid placenta at the deep pole in contact  
261 with the endometrium, and the smooth membranes. This remodelling coincides with  
262 onset of the maternal arterial circulation to the placenta, which starts preferentially in  
263 the periphery and then extends centripetally, reflecting the degree of trophoblast  
264 invasion and arterial plugging across the placental bed (41). Villi sampled from the  
265 peripheral region display higher levels of oxidative stress and activation of the apoptotic  
266 cascade than their counterparts from the central region, and it has been proposed that  
267 these effects mediate the regression. Excessive regression at this stage of development  
268 may lead to placentas with eccentric insertions of the umbilical cord and more irregular  
269 margins (42, 43).

270

271 Even within the definitive placenta there will be oxygen gradients that reflect the  
272 pattern of maternal arterial blood flow. The placental villi are not arranged at random,  
273 but form 30-40 lobules, each centred over the opening of a maternal spiral artery. The

274 arteries deliver their blood into the relatively villus-free central cavities of a lobule.  
275 From there, the blood percolates through the network of intervillous clefts, exchanging  
276 oxygen with the fetal circulation as it does so, before draining into the openings of the  
277 uterine veins. Each lobule thus represents an individual maternal-fetal exchange unit,  
278 and the pattern of the circulation suggests an oxygen gradient from the arterial centre to  
279 the more venous periphery. This concept is supported by differences in the expression  
280 and activity of the principal antioxidant enzymes (44). These differences in oxygenation  
281 may explain regional variations in villous morphology and enzyme activities (45).  
282 Oxygen gradients similarly occur within tumours due to the limitations of diffusion (15),  
283 and again may mediate cell behaviours, such as resistance to radiotherapy, or  
284 predisposition to metastasis (46).

285

## 286 **Conclusion**

287 Early placental development occurs in a low oxygen environment, and as a rapidly  
288 proliferating tissue it shares many of the same metabolic requirements as tumours.  
289 However, in the placenta there is continual replenishment of oxygen due to plasma  
290 flowing at a slow rate through the intervillous space, and so the tissues do not  
291 experience the increasing drive towards hypoxia that typifies the central regions of  
292 tumours. Nonetheless, oxygen appears to be a major regulator of cell behaviour in both  
293 the placenta and tumours. A better understanding of the similarities and differences  
294 between the two may lead to new insights that are beneficial to these contrasting fields  
295 of biology.

296

## 297 **Acknowledgements**

298 The authors gratefully acknowledge the contributions made by many colleagues during  
299 discussions to the ideas presented in this review.

300

### 301 **Conflict of interest**

302 The authors have no conflicts of interest to declare.

303

304

305

### 306 **References**

- 307 1. Holtan SG, Creedon DJ, Haluska P, Markovic SN. Cancer and pregnancy: parallels  
308 in growth, invasion, and immune modulation and implications for cancer therapeutic  
309 agents. *Mayo Clin Proc.* 2009;84(11):985-1000.
- 310 2. Yedwab GA, Paz G, Homonnai TZ, David MP, Kraicer PF. The temperature, pH, and  
311 partial pressure of oxygen in the cervix and uterus of women and uterus of rats during  
312 the cycle. *Fertil Steril.* 1976;27(3):304-9.
- 313 3. Ottosen LD, Hindkaer J, Husth M, Petersen DE, Kirk J, Ingerslev HJ. Observations  
314 on intrauterine oxygen tension measured by fibre-optic microsensors. *Reprod Biomed*  
315 *Online.* 2006;13(3):380-5.
- 316 4. Houghton FD, Thompson JG, Kennedy CJ, Leese HJ. Oxygen consumption and  
317 energy metabolism of the early mouse embryo. *Mol Reprod Dev.* 1996;44(4):476-85.
- 318 5. Leese HJ. Quiet please, do not disturb; a hypothesis of embryo metabolism and  
319 viability. *BioEssays.* 2002;24:845-9.
- 320 6. Sturmeiy RG, Hawkhead JA, Barker EA, Leese HJ. DNA damage and metabolic  
321 activity in the preimplantation embryo. *Hum Reprod.* 2009;24(1):81-91.
- 322 7. Rodesch F, Simon P, Donner C, Jauniaux E. Oxygen measurements in endometrial  
323 and trophoblastic tissues during early pregnancy. *Obstetrics and Gynecology.*  
324 1992;80:283-5.
- 325 8. Jauniaux E, Watson AL, Hempstock J, Bao Y-P, Skepper JN, Burton GJ. Onset of  
326 maternal arterial bloodflow and placental oxidative stress; a possible factor in human  
327 early pregnancy failure. *American Journal of Pathology.* 2000;157:2111-22.
- 328 9. Hustin J, Schaaps JP. Echographic and anatomic studies of the  
329 maternotrophoblastic border during the first trimester of pregnancy. *American Journal*  
330 *of Obstetrics and Gynecology.* 1987;157:162-8.
- 331 10. Burton GJ, Jauniaux E, Watson AL. Maternal arterial connections to the placental  
332 intervillous space during the first trimester of human pregnancy; the Boyd Collection  
333 revisited. *Am J Obstet Gynecol.* 1999;181:718-24.
- 334 11. Castellucci M, Kaufmann P. A three-dimensional study of the normal human  
335 placental villous core: II. Stromal architecture. *Placenta.* 1982;3:269-86.

- 336 12. Jauniaux E, Jurkovic D, Campbell S. In vivo investigations of anatomy and  
337 physiology of early human placental circulations. *Ultrasound in Obstetrics and*  
338 *Gynecology*. 1991;1:435-45.
- 339 13. Hofmann OM, Brittain T. Partitioning of oxygen and carbon monoxide in the three  
340 human embryonic hemoglobins. *Hemoglobin*. 1998;22(4):313-9.
- 341 14. Carter AM. Evolution of factors affecting placental oxygen transfer. *Placenta*.  
342 2009;30 Suppl A:S19-25.
- 343 15. Bertuzzi A, Gandolfi A. Cell kinetics in a tumour cord. *J Theor Biol*.  
344 2000;204(4):587-99.
- 345 16. Burton GJ, Jauniaux E, Charnock-Jones DS. Human early placental development:  
346 potential roles of the endometrial glands. *Placenta*. 2007;28 Suppl A:S64-9.
- 347 17. Maruo T, Matsuo H, Murata K, Mochizuki M. Gestational age-dependent dual  
348 action of epidermal growth factor on human placenta early in gestation. *J Clin*  
349 *Endocrinol Metab*. 1992;75:1362-7.
- 350 18. Forbes K, Westwood M, Baker PN, Aplin JD. Insulin-like growth factor I and II  
351 regulate the life cycle of trophoblast in the developing human placenta. *Am J Physiol Cell*  
352 *Physiol*. 2008;294(6):C1313-22.
- 353 19. Seppälä M, Jukunen M, Riitinen L, Koistinen R. Endometrial proteins: a  
354 reappraisal. *Human Reproduction*. 1992;7 Suppl. 1:31-8.
- 355 20. Cindrova-Davies T, van Patot MT, Gardner L, Jauniaux E, Burton GJ, Charnock-  
356 Jones DS. Energy status and HIF signalling in chorionic villi show no evidence of hypoxic  
357 stress during human early placental development. *Mol Hum Reprod*. 2015;21(3):296-  
358 308.
- 359 21. Burton GJ, Watson AL, Hempstock J, Skepper JN, Jauniaux E. Uterine glands  
360 provide histiotrophic nutrition for the human fetus during the first trimester of  
361 pregnancy. *J Clin Endocrinol Metab*. 2002;87:2954-9.
- 362 22. Jauniaux E, Watson AL, Burton GJ. Evaluation of respiratory gases and acid-base  
363 gradients in fetal fluids and uteroplacental tissue between 7 and 16 weeks. *American*  
364 *Journal of Obstetrics and Gynecology*. 2001;184:998-1003.
- 365 23. Jauniaux E, Hempstock J, Teng C, Battaglia F, Burton GJ. Polyol concentrations in  
366 the fluid compartments of the human conceptus during the first trimester of pregnancy;  
367 maintenance of redox potential in a low oxygen environment. *J Clin Endocrinol Metab*.  
368 2005;90:1171-5.
- 369 24. Semenza GL, Roth PH, Fang HM, Wang GL. Transcriptional regulation of genes  
370 encoding glycolytic enzymes by hypoxia-inducible factor 1. *J Biol Chem*.  
371 1994;269(38):23757-63.
- 372 25. Kim JW, Tchernyshyov I, Semenza GL, Dang CV. HIF-1-mediated expression of  
373 pyruvate dehydrogenase kinase: a metabolic switch required for cellular adaptation to  
374 hypoxia. *Cell Metab*. 2006;3(3):177-85.
- 375 26. Papatreou I, Cairns RA, Fontana L, Lim AL, Denko NC. HIF-1 mediates  
376 adaptation to hypoxia by actively downregulating mitochondrial oxygen consumption.  
377 *Cell Metab*. 2006;3(3):187-97.
- 378 27. Simon MC. Coming up for air: HIF-1 and mitochondrial oxygen consumption. *Cell*  
379 *Metab*. 2006;3(3):150-1.
- 380 28. Vander Heiden MG, Cantley LC, Thompson CB. Understanding the Warburg effect:  
381 the metabolic requirements of cell proliferation. *Science*. 2009;324(5930):1029-33.
- 382 29. Eriksson UJ. Oxidative DNA damage and embryo development. *Nature Medicine*.  
383 1999;5:715.

- 384 30. Nicol CJ, Zielenski J, Tsui L-C, Wells PG. An embryoprotective role for glucose-6-  
385 phosphate dehydrogenase in developmental oxidative stress and chemical  
386 teratogenesis. *FASEB Journal*. 2000;14:111-27.
- 387 31. Collier HA. Is cancer a metabolic disease? *Am J Pathol*. 2014;184(1):4-17.
- 388 32. Boyd JD. Glycogen in early human implantation sites. *Memoirs of the Society for*  
389 *Endocrinology*. 1959;6:26-34.
- 390 33. Robb SA, Hytten FE. Placental glycogen. *Br J Obstet Gynaecol*. 1976;83(1):43-53.
- 391 34. van Uitert EM, Exalto N, Burton GJ, Willemsen SP, Koning AH, Eilers PH, et al.  
392 Human embryonic growth trajectories and associations with fetal growth and  
393 birthweight. *Hum Reprod*. 2013;28(7):1753-61.
- 394 35. Mohyeldin A, Garzon-Muvdi T, Quinones-Hinojosa A. Oxygen in stem cell biology:  
395 a critical component of the stem cell niche. *Cell Stem Cell*. 2010;7(2):150-61.
- 396 36. Genbacev O, Zhou Y, Ludlow JW, Fisher SJ. Regulation of human placental  
397 development by oxygen tension. *Science*. 1997;277:1669-72.
- 398 37. Zhou S, Xie Y, Puscheck EE, Rappolee DA. Oxygen levels that optimize TSC culture  
399 are identified by maximizing growth rates and minimizing stress. *Placenta*.  
400 2011;32(6):475-81.
- 401 38. Hemberger M, Udayashankar R, Tesar P, Moore H, Burton GJ. ELF5-enforced  
402 transcriptional networks define an epigenetically regulated trophoblast stem cell  
403 compartment in the human placenta. *Mol Hum Genet*. 2010;19:2456-67.
- 404 39. Davies JE, Pollheimer J, Yong HE, Kokkinos MI, Kalionis B, Knofler M, et al.  
405 Epithelial-mesenchymal transition during extravillous trophoblast differentiation. *Cell*  
406 *Adh Migr*. 2016;10(3):310-21.
- 407 40. Caniggia I, Mostachfi H, Winter J, Gassmann M, Lye SJ, Kuliszewski M, et al.  
408 Hypoxia-inducible factor-1 mediates the biological effects of oxygen on human  
409 trophoblast differentiation through TGFbeta(3). *J Clin Invest*. 2000;105(5):577-87.
- 410 41. Jauniaux E, Hempstock J, Greenwold N, Burton GJ. Trophoblastic oxidative stress  
411 in relation to temporal and regional differences in maternal placental blood flow in  
412 normal and abnormal early pregnancies. *American Journal of Pathology*. 2003;162:115-  
413 25.
- 414 42. Burton GJ, Jauniaux E, Charnock-Jones DS. The influence of the intrauterine  
415 environment on human placental development. *Int J Dev Biol*. 2010;54:303-12.
- 416 43. Salafia CM, Yampolsky M, Shlakhter A, Mandel DH, Schwartz N. Variety in  
417 placental shape: when does it originate? *Placenta*. 2012;33(3):164-70.
- 418 44. Hempstock J, Bao Y-P, Bar-Issac M, Segaren N, Watson AL, Charnock Jones DS, et  
419 al. Intralobular differences in antioxidant enzyme expression and activity reflect oxygen  
420 gradients within the human placenta. *Placenta*. 2003;24:517-23.
- 421 45. Schuhmann RA. Placentone structure of the human placenta. *Bibliotheca*  
422 *Anatomica*. 1982;22:46-57.
- 423 46. Lewis DM, Park KM, Tang V, Xu Y, Pak K, Eisinger-Mathason TS, et al.  
424 Intratumoral oxygen gradients mediate sarcoma cell invasion. *Proc Natl Acad Sci U S A*.  
425 2016;113(33):9292-7.
- 426

427

428 **Figure legend**



429 Figure 1. Schematic representation of the interconnections between glycolysis and some  
430 of the polyol pathways. Polyols that are at high concentrations in the first trimester  
431 placenta are shown in green, and their synthesis enables the regeneration of NAD<sup>+</sup> and  
432 NADP<sup>+</sup> under low oxygen conditions independent of the TCA acid cycle. NAD<sup>+</sup> is required  
433 to maintain glycolysis and production of ATP, whereas NADP<sup>+</sup> is important for the  
434 generation of reduced glutathione. By contrast, in tumours NAD<sup>+</sup> is regenerated  
435 principally through fermentation of pyruvate to lactate under the action of lactate  
436 dehydrogenase (LDH). Pathways activated in tumours are shown in red, and include  
437 stabilisation of HIF through increasing hypoxia. HIF promotes glycolysis and LDH, but  
438 inhibits pyruvate dehydrogenase (PDH) and so blocks the conversion of pyruvate to  
439 acetyl-CoA. Consequently, there is a build-up of intermediates in the glycolytic pathway,  
440 favouring the diversion of carbon skeletons for synthesis of nucleic acids. Some  
441 oncogenes promote cell proliferation through similar effects. PEP;  
442 phosphoenolpyruvate.