

# Semi-artificial photoelectrochemistry for CO<sub>2</sub>-mediated enantioselective organic synthesis

Tessel Bouwens<sup>1,a</sup>, Samuel J. Cobb<sup>1,b</sup>, Celine W. S. Yeung<sup>1</sup>, Yongpeng Liu<sup>1</sup>, Guilherme Martins<sup>2</sup>, Inês A. C. Pereira<sup>2</sup>, Erwin Reisner<sup>1\*</sup>

<sup>1</sup>Yusuf Hamied Department of Chemistry, University of Cambridge, Cambridge CB2 1EW, U.K. <sup>2</sup>Instituto de Tecnologia Química e Biológica António Xavier (ITQB NOVA), Universidade NOVA de Lisboa, Av. da República, 2780-157 Oeiras, Portugal.

**ABSTRACT:** Photoelectrochemical (PEC) cells are under intensive development for the synthesis of solar fuels, but CO<sub>2</sub> reduction typically only results in simple building blocks such as HCOO<sup>-</sup>. Here, we demonstrate that CO<sub>2</sub>-converting PEC cells can drive integrated enzymatic domino catalysis to produce chiral organic molecules by using CO<sub>2</sub>/HCOO<sup>-</sup> as a sustainable redox couple. First, we establish a semi-artificial electrode consisting of three enzymes co-immobilized on a high surface area electrode based on carbon felt covered by a mesoporous indium tin oxide (ITO) coating. When applying a mild cathodic potential (−0.25 V vs. the reversible hydrogen electrode (RHE)), CO<sub>2</sub> is reduced to HCOO<sup>-</sup> using a W-formate dehydrogenase (FDH<sub>NvH</sub>) from *Nitratidesulfovibrio vulgaris* Hildenborough, which then enables the reduction of NAD<sup>+</sup> to NADH by an NAD<sup>+</sup>-cofactor dependent formate dehydrogenase from *Candida Boidinii* (FDH<sub>CB</sub>). Subsequently, an alcohol dehydrogenase (ADH) uses NADH generated from CO<sub>2</sub>/HCOO<sup>-</sup> cycling to reduce acetophenone into chiral 1-phenylethanol in good enantiomeric excess (93%) and conversion yields (39%). Depending on the specific ADH (ADH<sub>S</sub> or ADH<sub>R</sub>), either (*S*) or (*R*)-1-phenylethanol can be synthesized at pH 6 and 20 °C. To illustrate solar energy utilization, we integrate the three nanoconfined enzymes with a PEC platform based on an integrated organic semiconductor (OSC) photocathode to allow for enantioselective synthesis (at +0.8 V vs. RHE) based on a solar fuel device. This proof-of-principle demonstration shows that concepts and devices from artificial photosynthesis can be readily translated to precise and sustainable biocatalysis, including the production of chiral organic molecules using light.

Photosynthesis has long inspired scientists as it harnesses solar energy to convert simple building blocks (CO<sub>2</sub>, H<sub>2</sub>O) into complex organic chemicals (glucose) while releasing O<sub>2</sub>.<sup>1,2</sup> Accordingly, artificial photosynthetic devices such as photoelectrochemical (PEC) cells aim to mimic this process, but they can currently only produce relatively simple chemical fuels (e.g., H<sub>2</sub>, HCOO<sup>-</sup>, CO, CH<sub>4</sub>, ethylene, ethanol, or propanol).<sup>3–7</sup> Current efforts to mimic the natural process have mainly focused on the light-dependent reactions in the thylakoid membrane, while the compartmentalized multi-enzyme machinery responsible for glucose synthesis from CO<sub>2</sub> occurring in the light-independent stromal reactions has received limited attention.<sup>8</sup> As mature PEC devices with high efficiency and stability are emerging, a growing opportunity arises to expand the reactivity space from simple fuels to more complex organic molecules by integrating cascades in devices, and thereby bringing us a step closer to truly mimicking the reactivity of natural photosynthesis.

State-of-the-art PEC cells provide an excellent platform to expand from classical solar fuels reactivity (water splitting, CO<sub>2</sub> reduction) to enantioselective organic synthesis, in line with our ongoing mission to integrate the fields of artificial photosynthesis and organic chemistry.<sup>9</sup> While the recent renaissance of organic photoredox catalysis has made

substantial progress in developing visible light responsive photocatalysts with many demonstrations of noteworthy reactions, it has not yet taken full advantage of the opportunities provided by chiral bio-catalytic synthesis under benign, aqueous conditions and solid-state devices employing state-of-the-art semiconductors.

In this study, we aim to close this gap and develop a platform that uses solar energy to drive biocatalysis integrated in a PEC device for chiral organic synthesis. We employ evolutionarily-optimized enzymes as biocatalysts to tackle the bottleneck of synthetic catalysts to overcome efficiency, selectivity and reactivity issues.<sup>10–12</sup> While semi-artificial PEC cells employing enzymes are known, only the production of simple fuels such as H<sub>2</sub>,<sup>13,14</sup> CO,<sup>15</sup> HCOO<sup>-</sup><sup>16,17</sup> and MeOH<sup>18,19</sup> has so far been demonstrated. However, oxidoreductases provide an opportunity to catalyze an exciting diversity of reactions, including control of chemo-, regio-, and stereoselectivities, which outcompetes the possibilities provided by synthetic catalysts.<sup>20–22</sup> To provide a glimpse of possibilities for the future, we demonstrate here that PEC cells producing classical solar fuels, such as HCOO<sup>-</sup> from CO<sub>2</sub>, can be readily modified using cofactor dependent enzymes for chiral synthesis using light (Figure 1).




**Figure 1.** Schematic representation of the PEC-cascade system using  $\text{CO}_2/\text{HCOO}^-$  as a mediator to generate chiral building blocks featuring  $\text{FDH}_{\text{NvH}}$ ,  $\text{FDH}_{\text{CB}}$  and ADH. All three enzymes are integrated into a Ti|ITO-CF scaffold (bottom left) for electrochemical characterization and an OPV-based photocathode, giving OPV|ITO-CF| $\text{FDH}_{\text{NvH}}/\text{FDH}_{\text{CB}}/\text{ADH}$ . The SEM and EDX images of Ti|ITO-CF electrodes demonstrating the ITO NPs on CF.

The integrated enzyme cascade consists of three enzymes. The first enzyme is formate dehydrogenase ( $\text{FDH}_{\text{NvH}}$ ) from *Nitratidesulfovibrio vulgaris* Hildenborough (NvH),<sup>23,24</sup> which is known to attach in an electroactive configuration to porous ITO electrodes via non-covalent interactions, allowing for  $\text{CO}_2$  reduction to  $\text{HCOO}^-$  with marginal overpotential and a  $K_{\text{M}}$  for  $\text{CO}_2$  of 0.42 mM.<sup>25–27</sup> The next enzyme is the  $\text{NAD}^+$ -cofactor dependent formate dehydrogenase from *Candida Boidinii* ( $\text{FDH}_{\text{CB}}$ ), which uses  $\text{HCOO}^-$  to catalyze the reduction of  $\text{NAD}^+$  to NADH. The last enzyme is an alcohol dehydrogenase (ADH), which can use the NADH produced from  $\text{CO}_2/\text{HCOO}^-$  cycling to reduce pro-chiral acetophenone (selected as model building block for pharmaceutical applications<sup>28,29</sup>).

Porous metal oxide electrodes made from indium tin oxide (ITO) provide conductivity, a robust interaction with electroactive enzymes, and their porosity provide a high surface area to support large quantities of enzymes.<sup>30–32</sup> However, the brittleness and conductivity constraints of porous ITO films only allows for film thicknesses up to approximately 50  $\mu\text{m}$ .<sup>33,34</sup> Therefore, we developed a conducting scaffold for enzyme immobilization in this study, using commercially available carbon felt (CF) cuboids ( $5 \times 5 \times 3.2 \text{ mm}^3$ ) coated with ITO nanoparticles. These macroscopic cuboids provide mm-thickness and can be readily connected to a titanium foil ( $20 \times 10 \text{ mm}^2$ ) using graphite epoxy (Figures S1–S4, further discussion in Supporting Information). These Ti|ITO-CF electrodes can be used as a (dark) cathode or integrated into a photocathode (see below). The maximum loading of ITO NPs onto CF was determined (Figure S1), and the CF cuboids demonstrated an apparent porosity of 78%, close to the reported porosity of these CF materials (>80%). Ti|ITO-CF were characterized by SEM and EDX (Figure 1, Figures S2–S4), which shows the individual ITO NPs and the presence of In, Sn and O on the carbon fibers.

$\text{FDH}_{\text{NvH}}$  (125 pmol) was then dropcast onto the Ti|ITO-CF electrode ( $0.25 \text{ cm}^2$ ) and analyzed using protein film voltammetry (PFV) in an aqueous electrolyte solution containing MOPS (13.5 mL, 0.1 M, pH 6), DMSO (1.5 mL),  $\text{NAD}^+$  (1 mM) and acetophenone (50 mM), saturated with  $\text{CO}_2$  at ambient temperature (20 °C, Table S1). The PFV response shows an onset of catalytic current for  $\text{CO}_2$  reduction close to 0 V vs. the reversible hydrogen electrode (RHE), as expected for the reversible  $\text{CO}_2$  to  $\text{HCOO}^-$  biocatalyst (Figure 2).<sup>32</sup> During the back scan, an anodic wave is observed with a peak current at approximately +0.2 V vs. RHE, which is only present upon scan reversal following the catalytic cathodic wave (Figure 2b). This observation suggests that this cathodically induced anodic wave is due to the oxidation of  $\text{HCOO}^-$  produced from  $\text{CO}_2$  reduction in the porous stationary Ti|ITO-CF| $\text{FDH}_{\text{NvH}}$  electrode. In the absence of  $\text{CO}_2$ , no catalytic currents are observed (Figure 2a).

Next, we studied the electrochemical response with all three enzymes, using  $\text{FDH}_{\text{NvH}}$  (125 pmol),  $\text{FDH}_{\text{CB}}$  (1 nmol) and ADH (5 nmol) dropcast onto Ti|ITO-CF ( $0.25 \text{ cm}^2$ ). Note that solely  $\text{FDH}_{\text{NvH}}$  is an electroactive enzyme receiving electrons directly from Ti|ITO-CF, whereas  $\text{FDH}_{\text{CB}}$  and ADH rely on  $\text{HCOO}^-$  and NADH as electron source. In a  $\text{CO}_2$ -saturated solution and in the presence of  $\text{NAD}^+$  and acetophenone we observe a qualitatively similar voltametric cathodic response for Ti|ITO-CF| $\text{FDH}_{\text{NvH}}/\text{FDH}_{\text{CB}}/\text{ADH}$  compared to Ti|ITO-CF| $\text{FDH}_{\text{NvH}}$ , but the current density of the anodic wave at +0.2 V vs. RHE is significantly reduced (Figure 2a). This observation supports that  $\text{HCOO}^-$  is readily consumed by  $\text{FDH}_{\text{CB}}$  and used to generate the NADH cofactor that is needed to produce the chiral alcohol product.

Protein film chronoamperometry (PF-CA) was performed with ITO-CF| $\text{FDH}_{\text{NvH}}$  at an applied potential of  $-0.25 \text{ V}$  vs. RHE (Figure S5), showing quantitative conversion of  $\text{CO}_2$  into  $\text{HCOO}^-$  (Faradaic efficiency,  $\text{FE}_{\text{HCOO}^-} = 99\%$ ; determined by  $^1\text{H}$  NMR spectroscopy, Figures S6–S8). PF-CA with all

three enzymes produces 1-phenylethanol (PE), demonstrating that the cascade is functional electrochemically ( $FE_{HCOO^-} = 72\%$  and  $FE_{PE} = 27\%$ ,  $FE_{total} = 99\%$ ).




**Figure 2.** (a) Protein film voltammetry (PFV) scans of FDH<sub>NvH</sub> (red and blue traces) and 3-enzyme cascade (black trace) under CO<sub>2</sub> (red and black traces) or N<sub>2</sub> (blue trace). (b) PFV analysis of FDH<sub>NvH</sub>, start at +50 mV to 550 mV to -250 mV and back (red trace). PFV scans recorded using 10 mV s<sup>-1</sup> in electrolyte comprised of MOPS, (13.5 mL, 0.1 M, pH 6), NAD<sup>+</sup> (1 mM), acetophenone (50 mM) and DMSO-d<sub>6</sub> (1.5 mL) (Table S1).

The turnover number of ADH ( $TON_{ADH}$ ) for 1-phenylethanol synthesis is estimated from PF-CA at approximately  $5 \times 10^3$  after 12 h (Figure S5, Figures S9–S10). The cascade generates chiral 1-phenylethanol in high conversions after 12 h ( $38 \pm 8\%$ , Table S2, Figure S7f). Depending on the choice of ADH (Figures S11–S12), we were able to produce both the (*S*)-enantiomer (enantiomeric excess,  $ee = 93\%$ , determined by chiral HPLC) or the (*R*)-enantiomer ( $ee = 59\%$ ) (Figure 2b, Figures S13–S14).

Isotopic labeling experiments using <sup>13</sup>CO<sub>2</sub> with the Ti|ITO-CF|FDH<sub>NvH</sub>/FDH<sub>CB</sub>/ADH electrode at an applied potential of -0.25 V vs. RHE for 4 hours confirms the origin of HCOO<sup>-</sup> solely from CO<sub>2</sub> (<sup>1</sup>H-NMR,  $\delta = 8.44$  ppm, doublet,  $J_{C-H} = 194$  Hz, Figure S15). A control experiment using ITO-CF|FDH<sub>NvH</sub>/FDH<sub>CB</sub>/ADH without NAD<sup>+</sup> resulted in negligible amounts of 1-phenylethanol (Figure S16).

The FE and  $ee$  for Ti|ITO-CF|FDH<sub>NvH</sub>/FDH<sub>CB</sub>/ADH are comparable to previous work employing electrocatalytic enzymatic cascades based on NAD(P)H-recycling using ferredoxin NADP<sup>+</sup> reductase.<sup>22,35,36</sup> Synthetic co-factor regeneration strategies have also been applied previously, e.g., using Cp\*Rh(bpy)-derived mediators, but these systems suffer from side reactions, such as H<sub>2</sub> formation ( $FE_{NADH} < 86\%$ ).<sup>18,37–45</sup> Our semi-artificial platform operates selectively (quantitative  $FE_{NADH}$ , no side-products detected) with a high  $k_{cat}(NAD^+)_{FDH} = 8400$  h<sup>-1</sup>, whereas Cp\*Rh(bpy)-derived mediators only reach  $k_{cat}(NAD^+)_{Rh} = 36$  h<sup>-1</sup>.<sup>46</sup> Electrocatalytic production was also integrated with ADH-promoted synthesis of 1-propanol from 1-propanal via Cp\*Rh(bpy)-mediated cofactor regeneration,<sup>44</sup> but this system requires a flow-through electrolytic cell with a negative potential of -1.6 V vs. RHE.

The interfacial electron transfer kinetics were investigated by electrochemical impedance spectroscopy (EIS)<sup>47</sup> over

the same potential range used for the PFV scans with 0.1 V intervals and a sinusoidal perturbation of 15 mV. Quantitative analysis with Nyquist plot fitting (Figures S17–S20) has previously been validated for electroenzymatic reactions involving H<sub>2</sub>ase and FDH<sub>NvH</sub>.<sup>48–50</sup> The values for the resistance  $R_e$  remain similar across different systems, thus the values of time constant  $\tau_e$ , defined by  $\tau_e = R_e \times C_e$ , are driven by the distinct differences in  $C_e$  values. The time constant describes all the processes occurring within the measurement time domain. The addition of FDH<sub>CB</sub> and FDH<sub>CB</sub>/ADH lead to a smaller  $\tau_e$ , suggesting that the CO<sub>2</sub>/HCOO<sup>-</sup>-cycling for Ti|ITO-CF|FDH<sub>NvH</sub>/FDH<sub>CB</sub>/ADH is faster than the single process at the Ti|ITO-CF|FDH<sub>NvH</sub> electrode (further discussion in Supporting Information).<sup>51</sup>

After establishing the semi-artificial electrocatalytic domino system, we integrated an organic photovoltaic (OPV)-based PEC system based on a recently established  $\pi$ -conjugated organic semiconductor (OSC) PEC platform.<sup>50</sup> Specifically, we employ an OPV device based on PCE10:EH-IDTBR that is encapsulated by graphite epoxy to (i) protect the OPV from the aqueous electrolyte solution and (ii) integrate the ITO-CF electrode scaffold (Figure 1).<sup>52</sup> The ITO-CF cuboid is connected to the OPV using graphite-epoxy paste and the photocathode is wired to a metal rod. The enzymes were dropcast onto the OPV|ITO-CF using the same procedure as for the electrochemical studies on Ti|ITO-CF.

The OPV|ITO-CF|FDH<sub>NvH</sub>/FDH<sub>CB</sub>/ADH photocathode displays an onset potential of 1 V vs. RHE with a high photocurrent density of approximately -6 mA cm<sup>-2</sup> at 0 V vs. RHE under standard solar spectrum air mass 1.5 global (AM1.5G) irradiation at 20 °C (pH 6, Figure 3a, Figure S21). PF-CA showed a relatively stable current density of ~1 mA cm<sup>-2</sup> at +0.8 V vs. RHE during 12 hours AM1.5G irradiation (Figure 3b). The generation of the intermediate formate and (*S*)-1-phenylethanol was followed by <sup>1</sup>H NMR spectroscopy (Figure 3c, Figures S22–S23). After 12 hours, a  $FE_{PE}$  of 10% with a  $FE_{HCOO^-}$  of 54% was obtained. Our OPV|ITO-CF|FDH<sub>NvH</sub>/FDH<sub>CB</sub>/ADH photocathode therefore demonstrates solar-powered recycling of NADH using CO<sub>2</sub>/HCOO<sup>-</sup> as a mediator for the enantioselective synthesis of (*S*)-1-phenylethanol ( $TON_{ADH}$  approximately  $1.2 \times 10^3$ , Figures S23–S24). The longevity of the current domino PEC system is limited by NADH degradation during irradiation (Figure S25).

In summary, we have demonstrated the integration of three enzymes into a state-of-the-art PEC cell using CO<sub>2</sub> as a redox mediator to produce NADH to drive the synthesis of chiral organics (both (*R*) and (*S*)-enantiomers). This strategy provides a platform to leverage enzyme-driven photoelectrochemistry to synthesize chiral organic chemicals from simple substrates for the chemical and pharmaceutical industry in the future.<sup>28,29,53</sup> This principle can be readily adopted by contemporary solar fuel devices producing (i) other fuels; e.g., H<sub>2</sub> using a cofactor dependent hydrogenase,<sup>13,54</sup> and (ii) other photocathodes materials; e.g., encapsulated silicon, copper oxide, lead halide perovskite.<sup>4,5,55</sup> This work also intends to inspire and motivate efforts to connect the materials- and device-focused artificial photosynthesis

community with organic chemists employing biocatalysis and photoredox catalysis.




**Figure 3.** Production of chiral (*S*)-1-phenylethanol using a PEC system: OPV|ITO-CF|FDH<sub>NVH</sub>/FDH<sub>CB</sub>/ADH. a) PFV scans in dark, light and chopped with on/off cycles of 10 sec. b) Chopped light PF-CA at +0.8 V vs. RHE with 50 min (green/orange) to 10 min (white) on/off-cycles. NMR samples were collected after  $t=0, 1, 2, 3, 4, 6, 9$  and 12 h. c) FE<sub>PE</sub>, FE<sub>HCOO<sup>-</sup></sub> and FE<sub>Total</sub>. \*Calculated from deconvoluted NMR signals.

#### ASSOCIATED CONTENT

**Data Availability Statement.** Experimental data of this study can be accessed through the University of Cambridge data repository: <https://doi.org/10.17863/CAM.117123>.

**Supporting Information.** The Supporting Materials consists of experimental procedures, material characterization and additional spectroscopic data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

#### AUTHOR INFORMATION

##### Corresponding Author

\* Erwin Reisner – Yusuf Hamied Department of Chemistry, University of Cambridge, Cambridge CB2 1EW, U.K.; [orcid.org/0000-0002-7781-1616](https://orcid.org/0000-0002-7781-1616); Email: [reisner@ch.cam.ac.uk](mailto:reisner@ch.cam.ac.uk)

##### Authors

**Tessel Bouwens** – Yusuf Hamied Department of Chemistry, University of Cambridge, Cambridge CB2 1EW, U.K.; [orcid.org/0000-0003-4754-8930](https://orcid.org/0000-0003-4754-8930)

**Sam J. Cobb** – Yusuf Hamied Department of Chemistry, University of Cambridge, Cambridge CB2 1EW, U.K.; [orcid.org/0000-0001-5015-8090](https://orcid.org/0000-0001-5015-8090).

**Celine Wing See Yeung**, Yusuf Hamied Department of Chemistry, University of Cambridge, Cambridge CB2 1EW, U.K.; [orcid.org/0000-0003-4954-7288](https://orcid.org/0000-0003-4954-7288).

**Yongpeng Liu** – Yusuf Hamied Department of Chemistry, University of Cambridge, Cambridge CB2 1EW, U.K.; [orcid.org/0000-0002-4544-4217](https://orcid.org/0000-0002-4544-4217).

**Guilherme Martins** – Instituto de Tecnologia Química e Biológica António Xavier (ITQB NOVA), Universidade NOVA de Lisboa, 2780-157 Oeiras, Portugal; [orcid.org/0009-0000-6042-7151](https://orcid.org/0009-0000-6042-7151).

**Inês A. C. Pereira** – Instituto de Tecnologia Química e Biológica António Xavier (ITQB NOVA), Universidade NOVA de Lisboa, 2780-157 Oeiras, Portugal; [orcid.org/0000-0003-3283-4520](https://orcid.org/0000-0003-3283-4520).

##### Present Addresses

<sup>a</sup> **Tessel Bouwens** – Department of Chemical Engineering, Delft University of Technology, Van der Maasweg 9, 2629 HZ Delft, The Netherlands; [orcid.org/0000-0003-4754-8930](https://orcid.org/0000-0003-4754-8930)

<sup>b</sup> **Sam J. Cobb** – Department of Chemistry, The University of Manchester, Oxford Road, Manchester M13 9PL, UK; [orcid.org/0000-0001-5015-8090](https://orcid.org/0000-0001-5015-8090).

#### Author Contributions

The manuscript was written through contributions of all authors.

#### ACKNOWLEDGMENTS

We acknowledge funding from the Dutch Research Council (NWO Rubicon to T.B.), UK Research & Innovation (UKRI, ERC Advanced Grant EP/X030563/1 to E.R.), UK Department of Science, Innovation & Technology and the Royal Academy of Engineering Chair in Emerging Technologies programme (CIET-2324-83 to E.R.), the Singapore Agency for Science, Technology and Research (A\*STAR) for a Ph.D. studentship (to C.W.S.Y.), the EPSRC Cambridge NanoDTC (EP/S022953/1 to C.W.S.Y.), the Leverhulme Trust for an Early Career Fellowship (ECF-2021-072 to S.J.C. and ECF-2024-230 to Y.L.), the Isaac Newton Trust (20.08(r) to S.J.C. and 24.08(s) to Y.L.), the Fundação para a Ciência e Tecnologia (FCT, Portugal) through fellowship DFA/BD/7897/2020 (R. R. M.), MOSTMICRO-ITQB (UIDB/04612/2020 and UIDP/04612/2020) and Associated Laboratory LS4FUTURE (LA/P/0087/2020) (to IACP). We thank Dr Carolina Pulignani and Dr Leonardo Castañeda-Losada (University of Cambridge) for help with HPLC, Benjamin Craig for preliminary electrochemical experiments (University of Cambridge), Dr Rita R. Manuel (ITQB NOVA) for help in protein purification and Beverly Low and Dr Motiar Rahman for discussions regarding this work.

#### REFERENCES

- (1) Lewis, N. S.; Nocera, D. G. Powering the Planet: Chemical Challenges in Solar Energy Utilization. *Proc. Natl. Acad. Sci. U. S. A.* **2006**, *103* (43), 15729–15735. <https://doi.org/10.1073/pnas.0603395103>.
- (2) Yoshino, S.; Takayama, T.; Yamaguchi, Y.; Iwase, A.; Kudo, A. CO<sub>2</sub> Reduction Using Water as an Electron Donor over Heterogeneous Photocatalysts Aiming at Artificial Photosynthesis. *Acc. Chem. Res.* **2022**, *55* (7), 966–977. <https://doi.org/10.1021/acs.accounts.1c00676>.
- (3) Roh, I.; Yu, S.; Lin, C.-K.; Louisia, S.; Cestellos-Blanco, S.; Yang, P. Photoelectrochemical CO<sub>2</sub> Reduction toward Multicarbon

- Products with Silicon Nanowire Photocathodes Interfaced with Copper Nanoparticles. *J. Am. Chem. Soc.* **2022**, *144* (18), 8002–8006. <https://doi.org/10.1021/jacs.2c03702>.
- (4) Xia, M.; Pan, L.; Liu, Y.; Gao, J.; Li, J.; Mensi, M.; Sivula, K.; Zakeeruddin, S. M.; Ren, D.; Grätzel, M. Efficient Cu<sub>2</sub>O Photocathodes for Aqueous Photoelectrochemical CO<sub>2</sub> Reduction to Formate and Syngas. *J. Am. Chem. Soc.* **2023**, *145* (51), 27939–27949. <https://doi.org/10.1021/jacs.3c06146>.
- (5) Andrei, V.; Reuillard, B.; Reisner, E. Bias-Free Solar Syngas Production by Integrating a Molecular Cobalt Catalyst with Perovskite–BiVO<sub>4</sub> Tandems. *Nat. Mater.* **2020**, *19* (2), 189–194. <https://doi.org/10.1038/s41563-019-0501-6>.
- (6) Rahaman, M.; Andrei, V.; Wright, D.; Lam, E.; Pornrungroj, C.; Bhattacharjee, S.; Pichler, C. M.; Greer, H. F.; Baumberg, J. J.; Reisner, E. Solar-Driven Liquid Multi-Carbon Fuel Production Using a Standalone Perovskite–BiVO<sub>4</sub> Artificial Leaf. *Nat. Energy*, **2023**, *8* (6), 629–638. <https://doi.org/10.1038/s41560-023-01262-3>.
- (7) Zhou, B.; Ou, P.; Pant, N.; Cheng, S.; Vanka, S.; Chu, S.; Rashid, R. T.; Botton, G.; Song, J.; Mi, Z. Highly Efficient Binary Copper–iron Catalyst for Photoelectrochemical Carbon Dioxide Reduction toward Methane. *Proc. Natl. Acad. Sci. U. S. A.* **2020**, *117* (3), 1330–1338. <https://doi.org/10.1073/pnas.1911159117>.
- (8) Miller, T. E.; Beneyton, T.; Schwander, T.; Diehl, C.; Girault, M.; McLean, R.; Chotel, T.; Claus, P.; Cortina, N. S.; Baret, J.-C.; Erb, T. J. Light-Powered CO<sub>2</sub> Fixation in a Chloroplast Mimic with Natural and Synthetic Parts. *Science*, **2020**, *368* (6491), 649–654. <https://doi.org/10.1126/science.aaz6802>.
- (9) Reisner, E. When Does Organic Photoredox Catalysis Meet Artificial Photosynthesis? *Angew. Chem. Int. Ed.* **2019**, *58* (12), 3656–3657. <https://doi.org/10.1002/anie.201814692>.
- (10) Goldberg, K.; Schroer, K.; Lütz, S.; Liese, A. Biocatalytic Ketone Reduction - A Powerful Tool for the Production of Chiral Alcohols - Part I: Processes with Isolated Enzymes. *Appl. Microbiol. Biotechnol.* **2007**, *76* (2), 237–248. <https://doi.org/10.1007/s00253-007-1002-0>.
- (11) Hollmann, F.; Opperman, D. J.; Paul, C. E. Biocatalytic Reduction Reactions from a Chemist's Perspective. *Angew. Chem. Int. Ed.* **2021**, *60* (11), 5644–5665. <https://doi.org/10.1002/anie.202001876>.
- (12) Chen, H.; Simoska, O.; Lim, K.; Grattieri, M.; Yuan, M.; Dong, F.; Lee, Y. S.; Beaver, K.; Weliwatte, S.; Gaffney, E. M.; Minter, S. D. Fundamentals, Applications, and Future Directions of Bioelectrocatalysis. *Chem. Rev.* **2020**, *120* (23), 12903–12993. <https://doi.org/10.1021/acs.chemrev.0c00472>.
- (13) Edwardes Moore, E.; Andrei, V.; Zacarias, S.; Pereira, I. A. C.; Reisner, E. Integration of a Hydrogenase in a Lead Halide Perovskite Photoelectrode for Tandem Solar Water Splitting. *ACS Energy Lett.* **2020**, *5* (1), 232–237. <https://doi.org/10.1021/acsenenergylett.9b02437>.
- (14) Sokol, K. P.; Robinson, W. E.; Warnan, J.; Kornienko, N.; Nowaczyk, M. M.; Ruff, A.; Zhang, J. Z.; Reisner, E. Bias-Free Photoelectrochemical Water Splitting with Photosystem II on a Dye-Sensitized Photoanode Wired to Hydrogenase. *Nat. Energy* **2018**, *3* (11), 944–951. <https://doi.org/10.1038/s41560-018-0232-y>.
- (15) Bachmeier, A.; Hall, S.; Ragsdale, S. W.; Armstrong, F. A. Selective Visible-Light-Driven CO<sub>2</sub> Reduction on a p-Type Dye-Sensitized NiO Photocathode. *J. Am. Chem. Soc.* **2014**, *136* (39), 13518–13521. <https://doi.org/10.1021/ja506998b>.
- (16) Edwardes Moore, E.; Andrei, V.; Oliveira, A. R.; Coito, A. M.; Pereira, I. A. C.; Reisner, E. A Semi-artificial Photoelectrochemical Tandem Leaf with a CO<sub>2</sub>-to-Formate Efficiency Approaching 1%. *Angew. Chem. Int. Ed.* **2021**, *60* (50), 26303–26307. <https://doi.org/10.1002/anie.202110867>.
- (17) Sokol, K. P.; Robinson, W. E.; Oliveira, A. R.; Warnan, J.; Nowaczyk, M. M.; Ruff, A.; Pereira, I. A. C.; Reisner, E. Photoreduction of CO<sub>2</sub> with a Formate Dehydrogenase Driven by Photosystem II Using a Semi-Artificial Z-Scheme Architecture. *J. Am. Chem. Soc.* **2018**, *140* (48), 16418–16422. <https://doi.org/10.1021/jacs.8b10247>.
- (18) Kuk, S. K.; Singh, R. K.; Nam, D. H.; Singh, R.; Lee, J.; Park, C. B. Photoelectrochemical Reduction of Carbon Dioxide to Methanol through a Highly Efficient Enzyme Cascade. *Angew. Chem. Int. Ed.* **2017**, *56* (14), 3827–3832. <https://doi.org/10.1002/anie.201611379>.
- (19) Wang, P.; Wang, X.; Chandra, S.; Lielpetere, A.; Quast, T.; Conzuelo, F.; Schuhmann, W. Hybrid Enzyme-Electrocatalyst Cascade Modified Gas-Diffusion Electrodes for Methanol Formation from Carbon Dioxide. *Angew. Chem. Int. Ed.* **2025**, *64* (12), e202422882. <https://doi.org/10.1002/anie.202422882>.
- (20) Zhang, C.; Zhang, H.; Pi, J.; Zhang, L.; Kuhn, A. Bulk Electrocatalytic NADH Cofactor Regeneration with Bipolar Electrochemistry. *Angew. Chem. Int. Ed.* **2022**, *61* (3), e202111804. <https://doi.org/10.1002/anie.202111804>.
- (21) Cheng, B.; Wan, L.; Armstrong, F. A. Progress in Scaling up and Streamlining a Nanoconfined, Enzyme-Catalyzed Electrochemical Nicotinamide Recycling System for Biocatalytic Synthesis. *ChemElectroChem* **2020**, *7* (22), 4672–4678. <https://doi.org/10.1002/celec.202001166>.
- (22) Morello, G.; Megarity, C. F.; Armstrong, F. A. The Power of Electrified Nanoconfinement for Energising, Controlling and Observing Long Enzyme Cascades. *Nat. Commun.* **2021**, *12* (1), 340. <https://doi.org/10.1038/s41467-020-20403-w>.
- (23) Oliveira, A. R.; Mota, C.; Mourato, C.; Domingos, R. M.; Santos, M. F. A.; Gesto, D.; Guigliarelli, B.; Santos-Silva, T.; Romão, M. J.; Cardoso Pereira, I. A. Toward the Mechanistic Understanding of Enzymatic CO<sub>2</sub> Reduction. *ACS Catal.* **2020**, *10* (6), 3844–3856. <https://doi.org/10.1021/acscatal.0c00086>.
- (24) Note that *Nitratidesulfovibrio vulgaris* Hildenborough was formerly named *Desulfovibrio vulgaris* Hildenborough.
- (25) Cobb, S. J.; Badiani, V. M.; Dharani, A. M.; Wagner, A.; Zacarias, S.; Oliveira, A. R.; Pereira, I. A. C.; Reisner, E. Fast CO<sub>2</sub> Hydration Kinetics Impair Heterogeneous but Improve Enzymatic CO<sub>2</sub> Reduction Catalysis. *Nat. Chem.* **2022**, *14* (4), 417–424. <https://doi.org/10.1038/s41557-021-00880-2>.
- (26) Edwardes Moore, E.; Cobb, S. J.; Coito, A. M.; Oliveira, A. R.; Pereira, I. A. C.; Reisner, E. Understanding the Local Chemical Environment of Bioelectrocatalysis. *Proc. Natl. Acad. Sci. U. S. A.* **2022**, *119* (4), e2114097119. <https://doi.org/10.1073/pnas.2114097119>.
- (27) Cobb, S. J.; Dharani, A. M.; Oliveira, A. R.; Pereira, I. A. C.; Reisner, E. Carboxysome-Inspired Electrocatalysis Using Enzymes for the Reduction of CO<sub>2</sub> at Low Concentrations. *Angew. Chem. Int. Ed.* **2023**, *62* (26), e202218782. <https://doi.org/10.1002/anie.202218782>.
- (28) Raynbird, M. Y.; Sampson, J. B.; Smith, D. A.; Forsyth, S. M.; Moseley, J. D.; Wells, A. S. Ketone Reductase Biocatalysis in the Synthesis of Chiral Intermediates Toward Generic Active Pharmaceutical Ingredients. *Org. Process. Res. Dev.* **2020**, *24* (6), 1131–1140. <https://doi.org/10.1021/acs.oprd.0c00120>.
- (29) Jakobblinnert, A.; Mladenov, R.; Paul, A.; Sibilla, F.; Schwaneberg, U.; Ansorge-Schumacher, M. B.; De Maria, P. D. Asymmetric Reduction of Ketones with Recombinant E. Coli Whole Cells in Neat Substrates. *Chem. Commun.* **2011**, *47* (44), 12230–12232. <https://doi.org/10.1039/c1cc14097c>.
- (30) Mersch, D.; Lee, C.-Y.; Zhang, J. Z.; Brinkert, K.; Fontecilla-Camps, J. C.; Rutherford, A. W.; Reisner, E. Wiring of Photosystem II to Hydrogenase for Photoelectrochemical Water Splitting. *J. Am. Chem. Soc.* **2015**, *137* (26), 8541–8549. <https://doi.org/10.1021/jacs.5b03737>.
- (31) Kato, M.; Cardona, T.; Rutherford, A. W.; Reisner, E. Photoelectrochemical Water Oxidation with Photosystem II Integrated in a Mesoporous Indium–Tin Oxide Electrode. *J. Am. Chem. Soc.* **2012**, *134* (20), 8332–8335. <https://doi.org/10.1021/ja301488d>.
- (32) Miller, M.; Robinson, W. E.; Oliveira, A. R.; Heidary, N.; Kornienko, N.; Warnan, J.; Pereira, I. A. C.; Reisner, E. Interfacing Formate Dehydrogenase with Metal Oxides for the Reversible

- Electrocatalysis and Solar-Driven Reduction of Carbon Dioxide. *Angew. Chem. Int. Ed.* **2019**, *58* (14), 4601–4605. <https://doi.org/10.1002/anie.201814419>.
- (33) Mersch, D.; Lee, C.-Y.; Zhang, J. Z.; Brinkert, K.; Fontecilla-Camps, J. C.; Rutherford, A. W.; Reisner, E. Wiring of Photosystem II to Hydrogenase for Photoelectrochemical Water Splitting. *J. Am. Chem. Soc.* **2015**, *137* (26), 8541–8549. <https://doi.org/10.1021/jacs.5b03737>.
- (34) Fang, X.; Sokol, K. P.; Heidary, N.; Kandel, T. A.; Zhang, J. Z.; Reisner, E. Structure–Activity Relationships of Hierarchical Three-Dimensional Electrodes with Photosystem II for Semiartificial Photosynthesis. *Nano Lett.* **2019**, *19* (3), 1844–1850. <https://doi.org/10.1021/acs.nanolett.8b04935>.
- (35) Megarity, C. F.; Siritanaratkul, B.; Cheng, B.; Morello, G.; Wan, L.; Sills, A. J.; Heath, R. S.; Turner, N. J.; Armstrong, F. A. Electrified Nanoconfined Biocatalysis with Rapid Cofactor Recycling. *ChemCatChem* **2019**, *11* (23), 5662–5670. <https://doi.org/10.1002/cctc.201901245>.
- (36) Megarity, C. F.; Weald, T. R. I.; Heath, R. S.; Turner, N. J.; Armstrong, F. A. A Nanoconfined Four-Enzyme Cascade Simultaneously Driven by Electrical and Chemical Energy, with Built-in Rapid, Confocal Recycling of NADP(H) and ATP. *ACS Catal.* **2022**, *12* (15), 8811–8821. <https://doi.org/10.1021/acscatal.2c00999>.
- (37) Wienkamp, R.; Steckhan, E. Indirect Electrochemical Regeneration of NADH by a Bipyridinerhodium(I) Complex as Electron-Transfer Agent. *Angew. Chem., Int. Ed. Engl.* **1982**, *21* (10), 782–783. <https://doi.org/10.1002/anie.198207822>.
- (38) Hildebrand, F.; Kohlmann, C.; Franz, A.; Lütz, S. Synthesis, Characterization and Application of New Rhodium Complexes for Indirect Electrochemical Cofactor Regeneration. *Adv. Synth. Catal.* **2008**, *350* (6), 909–918. <https://doi.org/10.1002/adsc.200700505>.
- (39) Zhang, B.; Xu, S.; He, D.; Chen, R.; He, Y.; Fa, W.; Li, G.; Wang, D. Photoelectrochemical NADH Regeneration Is Highly Sensitive to the Nature of Electrode Surface. *J. Chem. Phys.* **2020**, *153* (6), 064703. <https://doi.org/10.1063/5.0016459>.
- (40) Xing, F.; Xue, X.; Li, J.; Liu, J.; Wang, W.; Dong, W.; Yuan, H.; Liu, J. Sustainable Photocatalytic Biological Cofactor Regeneration Fueled by Selective Alcohol Oxidation over Polarized ZnIn<sub>2</sub>S<sub>4</sub>. *ACS Catal.* **2024**, *14* (15), 11366–11377. <https://doi.org/10.1021/acscatal.4c01703>.
- (41) Chen, M.; Liu, F.; Wu, Y.; Li, Y.; Liu, C.; Zhao, Z.; Zhang, P.; Zhao, Y.; Sun, L.; Li, F. Bioinspired Photoelectrochemical NADH Regeneration Based on a Molecular Catalyst-Modified Photocathode. *Chem. Commun.* **2024**, *60* (24), 3319–3322. <https://doi.org/10.1039/d4cc00448e>.
- (42) Castañeda-Losada, L.; Adam, D.; Paczia, N.; Buesen, D.; Steffler, F.; Sieber, V.; Erb, T. J.; Richter, M.; Plumeré, N. Bioelectrocatalytic Cofactor Regeneration Coupled to CO<sub>2</sub> Fixation in a Redox-Active Hydrogel for Stereoselective C–C Bond Formation. *Angew. Chem. Int. Ed.* **2021**, *60* (38), 21056–21061. <https://doi.org/10.1002/anie.202103634>.
- (43) Lee, D.; Yamauchi, K.; Sakai, K. Water-Induced Switching in Selectivity and Steric Control of Activity in Photochemical CO<sub>2</sub> Reduction Catalyzed by RhCp\*(Bpy) Derivatives. *J. Am. Chem. Soc.* **2024**, *146* (46), 31597–31611. <https://doi.org/10.1021/jacs.4c09486>.
- (44) Wang, C.; Dong, W.; Zhang, P.; Ma, Y.; Han, Z.; Zou, Y.; Wang, W.; Li, H.; Hollmann, F.; Liu, J. Formate-Mediated Electroenzymatic Synthesis via Biological Cofactor NADH. *Angew. Chem. Int. Ed.* **2024**, *63* (41), e202408756. <https://doi.org/10.1002/anie.202408756>.
- (45) Meyer, J.; Romero, M.; Thöming, J.; Baune, M.; Reimer, N.; Dringen, R.; Bösing, I. Experimental Insights into Electrocatalytic [Cp\*Rh(Bpy)Cl]<sup>+</sup> Mediated NADH Regeneration. *Sci. Rep.* **2023**, *13*, 22394. <https://doi.org/10.1038/s41598-023-49021-4>.
- (46) Hollmann, F.; Witholt, B.; Schmid, A. [Cp\*Rh(Bpy)(H<sub>2</sub>O)]<sup>2+</sup>: A Versatile Tool for Efficient and Non-Enzymatic Regeneration of Nicotinamide and Flavin Coenzymes. *J. Mol. Catal. B Enzym.* **2002**, *19–20*, 167–176. [https://doi.org/10.1016/S1381-1177\(02\)00164-9](https://doi.org/10.1016/S1381-1177(02)00164-9).
- (47) Pandey, K.; Islam, S. T. A.; Happe, T.; Armstrong, F. A. Frequency and Potential Dependence of Reversible Electrocatalytic Hydrogen Interconversion by [FeFe]-Hydrogenases. *Proc. Natl. Acad. Sci. U. S. A.* **2017**, *114* (15), 3843–3848. <https://doi.org/10.1073/pnas.1619961114>.
- (48) Liu, Y.; Pulignani, C.; Webb, S.; Cobb, S. J.; Rodríguez-Jiménez, S.; Kim, D.; Milton, R. D.; Reisner, E. Electrostatic [FeFe]-Hydrogenase–Carbon Nitride Assemblies for Efficient Solar Hydrogen Production. *Chem. Sci.* **2024**, *15* (16), 6088–6094. <https://doi.org/10.1039/D4SC00640B>.
- (49) Liu, Y.; Bin Mohamad Annuar, A.; Rodríguez-Jiménez, S.; Yeung, C. W. S.; Wang, Q.; Coito, A. M.; Manuel, R. R.; Pereira, I. A. C.; Reisner, E. Solar Fuel Synthesis Using a Semiartificial Colloidal Z-Scheme. *J. Am. Chem. Soc.* **2024**, *146* (43), 29865–29876. <https://doi.org/10.1021/jacs.4c11827>.
- (50) Yeung, C. W. S.; Liu, Y.; Cobb, S.; Andrei, V.; Coito, A.; Manuel, R.; Pereira, I.; Reisner, E. Semi-Artificial Leaf Interfacing Organic Semiconductors and Enzymes for Solar Fuel Synthesis. ChemRxiv preprint, Submission: 20<sup>th</sup> Dec 2024. DOI: 10.26434/chemrxiv-2024-f49zl (accessed 2025-04-01). <https://doi.org/10.26434/chemrxiv-2024-f49zl>.
- (51) Wiecek, A.; Liu, Y.; Cho, H.-H.; Sivula, K. Assessing the Charge Carrier Dynamics at Hybrid Interfaces of Organic Photoanodes for Solar Fuels. *J. Phys. Chem. Lett.* **2024**, *15* (24), 6347–6354. <https://doi.org/10.1021/acs.jpclett.4c01170>.
- (52) Yeung, C. W. S.; Andrei, V.; Lee, T. H.; Durrant, J. R.; Reisner, E. Organic Semiconductor–BiVO<sub>4</sub> Tandem Devices for Solar-Driven H<sub>2</sub>O and CO<sub>2</sub> Splitting. *Adv. Mater.* **2024**, *36* (35), 2404110. <https://doi.org/10.1002/adma.202404110>.
- (53) Huang, C.; Xiong, P.; Lai, X.-L.; Xu, H.-C. Photoelectrochemical Asymmetric Catalysis. *Nat Catal* **2024**, *7* (12), 1250–1254. <https://doi.org/10.1038/s41929-024-01260-y>.
- (54) Lauterbach, L.; Lenz, O.; Vincent, K. A. H<sub>2</sub>-driven Cofactor Regeneration with NAD(P)<sup>+</sup>-reducing Hydrogenases. *FEBS J.* **2013**, *280* (13), 3058–3068. <https://doi.org/10.1111/febs.12245>.
- (55) Jia, X.; Stewart-Jones, E.; Alvarez-Hernandez, J. L.; Bein, G. P.; Dempsey, J. L.; Donley, C. L.; Hazari, N.; Houck, M. N.; Li, M.; Mayer, J. M.; Nedzbala, H. S.; Powers, R. E. Photoelectrochemical CO<sub>2</sub> Reduction to CO Enabled by a Molecular Catalyst Attached to High-Surface-Area Porous Silicon. *J. Am. Chem. Soc.* **2024**, *146* (12), 7998–8004. <https://doi.org/10.1021/jacs.3c10837>.

---

## ToC artwork:

