
*Genome analysis***SNP Interaction Pattern Identifier (SIPI): An Intensive Search for SNP-SNP Interaction Patterns**

Hui-Yi Lin^{1,*}, Dung-Tsa Chen², Po-Yu Huang³, Yung-Hsin Liu⁴, Augusto Ochoa⁵, Jovanny Zabaleta⁵, Donald E. Mercante¹, Zhide Fang¹, Thomas A. Sellers⁶, Julio Pow-Sang⁷, Chia-Ho Cheng², Rosalind Eeles^{8,9}, Doug Easton¹⁰, Zsofia Kote-Jarai⁸, Ali Amin Al Olama¹⁰, Sara Benlloch¹⁰, Kenneth Muir¹¹, Graham G. Giles^{12,13}, Fredrik Wiklund¹⁴, Henrik Gronberg¹⁴, Christopher A. Haiman¹⁵, Johanna Schleutker^{16,17}, Børge G. Nordestgaard¹⁸, Ruth C. Travis¹⁹, Freddie Hamdy²⁰, Nora Pashayan^{21,38}, Kay-Tee Khaw²², Janet L. Stanford^{23,24}, William J. Blot²⁵, Stephen N. Thibodeau²⁶, Christiane Maier²⁷, Adam S. Kibel^{28,29}, Cezary Cybulski³⁰, Lisa Cannon-Albright³¹, Hermann Brenner^{32,33,39}, Radka Kaneva³⁴, Jyotsna Batra³⁵, Manuel R. Teixeira³⁶, Hardev Pandha³⁷, Yong-Jie Lu⁴⁰, the PRACTICAL consortium⁴¹, and Jong Y. Park⁶

¹ Biostatistics Program, School of Public Health, Louisiana State University Health Sciences Center, New Orleans, LA 70112, USA. ² Department of Biostatistics and Bioinformatics, Moffitt Cancer Center & Research Institute, Tampa, FL 33612, USA. ³ Computational Intelligence Technology Center, Industrial Technology Research Institute, Hsinchu City, Taiwan. ⁴ Department of Biometrics, INC Research, LLC, Raleigh, NC 27609, USA. ⁵ Stanley S. Scott Cancer Center, Louisiana State University Health Sciences Center, New Orleans, LA 70112, USA. ⁶ Department of Cancer Epidemiology, Moffitt Cancer Center & Research Institute, Tampa, FL 33612, USA. ⁷ Department of Genitourinary Oncology, Moffitt Cancer Center & Research Institute, Tampa, FL 33612, USA. ⁸ The Institute of Cancer Research, London, SM2 5NG, UK, ⁹ Royal Marsden NHS Foundation Trust, London, SW3 6JJ, UK, ¹⁰ Centre for Cancer Genetic Epidemiology, Department of Public Health and Primary Care, University of Cambridge, Strangeways Research Laboratory, Worts Causeway, Cambridge, UK, ¹¹ University of Warwick, Coventry, UK, ¹² Cancer Epidemiology Centre, The Cancer Council Victoria, 615 St Kilda Road, Melbourne, Victoria, 3004, Australia, ¹³ Centre for Epidemiology and Biostatistics, Melbourne School of Population and Global Health, The University of Melbourne, Melbourne, Victoria, Australia, ¹⁴ Department of Medical Epidemiology and Biostatistics, Karolinska Institute, Stockholm, Sweden, ¹⁵ Department of Preventive Medicine, Keck School of Medicine, University of Southern California/Norris Comprehensive Cancer Center, Los Angeles, California, USA, ¹⁶ Department of Medical Biochemistry and Genetics, Institute of Biomedicine, Kiinamyllynkatu 10, FI-20014 University of Turku; and Tyks Microbiology and Genetics, Department of Medical Genetics, Turku University Hospital, ¹⁷ BioMediTech, 30014 University of Tampere, Tampere, Finland, ¹⁸ Department of Clinical Biochemistry, Herlev Hospital, Copenhagen University Hospital, Herlev Ringvej 75, DK-2730 Herlev, Denmark, ¹⁹ Cancer Epidemiology, Nuffield Department of Population Health University of Oxford, Oxford, UK, ²⁰ Nuffield Department of Surgical Sciences, University of Oxford, Oxford, UK, Faculty of Medical Science, University of Ox-

ford, John Radcliffe Hospital, Oxford, UK, ²¹ Centre for Cancer Genetic Epidemiology, Department of Oncology, University of Cambridge, Strangeways Research Laboratory, Worts Causeway, Cambridge, UK, ²² Cambridge Institute of Public Health, University of Cambridge, Forvie Site, Robinson Way, Cambridge CB2 0SR, UK, ²³ Division of Public Health Sciences, Fred Hutchinson Cancer Research Center, Seattle, Washington, USA, ²⁴ Department of Epidemiology, School of Public Health, University of Washington, Seattle, Washington, USA, ²⁵ International Epidemiology Institute, 1455 Research Blvd., Suite 550, Rockville, MD 20850, USA, ²⁶ Department of Laboratory Medicine and Pathology, Mayo Clinic, Rochester, MN, USA, ²⁷ Institute of Human Genetics University Hospital Ulm, Germany, ²⁸ Brigham and Women's Hospital/Dana-Farber Cancer Institute, USA, 45 Francis Street- ASB II-3, Boston, MA 02115, USA, ²⁹ Washington University, St Louis, Missouri, USA, ³⁰ International Hereditary Cancer Center, Department of Genetics and Pathology, Pomeranian Medical University, Szczecin, Poland, ³¹ Division of Genetic Epidemiology, Department of Medicine, University of Utah School of Medicine, USA, ³² Division of Clinical Epidemiology and Aging Research, German Cancer Research Center (DKFZ), Heidelberg, Germany, ³³ Division of Preventive Oncology, German Cancer Research Center (DKFZ) and National Center for Tumor Diseases (NCT), Heidelberg, Germany, ³⁴ Molecular Medicine Center and Department of Medical Chemistry and Biochemistry, Medical University - Sofia, 2 Zdrave St, 1431, Sofia, Bulgaria, ³⁵ Australian Prostate Cancer Research Centre-Qld, Institute of Health and Biomedical Innovation and Schools of Life Science and Public Health, Queensland University of Technology, Brisbane, Australia, ³⁶ Department of Genetics, Portuguese Oncology Institute, Porto, Portugal and Biomedical Sciences Institute (ICBAS), Porto University, Porto, Portugal, ³⁷ The University of Surrey, Guildford, Surrey, GU2 7XH, UK, ³⁸ University College London, Department of Applied Health Research, 1-19 Torrington Place, London, WC1E 7HB, UK, ³⁹ German Cancer Consortium (DKTK), German Cancer Research Center (DKFZ), Heidelberg Germany, ⁴⁰ Centre for Molecular Oncology, Barts Cancer Institute, Queen Mary University of London, John Vane Science Centre, Charterhouse Square, London, EC1M 6BQ, UK, ⁴¹ Additional members from the Prostate Cancer Association Group to Investigate Cancer Associated Alterations in the Genome (PRACTICAL) consortium to be provided in the supplement.

*To whom correspondence should be addressed.

Associate Editor: XXXXXXXX

Received on XXXXXX; revised on XXXXXX; accepted on XXXXXX

Abstract

Motivation: Testing SNP-SNP interactions is considered as a key for overcoming bottlenecks of genetic association studies. However, related statistical methods for testing SNP-SNP interactions are underdeveloped. **Results:** We propose the SNP Interaction Pattern Identifier (SIPI), which tests 45 biologically meaningful interaction patterns for a binary outcome. SIPI takes non-hierarchical models, inheritance modes, and mode coding direction into consideration. The simulation results show that SIPI has higher power than MDR (Multifactor Dimensionality Reduction), AA_Full, Geno_Full (full interaction model with additive or genotypic mode) and SNPAssoc in detecting interactions. Applying SIPI to the prostate cancer PRACTICAL consortium data with approximately 21,000 patients, the four SNP pairs in *EGFR-EGFR*, *EGFR-MMP16*, and *EGFR-CSF1* were found to be associated with prostate cancer aggressiveness with the exact or similar pattern in the discovery and validation sets. A similar match for external validation of SNP-SNP interaction studies is suggested. We demonstrated that SIPI not only searches for more meaningful interaction patterns but can also overcome the unstable nature of interaction patterns.

Availability: The SIPI software is freely available at <http://publichealth.lsuhscc.edu/LinSoftware/>.

Contact: hlin1@lsuhsc.edu

Supplementary information: Supplementary figures and tables are available at Bioinformatics online.

1 Introduction

During the past decade, the genome-wide association studies (GWAS) have successfully identified many inherited genetic variants (or SNPs) associated with complex diseases, such as cancer or related phenotypes. However, the predictive power of cancer risk for the GWAS-identified SNPs is small by a median of 1.2 per-allele odds ratio (Ioannidis, et al., 2010). The predictive power of these GWAS SNPs can be improved by combining multiple SNPs in a prediction model (Van den Broeck, et al., 2014). We recently reported the polygenic genetic models to estimate their risk for prostate cancer (Al Olama, et al., 2014; Amin Al Olama, et al., 2015; Eeles, et al., 2013). Despite these efforts, major proportion of familiar risk of prostate cancer remains unknown. The similar situation applies for using SNPs to predict prostate cancer prognosis (Van den Broeck, et al., 2014). It is well known that biological associations among genes are complicated. The majority of GWAS focus on identification of individual SNP effects, which are not sufficient to explain the complexity of disease causality. It has been shown that gene-gene/SNP-SNP interactions play an important role in the etiology of complex diseases (Cordell, 2009; Moore, 2003; Moore and Williams, 2002; Onay, et al., 2006). Although SNP-SNP or gene-gene interaction studies have been emerging, the statistical methods for evaluating SNP-SNP interactions are still underdeveloped.

The majority of genetic association studies focus on two-way interactions with two SNPs involved. In the past decade, various statistical methods have been proposed for evaluating two-way SNP-SNP interactions. These methods can be classified either model-based or pattern-based. The most common model-based approach tests an interaction based on a full interaction model (hierarchical model) with both main effects and their interaction and SNPs could be treated as different inheritance mode (such as additive, dominant, recessive, and genotypic). Examples include the full interaction model in PLINK (Purcell, et al., 2007), SNPAssoc (Gonzalez, et al., 2007) and the 2nd stage of the Boolean Operation-based Screening and Testing (BOOST) (Wan, et al., 2010). For the model-based approaches, the impact of an interaction can be distinguished from the main effects, but the types of detectable interaction patterns are limited. In the pattern-based approach, interaction detection is based on risk patterns of the 3x3 genotype combination table of the two SNPs. The Hypothesis Free Clinical Cloning (HFCC) tests for 255 patterns for one SNP pair (Gayan, et al., 2008), but some patterns may not be biologically meaningful or are rare. SNPmaxsel evaluates 16 interaction patterns and four main effects for a given SNP pair (Boulesteix, et al., 2007). Multifactor dimensionality reduction (MDR) is also a pattern-based approach (Ritchie, et al., 2003; Ritchie, et al., 2001) to define based the best model based on classification accuracy. The strength of pattern-based approaches is that they are designed to detect wider range of interaction patterns. The limitation of the pattern-based approaches is that they search associations that allow for but are not limited to interactions. A significant result detected using the pattern-based approaches may be due to strong main effect without an interaction.

To overcome these weaknesses, we propose SNP Interaction Pattern Identifier (SIPI), which combines the advantages of the model-based and pattern-based approaches. Our approach can examine 45 interaction models that consider biologically meaningful factors. Each model has a straightforward corresponding pattern, and there is a formal statistical test for evaluating the interaction effect. This approach is powerful, and the identified patterns can be easily applied to assemble risk-prediction models. For evaluating the performance of SIPI, we conducted a simulation study to evaluate power and type I errors of SIPI with other four approaches: MDR, AA_full, Geno_Full and SNPAssoc. The details of these methods are listed in Section 2.2.

2 Methods

2.1 SNP Interaction Pattern Identification (SIPI)

SIPI can intensively and effectively search pairwise SNP-SNP interactions. The conventional approach for identifying SNP-SNP interaction is to search a specific type of interaction using the full interaction model with the additive-additive mode based on the minor allele. The SIPI detects 45 interaction models, which take inheritance mode (both original and reverse coding), and risk category grouping (model structure) into consideration. The best interaction pattern is selected based on the Bayesian information criterion (BIC), which is used to deal with the trade-off between model fit and complexity of the model. BIC is also shown to be consistent in selecting the true model and tends to select a parsimonious model compared with the Akaike information criterion (AIC), especially in studies with a large sample size (Yang, 2005). The concept of SIPI can be applied to different types of outcomes, such as numeric, binary and time-to-event variables. In this study, we focused on the binary outcome using logistic regression models. The two primary components of SIPI are introduced separately below.

2.1a SNP Inheritance Modes

The SNP inheritance modes can impact on power to detect SNP interactions (Lin, et al., 2008). We designate a lowercase letter 'a' to denote the minor (low frequency) allele, and an uppercase 'A' to denote the major (common) allele. Each SNP has three genotype categories: homozygous major type ('AA'), heterozygous type ('Aa') and homozygous minor type ('aa'). For a SNP, the inheritance mode for a disease risk refers to a specific relationship between genotype and phenotype. The inheritance modes include additive, dominant, recessive, genotypic and over-dominant modes. The dominant mode assumes that the impact of having one or two copies of a given allele on the outcome is the same, and the recessive mode implies that the subjects with the homozygous genotype of a given allele have a different risk to develop the outcome compared with the other two genotypes. Additive mode refers to the impact of each additional copy of a given allele on the outcome being equal. The genotypic mode, treats a SNP as a categorical variable with three groups, and assumes that each genotype has a distinct effect on risk. This genotypic mode needs four degrees of freedom for the interaction term itself, and interpretation of the result is not straightforward. The over-dominant mode, which assumes that heterozygote has a different risk than the other two homozygous genotypes (Aa vs. AA/aa), is a rare case. Therefore, the SIPI takes three inheritance modes (dominant, recessive and additive) into consideration.

In the majority of genetic association studies, inheritance modes are defined based on the minor (or variant) allele. Under this scenario, the binary inheritance mode (dominant and recessive) is coded as "1" for the group containing the homozygous minor type, and the other group as "0" in modeling. For the AA, Aa and aa genotypes, the additive mode coding is 0, 1 and 2. The reverse coding (=1 - original coding for dominant and recessive mode; and 2-original coding for additive mode) of inheritance mode is seldom to be considered in testing SNP-SNP interactions. The coding direction (original/reverse coding) of inheritance mode does not impact on statistical significance (p-values) for testing the main effect in a main-effect model and the interaction term in a full interaction model, but dramatically impacts testing the interaction term in a non-hierarchical interaction model. As shown in Table 1, there are six total possible coding methods for inheritance modes for each SNP. The three inheritance modes with the original coding based on the minor allele are additive (noted as aSNP1 for SNP1), dominant (dSNP1) and recessive (rSNP1). For the inheritance modes with the reverse coding, the three modes are reverse additive (raSNP1), reverse dominant (rdSNP1), and reverse recessive (rrSNP1).

Table 1. SNP coding scheme of the inheritance modes

SNP1 Maj/Min=A/a	Inheritance mode with original coding ²			Reverse coding ¹		
	Additive (aSNP1)	Dominant (dSNP1)	Recessive (rSNP1)	Reverse Additive (raSNP1)	Reverse Dominant (rdSNP1)	Reverse Recessive (rrSNP1)
AA	0	0	0	2	1	1
Aa	1	1	0	1	0	1
aa	2	1	1	0	0	0
Data type	Continuous	Binary	Binary	Continuous	Binary	Binary

¹Maj/Min= major/minor allele

²Original modes are based on a minor allele 'a'; Reverse coding is (1 - original coding) for the dominant and recessive mode and is (2 - original coding) for the additive mode

2.1b Risk Category Grouping/Model Structure

Both hierarchical and non-hierarchical interaction model were considered in this study. For evaluating 2-way interactions, the hierarchical or full interaction models are the models with two main effects and their interactions. This is the most common model type for testing pairwise SNP-SNP interactions, but this full model tests only one specific interaction pattern. Non-hierarchical models are defined as models with an interaction, and none or one main effects. In genetic association studies, non-hierarchical models, which combines genotypes with similar outcome risk are possible (Lin, et al., 2013; Lin, et al., 2008). Using non-hierarchical models, a parsimonious model based on risk profile can be generated; therefore power of detecting these specific interaction patterns increases (Piegorisch, et al., 1994). As shown in Equations 1-4, four possible model structures for testing a two-way interaction include models with (1) two main effects plus an interaction (Full-int); (2) the main effect of variable 1 plus an interaction (Main1+int); (3) the main effect of variable 2 plus an interaction (Main2+int); and (4) an interaction only (Int-only). It is worthy to note that the interaction only model for a SNP pair does not mean their main effects alone.

By considering a binary inheritance mode, there are four inheritance mode combinations (dominant-dominant, dominant-recessive, recessive-dominant and recessive-recessive). When treating SNPs as numeric variables, the additive-additive mode is taken into consideration. Thus, SIPI considers a total of five possible types of inheritance mode combinations. For each inheritance mode combination, there are nine unique interaction models/patterns when taking into consideration different model structures and inheritance modes (types and original/reverse coding direction). An example of coding direction impact on a recessive-dominant interaction model for a SNP pair in the prostate cancer study is listed in Table S1. Thus, a total of 45 interaction patterns are considered in SIPI for each SNP pair (Table 2).

The best model among the 45 models is based on the lowest value of the Bayesian information criterion (BIC) (Schwarz, 1978). The significance of the interaction effect is tested using the Wald test of the interaction term (H0: $\beta_3=0$). Although the likelihood ratio test (LRT) is usually recommended as the most powerful approach, it requires performing the two models one wishes to compare. The Wald test is similar to LRT in large scale studies and only one model needs to be estimated. In order to ease computation burden for high-dimensional data, the Wald test was primarily used in SIPI. In the SIPI R package, the users can choose to report p-values based on the Wald test or LRT. The Bonferroni method is applied to adjust for multiple comparisons.

Full interaction model (Full-int):

$$\text{logit}[\text{pr}(Y = 1)] = \beta_0 + \beta_1 \text{SNP}_1 + \beta_2 \text{SNP}_2 + \beta_3 \text{SNP}_1 \times \text{SNP}_2 \quad (\text{eq. 1})$$

Main 1+ interaction (Main1+int):

$$\text{logit}[\text{pr}(Y = 1)] = \beta_0 + \beta_1 \text{SNP}_1 + \beta_3 \text{SNP}_1 \times \text{SNP}_2 \quad (\text{eq. 2})$$

Main 2+ interaction (Main2+int):

$$\text{logit}[\text{pr}(Y = 1)] = \beta_0 + \beta_2 \text{SNP}_2 + \beta_3 \text{SNP}_1 \times \text{SNP}_2 \quad (\text{eq. 3})$$

Interaction only (Int-only):

$$\text{logit}[\text{pr}(Y = 1)] = \beta_0 + \beta_3 \text{SNP}_1 \times \text{SNP}_2 \quad (\text{eq. 4})$$

,where Y is the binary outcome with a value of 0 or 1.

Table 2. List of 45 interaction models by considering the inheritance modes and model structures

SNP1x SNP2	Model structure ^c	Model label ^b	Model Details			
Dom-Dom	Full-int	DD_Full	dSNP1 +	dSNP2 +	dSNP1x dSNP2	
		Main1+int	DD_M1_int_o1	dSNP1 +	dSNP1x dSNP2	
	Main2+int	DD_M1_int_r1	rdSNP1 +	rdSNP1x dSNP2		
		DD_M2_int_o2		dSNP2 +	dSNP1x dSNP2	
	Int-only	DD_M2_int_r2		rdSNP2 +	dSNP1x rdSNP2	
		DD_int_oo			dSNP1x dSNP2	
	Dom-Rec	Full-int	DR_Full	dSNP1 +	rSNP2 +	dSNP1x rSNP2
			Main1+int	DR_M1_int_o1	dSNP1 +	dSNP1x rSNP2
		Main2+int	DR_M1_int_r1	rdSNP1 +	rdSNP1x rSNP2	
			DR_M2_int_o2		rSNP2 +	dSNP1x rSNP2
		Int-only	DR_M2_int_r2		rrSNP2 +	dSNP1x rrSNP2
			DR_int_oo			dSNP1x rSNP2
Rec-Dom	Full-int	RD_Full	rSNP1 +	dSNP2 +	rSNP1x dSNP2	
		Main1+int	RD_M1_int_o1	rSNP1 +	rSNP1x dSNP2	
	Main2+int	RD_M1_int_r1	rrSNP1 +	rrSNP1x dSNP2		
		RD_M2_int_o2		dSNP2 +	rSNP1x dSNP2	
	Int-only	RD_M2_int_r2		rdSNP2 +	rSNP1x rdSNP2	
		RD_int_rr			rdSNP1x rrSNP2	

Rec-Rec	Full-int	RR_int_oo	rSNP1 +	rSNP2 +	rSNP1x rSNP2
		RR_int_or	rSNP1 +	rSNP1 +	rSNP1x rSNP2
	Main1+int	RR_M1_int_o1	rrSNP1 +		rrSNP1x rSNP2
		RR_M1_int_r1	rrSNP1 +		rrSNP1x rSNP2
Main2+int	RR_M2_int_o2		rSNP2 +	rSNP1x rSNP2	
	RR_M2_int_r2		rrSNP2 +	rSNP1x rrSNP2	
Int-only	RR_int_oo			rSNP1x rSNP2	
	RR_int_or			rSNP1x rrSNP2	
	RR_int_ro			rrSNP1x rSNP2	
	RR_int_rr			rrSNP1x rrSNP2	
Add_Add	Full-int	AA_Full	aSNP1 +	aSNP2 +	aSNP1x aSNP2
		Main1+int	AA_M1_int_o1	aSNP1 +	aSNP1x aSNP2
	Main2+int	AA_M1_int_r1	raSNP1 +		raSNP1x aSNP2
		AA_M2_int_o2		aSNP2 +	aSNP1x aSNP2
Int-only	AA_M2_int_r2		raSNP2 +	aSNP1x raSNP2	
	AA_int_oo			aSNP1x aSNP2	
	AA_int_or			aSNP1x raSNP2	
	AA_int_ro			raSNP1x aSNP2	
	AA_int_rr			raSNP1x raSNP2	

¹Dom: dominant, Rec: recessive, Add: additive

²Full-int: full interaction model with two main effects plus an interaction; Main1+int: main effect of variable 1 plus an interaction; Main2+int: main effect of variable 2 plus an interaction; and (4) Int-only: an interaction only.

³_o1, _r1: minor allele (original coding), and reverse coding of SNP1

⁴_o2, _r2: minor allele (original coding), and reverse coding of SNP2

⁵_oo, _or, _ro, _rr: based on original-original, original-reverse, reverse-original and reverse-reverse coding for SNP1 and SNP2

2.1c Translating Interaction Models to Interaction Patterns

By treating SNPs as binary variables (such as dominant or recessive), we can simplify genotype combinations from a three-by-three panel into a two-by-two panel, resulting in four possible sub-groupings. For the two-by-two panel, we can categorize the genotype combinations to 4-, 3- and 2-risk subgroups. As shown in Figures S1-S2, we can translate the interaction models to the corresponding genotype interaction patterns. The full-int model has 4 risk subgroups, the Main1+int and Main2+int models have 3 risk subgroups, and the Int-only model have 2 risk subgroups. The non-hierarchical models have flexibility to combine genotype combinations with similar outcome risk. Our 45 model labels are based on a three-by-three table with an order of homozygous major, heterozygous and homozygous minor types (denote as AA, Aa, and aa) and the homozygous major genotypes of the two SNPs as the top left corner.

2.2 Other approaches for SNP-SNP interactions

2.2a MDR

MDR (Ritchie, et al., 2003; Ritchie, et al., 2001) searches overall associations that allow for but are not limited to interactions. A promising MDR generates a binary risk variable (high/low risk) by comparing the case-to-control ratio in each genotype combination to a threshold and classifies each genotype to either a high risk set or low risk set. The best model is decided based on classification accuracy. The K-fold cross-validation is used to relieve over-fitting issue in MDR. The permutation testing (Motsinger-Reif, 2008) can be used to determine MDR overall significance (not just for an interaction). In this study, MDR with the 5-fold cross-validation and a permutation testing procedure (1,000 randomized datasets) was performed. One major weakness of MDR is that its identified associations may be due to strong main effect without an interaction. Thus, another method for the MDR selected interaction is needed to distinguish the impact of main effects and interaction term.

2.2b AA_Full

The AA_full [available in PLINK (Purcell, et al., 2007)] approach uses a logistic regression model with both main effect and interaction. Each SNP is treated as an additive mode based on the minor allele. The significance test is evaluated using the Wald test of the interaction coefficient.

2.2c Geno_Full

The Geno_Full uses a full logistic regression model and each SNP is treated as a genotypic mode with two degrees of freedom. The significance test is evaluated using the likelihood ratio test of the interaction coefficient. This Geno_Full is equivalent to the 2nd stage of BOOST (Wan, et al., 2010), which uses the Kirkwood superposition approximation (KSA) is used to screen a subset of SNPs for the 2nd stage analyses.

2.2d SNPpassoc

SNPpassoc (Gonzalez, et al., 2007) used the same full logistic regression and allows for five different inheritance modes [additive, dominant, recessive, genotypic, and over-dominant (Aa vs. AA/aa)] based on the

SNP Interaction Pattern Identifier (SIPI)

minor allele. Two SNPs in the same pair are required to have the same inheritance mode.

2.3 Simulation

We conducted a simulation study to compare the power of SIPI with the conventional MDR, AA_Full, Geno_Full and SNPAssoc approach for detecting two-way SNP-SNP interactions. For simulation settings, one SNP pair was considered. The two candidate SNPs were generated independently based on the Hardy-Weinberg equilibrium. Seven sets of a wide range of minor allele frequencies (MAF=0.05-0.5) for SNP1 and SNP2 were investigated: (0.5, 0.3), (0.5, 0.2), (0.5, 0.05), (0.3, 0.3), (0.3, 0.1), (0.3, 0.05), and (0.1, 0.05). The sample sizes of 1,000 and 5,000 were chosen. All analyses were based on 1,000 simulation runs.

The binary outcome variable (such as case/control) was generated based on outcome prevalence (such as disease) in each genotype combination of the two given SNPs using multinomial distribution. We evaluated a total of six designed interaction patterns, including one real-data pattern (Figures 1-2). Most of these simulated models are based on the interaction patterns reported previously (Lin, et al., 2013; Lin, et al., 2012). One null model without a true interaction term was also tested. For the effect size of Models 1-4, the outcome prevalence was set to 0.3 or 0.4 in the high-risk subgroups and was 0.2 in the low-risk sub-groups. The corresponding odds ratio (OR) is 1.6 and 2.7, respectively. The settings of true interaction models are listed in Figures 1-2.

Models 1-3 were interaction-only models. For Model 1 (RR_int_rr pattern), both SNPs are considered as recessive with the reverse coding. The disease prevalence is 0.3 and 0.2 for the high- and low-risk groups, respectively. For Model 2 (DD_int_oo pattern), both SNPs are considered as dominant based on the minor alleles. In Model 3 (RD_int_rr), SNP1 is considered under a recessive mode, SNP2 is considered as dominant mode, and both SNPs have the reverse coding. Model 4 (DD_M1_int_o1) includes the SNP1 main effect and an interaction, in which both SNPs are considered as dominant based on the minor allele of SNP1. The significance of the interaction term is the same regardless of the inheritance mode coding (original or reverse) for SNP2. Model 5 (AA_Full) is a full interaction model and both SNPs are treated as an additive mode based on the minor allele. This AA_full model has the setting of $\beta_0 = -2.5$ and $\beta_1 = \beta_2 = \beta_3 = 0.6$ in eq. 1. Model 6 (RD_int_oo) was designed based on rs10488141 and rs6994019 from the PRACTICAL data (first SNP pair in Figure 4) with an OR of 1.9. For the null model, the outcome prevalence of 0.2 was applied for all nine genotype combinations.

2.4 Performance Evaluation

Both power and type I error were evaluated in the 1,000 simulation runs. Power is defined as the percentage of detecting a significant interaction when there is a true interaction. Type I error is defined as percentage of detecting a significant interaction when there is no interaction. The significant tests of the interaction for all four approaches (SIPI, AA_Full, Geno_Full and SNPAssoc) are based on testing the coefficient of the interaction term. Statistical significance for SIPI and SNPAssoc is defined as a $p < 0.001$ ($=0.05/45$) and $p < 0.01$ ($=0.05/5$) based on the Bonferroni correction. For the AA_Full and Geno_Full approaches, the significance level is 0.05. The significance of MDR is based on the permutation p-values (1,000 randomized datasets). In addition, we evaluated SIPI's pattern identification rate, which is defined as the percentage of identified correct interaction pattern among the significant simulation runs.

2.5 Prostate Cancer Study Application

SIPI was applied in evaluating SNP-SNP interactions in angiogenesis genes associated with prostate aggressiveness using Prostate Cancer Association Group to Investigate Cancer Associated Alterations in the Genome (PRACTICAL) consortium data. The study population includes 21,316 cases of European ancestry (3,812 aggressive and 17,504 non-aggressive) from the 32 study sites. We randomly selected half of the cases as the discovery set and the other half as the validation set in each study site. The sample sizes in the discovery and validation sets are 10,664 and 10,652, respectively. Individuals were excluded from the study based on strict quality control criteria including: overall call rate $< 95\%$ and extremely high or low heterozygosity ($p < 1.0 \times 10^{-5}$). Ag-

gressive prostate cancer was defined as a Gleason score > 8 , PSA > 100 , disease stage of "distant" (stage IV) or death from PCa. Ethnic groups were defined based on a subset of 37,000 uncorrelated markers that passed quality control (including $\sim 1,000$ selected as ancestry informative markers). Principal Component Analyses were carried out for the European subgroups. The details of this study population have been published previously (Eeles, et al., 2013).

We evaluated the 148 SNPs in the six angiogenesis genes (*EGFR*, *MMP16*, *ROBO1*, *CSF1*, *FBLN5*, and *HSPG2*), which were reported in a genetic interaction network associated with prostate cancer aggressiveness (Lin, et al., 2013). These result in 10,878 SNP pairs. The pairwise interactions among these SNPs associated with prostate cancer aggressiveness (yes/no) were investigated using the SIPI approach in the discovery set first. For the top SNP pairs identified in the discovery set, both SIPI and AA_Full were conducted in the validation set.

3 Results

3.1 Simulation

The power of comparing SIPI with other four methods (MDR, AA_Full, Geno_Full and SNPAssoc) using the six simulated models for two SNPs with MAF of (0.5, 0.3), (0.5, 0.2) and (0.5, 0.05) is shown in Figures 1-2. As the sample size increased, power of all five approaches increased. In general, SIPI and MDR had similar power, and both of them are more powerful than the other three full-model approaches (AA_Full, Geno_Full, and SNPAssoc). SIPI and MDR suffer less negative impact of SNPs with a low MAF than the other three full-model approaches.

The four methods (SIPI, AA_Full, Geno_Full and SNPAssoc) evaluate the impact of interactions, but MDR is used to test overall associations allow for an interaction. For fair comparisons, we discuss the MDR performance separately. In Models 1-4 for a SNP pair with a $MAF \geq 0.2$ under a sample size of 1,000, SIPI greater than 49% power while the other three approaches have low power ($< 25\%$). Under a sample size of 1,000 with MAF of (0.5, 0.05), power decreases for all three full-model approaches but SIPI still has the highest power. As the sample size increased to 5,000, SIPI has 100% power in most of the conditions for identifying an interaction with a SNP pair with MAF of (0.5, 0.3) and (0.5, 0.2). The order of power for detecting a SNP-SNP interaction is $SIPI > Geno_Full > AA_Full$ (similar with SNPAssoc) with a big sample size of 5,000.

The three full-model approaches (AA_Full, Geno_Full and SNPAssoc) have difficulty detecting Model 1, the "RR_int_rr" pattern. With a recessive interaction-only pattern (RR_int_rr) in Model 1 for a sample size of 1,000 (Figure 1), SIPI has a power of 49-54% but the other three approaches only have a power $< 10\%$. When the sample size increases to 5,000, the power of SIPI is approximately 100% while the other three approaches' power remains low ($< 30\%$).

For Model 2 (DD_int_oo), SIPI have power 58-65%, but the other three approaches only have $< 25\%$ power in a sample size of 1,000 and MAF of (0.5, 0.3) and (0.5, 0.2). As the sample size increases to 5,000, the power of all methods increase, and SIPI has the highest power compared with the other three approaches.

For Model 3 (RD_int_rr), SIPI have the highest power among all testing scenarios in Figure 1. For a sample size of 5,000, SIPI has 100% power, while power of other three approaches is $< 80\%$. Similarly, the power of Model 4 (DD_M1_int_o1) is 59-74% for SIPI and $< 20\%$ for the other three approaches when the sample size is 1,000. Power increases to close to close to 100% for SIPI and 22-60% for others when the sample size becomes 5,000.

For Model 5 (AA_Full), the AA_Full method is the most powerful among all testing approaches in most of the conditions, except the condition of low MAF of (0.5, 0.05) in a sample size of 1,000. Under this special condition, SIPI has the highest power and about 70% of the SIPI significant runs selected similar patterns (AA_M1_int and

AA_M1_int_r). For Model 6 generated based on the real data, SIPI is still the most powerful approach in most of the conditions.

For comparing SIPI with MDR, both methods are more powerful than other three full model approaches (AA_Full, Geno_Full and SNPAssoc). SIPI has similar or higher power compared with MDR in majority of simulated conditions. For Model 6 with high risk groups in the minor genotypes, SIPI is more powerful than MDR under the sample size of 1,000 and 5,000.

For type I errors, SIPI using the Bonferroni correction is the most conservative method among all testing approaches. As shown in Figure 3, SIPI has the smallest type I errors (0.01-0.02) compared to the other three methods. Some of SNPAssoc's type I errors (0.021-0.057) are also less than 0.05. The type I errors for MDR, AA_Full and Geno_Full are close to 0.05. As shown in Tables S2-4, the power and type I error comparisons for additional MAF conditions show similar observations.

3.2 Patten Detection Accuracy

The accuracy rate of pattern identification increases (Figures S4-S5) as the sample size increases. For Models 1, 2, 3 and 6 with 1,000 samples, 56-84% of the significant simulation runs identify the correct pattern. For the sample size of 5,000, all models have approximately 100% accuracy in identifying correct interaction patterns. For Models 4-5 and MAF=(0.3, 0.3) with a sample size of 1,000, the pattern identification rates are low (10% and 2%, respectively). However, these rates become 100% for a sample size of 5,000. Although pattern detection accuracy is low for the smaller sample, SIPI's power can still be high due to detection of other similar patterns. Using Model 4 with MAF=(0.3, 0.3) as an example, only 10% of the significant runs detect the correct pattern (DD_M1_int_oi) but other three similar patterns (39.9% DD_int_oo, 23% DR_int_rr, and 12.6% DR_int_or) are identified (Figure S5). Thus, its power of detecting any interaction can reach 61.2%.

From the simulation results, we observed an interesting scenario for common variants with a MAF close to 0.5. Under this condition, the minor allele determination is unstable, which can affect SIPI's model labels. The model label system are built upon the minor/major allele. As an example shown in Figure S3, a low risk subgroup of a (GG+ GG) combination of SNP1 and SNP2 are classified as the "DD_int_rr" pattern when SNP1 is with a major allele of 'G' and a minor allele of 'A' but is classified as "RD_int_or" (called a "sister pattern") when SNP1's major allele is 'A'. For an interaction with a SNP with a MAF close to 0.5, the pattern identification rate is the sum of the rates of the designed and sister patterns. We present the pattern identification rates for the significant simulation runs in Figures S4-S5. For Model 1 with a SNP pair with MAF=(0.5, 0.3), a total of 74% runs successfully identified the correct risk pattern (39% designed pattern and 5% sister pattern). A similar observations are presented for other models.

3.3 Example of Prostate Cancer Aggressiveness

For the proposed SIPI approach, we considered SNP pairs with a $p < 1 \times 10^{-7}$ to be statistically significant after the Bonferroni correction for 489,510 tests (=10,878 pairs x 45 models per pair). Although the SNP-SNP interaction results do not appear to be significant after adjusting for multiple comparisons, some of them show promising consistent results in both discovery and validation datasets. In the discovery set, 25 SNP pairs had a $p < 0.001$. Among these top 25 pairs, four pairs have a p -value < 0.01 in the validation set. Two pairs (rs10488141+ rs6994019 and rs2058502+ rs4947972) have the exact interaction pattern in both sets. The prevalence of prostate cancer aggressiveness by the nine genotype combinations are shown in Figure 4, and the prediction models are listed in Table 3. The prostate cancer patients with the TT + AC/AA genotype of the SNP pair of EGFR rs10488141 and MMP16 rs6994019 tend to suggest a higher risk of developing aggressive tumors (odds ratio (OR)=1.7, $p=4.5 \times 10^{-6}$). Those with GG+ GG of two SNPs in EGFR (rs2058502 and rs4947972) are less likely to have aggressive prostate cancer tumors (OR=0.8, $p=5.8 \times 10^{-6}$). Those with GG+ AG/AA of two SNPs in EGFR (rs723527 and rs845555) are likely to have aggressive prostate cancer tumors (OR=1.2, $p=3.1 \times 10^{-4}$). The patients with AA/AG and CC in EGFR rs2075110 and CSF1 rs7538029 have a lower chance of developing an aggressive prostate cancer (OR=0.9, $p=2.6 \times 10^{-5}$).

Table 3. Results of the PRACTICAL discovery and validation set for the top 25 SNP-SNP interaction pairs associated with prostate cancer aggressiveness with a $p < 0.001$ in the discovery set

SNP1	SNP2	Pattern	Discovery ¹	Validation ¹		Pattern Similarity ²
			P_d	Pattern	P_v	
rs10228436	rs723527	DR_int_oo	1.0×10^{-4}	DR_int_rr	0.020	
rs13222549	rs16880086	RR_int_oo	2.0×10^{-4}	AA_int_ro	0.378	
rs2017000	rs6981717	DR_int_ro	2.0×10^{-4}	AA_int_oo	0.043	
rs6956366	rs763317	RD_int_oo	2.7×10^{-4}	DR_int_or	0.032	
rs10488141	rs6994019	RD_int_oo	2.8×10^{-4}	RD_int_oo	0.005	same
rs723527	rs845552	RD_int_oo	2.9×10^{-4}	RR_int_oo	0.056	
rs2058502	rs4947972	DD_int_rr	8.9×10^{-4}	RD_int_or	0.002	Same (sister pattern)
rs6548616	rs7780270	DR_int_ro	3.2×10^{-4}	RR_int_ro	0.181	
rs12666347	rs7781264	DR_int_ro	3.6×10^{-4}	DD_int_ro	0.082	
rs2017000	rs723527	DR_int_oo	3.7×10^{-4}	RR_int_rr	0.079	
rs723527	rs845555	RD_int_oo	4.5×10^{-4}	RR_int_rr	0.009	similar
rs16880086	rs6954351	AA_int_ro	4.6×10^{-4}	RR_int_oo	0.123	
rs10228436	rs7780270	DR_int_oo	4.7×10^{-4}	DR_int_ro	0.070	
rs13222549	rs16880099	RD_int_oo	4.9×10^{-4}	AA_int_oo	0.424	
rs10225877	rs16880086	AA_int_oo	5.6×10^{-4}	RD_int_or	0.053	
rs1519938	rs9842630	DD_int_ro	5.9×10^{-4}	RR_int_oo	0.040	
rs13224708	rs17290392	DD_int_oo	6.1×10^{-4}	DR_int_oo	0.943	
rs10488141	rs1879202	RD_int_oo	6.4×10^{-4}	RR_int_oo	0.021	
rs10488141	rs2222294	RD_int_oo	7.3×10^{-4}	DR_int_ro	0.063	
rs2075110	rs7538029	RD_int_rr	7.7×10^{-4}	DD_int_oo	0.007	similar
rs13222549	rs17666091	RD_int_oo	8.7×10^{-4}	DR_int_oo	0.021	
rs11986591	rs6954351	AA_int_ro	9.1×10^{-4}	DR_int_oo	0.138	
rs11977660	rs9842630	DD_int_ro	9.2×10^{-4}	RD_int_ro	0.044	
rs7780270	rs9832396	RD_int_or	9.6×10^{-4}	RR_int_oo	0.191	
rs759169	rs9842630	AA_int_rr	9.8×10^{-4}	AA_int_rr	0.150	

¹ P_d : p-value in the discovery set, P_v : p-value in the validation set, $P_d < 0.001$ and $P_v < 0.01$ were in bold.
² Comparing patterns in the discovery and validation set for the SNP pairs with $P_d < 0.001$ and $P_v < 0.01$

Three of the four SNP interaction pairs remain promising (rs10488141+ rs6994019, rs2058502+ rs4947972, and rs2075110+ rs7538029) after including these four SNP pairs and the first five principal components of European ancestry in the model (Table 4). For evaluating whether the SNPs in the top pairs in the discovery are comparable in the validation set, the MAF of these SNPs are calculated. As shown in Table S5, the MAFs for these top SNPs are very similar in these two datasets. The individual effects of these SNPs in the combined dataset are also evaluated, and some SNPs did not have significant main effects. For example, the SNP pairs of rs10488141 and rs6994019 has an interaction with a p -value of 4.5×10^{-6} but without significant main effects (p -value=0.145 and 0.659, respectively). These show that some pure SNP-SNP interactions (without significant main effects) associated with prostate cancer aggressiveness. In summary, our results demonstrate SNP-SNP interactions in the two gene pairs (EGFR-MMP16 and EGFR-CSF1), and within EGFR. These findings support that EGFR may be the hub of this angiogenesis interaction network, which is consistent with the conclusion of the previous study (Lin, et al., 2013).

4 Discussion

For evaluating two-way SNP-SNP interactions, SIPI is more powerful than the MDR, AA_Full, Geno_Full and SNPAssoc approach, in general, even after applying stringent Bonferroni correction for multiple comparison justification. Although MDR and SIPI have similar power based on our simulation results, SIPI performs better than MDR is terms of testing significance of an interaction. MDR searches overall associations allowing interactions. For testing significance of an interaction, we need a two-stage MDR method, which has lower power than MDR alone. The primary strengths of SIPI are (1) taking non-hierarchical models, inheritance modes, and mode coding direction into consideration and (2) using BIC to search for a best interaction pattern. In practice, it is challenge to detect a true interaction pattern for studies with a limited sample size. These features equip SIPI for searching similar interaction patterns close to the truth, so it can overcome the unstable nature of detecting SNP-SNP interaction patterns.

Our study demonstrated that SIPI is a more comprehensive and flexible tool for detecting two-way SNP-SNP interactions compared with the three full model approaches: AA_Full in PLINK (Purcell, et al., 2007), Geno_Full and SNPAssoc (Gonzalez, et al., 2007). All these methods are based on hierarchical models, and the difference is how they deal with inheritance modes. AA_Full treats SNPs as an additive mode and Geno_Full treats SNPs as categorical variables. SNPAssoc considers five inheritance modes (additive, dominant, recessive, genotypic and over-

SNP Interaction Pattern Identifier (SIPI)

dominant) but two SNPs in a pair need to have the same mode. Thus, these three approaches can only detect limited interaction patterns. For example, AA_Full, Geno_Full, and SNPAssoc experienced difficulty in detecting the RR_int_rr pattern (Model 1, power<30%, Figure 1), but SIPI had 100% power for a large sample size of 5,000.

SIPI also provides advantages compared to other statistical approaches. BOOST (Wan, et al., 2010) is a two-stage method using the log-linear model to test interactions and treats SNPs as the genotypic mode (same as Geno_Full in our study) as the 2nd stage. SIPI is more powerful than Geno_Full (Figures 1 and 2, and Tables S2-S3), which is more powerful than the two-stage BOOST. SNPmaxsel (Boulesteix, et al., 2007) evaluates 16 interaction patterns, which are parts of SIPI patterns. These 16 patterns are the interaction-only models for SNPs with a binary mode (dominant or recessive). HFCC (Gayan, et al., 2008) is used to assess 255 patterns, but some are rare or biologically meaningless patterns. Compared with these approaches, SIPI tests the 45 biologically meaningful patterns, some of which have been reported previously (Lin, et al., 2013).

For external validation of SNP-SNP interactions, we suggest loosening the validation criteria for evaluating SNP-SNP interactions to allow for similar matches. The optimal goal of a genetic association study is to build prediction models for clinical usage. External validation using an independent dataset is a key in identifying true prediction factors. The majority of previous studies use AA_Full in the two independent datasets or the exact interaction pattern identified in the discovery set to verify the same pattern in the validation set (Su, et al., 2013). However, this exact match is too stringent for identifying SNP-SNP interactions. Our simulation findings (Figures S4-S5) indicate the unstable nature of interaction patterns due to unsteady risk profiles of the nine genotype subgroups. Thus, it should be more effective to allow for similar matches instead of exact matches in SNP-SNP interaction validation, especially in the studies with a small sample size. SIPI provides useful features that work to overcome this unstable pattern nature. SIPI uses the BIC to select the best pattern of 45 patterns so that the true pattern or the most similar pattern can be detected. This provides flexibility in terms of result validation. For a SNP pair with MAF of (0.3, 0.3) in Model 4 with a sample size of 1,000, SIPI can still reach 61% power to detect an interaction with any type of SNP1 and SNP2, even though only 10% of the significant results point to the correct pattern.

The outcome prevalence table stratified using three-by-three genotypes (called the “3x3 outcome table”, available in SIPI software) is a useful way to boost result interpretation for interaction patterns. Using the 3x3 outcome table for real prostate cancer data application, it is easy to observe that two of the top SNP pairs had similar interaction patterns in the discovery set and validation set (Figure 4). Combining the two testing sets with a larger sample size ensures that the interaction pattern is more reliable. In result validation, the sister pattern (one pattern with two different pattern labels) can be easily observed for an interaction with a SNP with a MAF close to 50%. In our prostate cancer application, three out of eight SNPs involved in the top SNP interactions have a MAF>45%. In practice, the sister pattern issue can be identified by reviewing the 3x3 outcome table. Thus, we cannot purely rely on model labels to decide whether the two patterns are exactly matched. Due to the sister pattern and similar matching issues, it is suggested that the 3x3 outcome table should be consulted to further review interaction patterns.

For potential biological relevance of our identified SNP-SNP interactions (within *EGFR*, *EGFR-MMP16* and *EGFR-CSF1*), the main key protein

was epidermal growth factor receptor (EGFR), which interacted with the other two proteins that were also involved in cancers. The EGFR is a critical protein in proliferation of epithelial cells, differentiation, and cell survival and is involved in oncogenesis. The EGFR has been known for a role in regulating prostate cellular growth and function (Bonaccorsi, et al., 2007; Leotoing, et al., 2007; Migliaccio, et al., 2006). Results from a meta-analysis of prostate cancer expression datasets were consistent with our results. Wang et al. identified the EGFR pathway, which was associated with prostate cancer risk (Wang, et al., 2011).


The interaction between matrix metalloproteinase16 (MMP16) and EGFR is interesting. These two proteins have also been implicated in several cancers including prostate cancer. MMPs are a family of proteolytic enzymes involved in tumor growth, invasion and metastasis (Rundhaug, 2005). Among 24 MMPs, the role of MMP16 in prostate cancer has not been well investigated. Jung et al. reported a down-regulation of MMP16 in malignant prostate tissues (Jung, et al., 2003). MMP16 has been shown to be associated with pancreatic cancer cell migration and invasion (Lin, et al., 2011) and lung development (Hadchouel, et al., 2008). Several cancers in which EGFR signaling is involved were often observed abnormal high expression of MMPs (Davidson, et al., 1999). Van Meter et al. reported MMP16 mRNA levels significantly increased after EGF stimulation in the glioma cell lines (Van Meter, et al., 2004).

Colony stimulating factor-1 (CSF1) is a protein that promotes metastatic potential in breast cancer (Lin, et al., 2001). Although there is no report on a role of CSF1 in prostate cancer, previous studies reported overexpression of serum CSF1 in several cancer sites, including pancreatic cancer (Pyonteck, et al., 2012), breast, ovary, and endometrial tissues (Espinosa, et al., 2011; Kacinski, 1997). Recently, Pei et al (Pei, et al., 2015) observed that CSF1 expression is positively correlated with progression and EGFR expression in lung cancer and concluded that co-expression of CSF1 and EGFR may be an independent prognostic biomarker for progression of lung cancer.

The SIPI software (SAS macro and SIPI R package) is freely available at <http://publichealth.lsuhsu.edu/LinSoftware/>. SIPI software can perform models adjusted for covariates. In SIPI R package, the original (‘SIPI’ function) and parallel computing functions (‘parSIPI’ function) are included. SIPI can finish the analyses of all pairwise analyses of 150 SNPs for a dataset with a sample size of 5,000 within about 3 hours on a desktop computer (3.6 GHz CPU with 8 cores) using the ‘parSIPI’ R function. For large scale studies, it is recommended to apply some approaches (such as statistical screening or biological pathway selection) to decrease the number of candidate SNPs before applying the SIPI analyses.


In summary, SIPI is a powerful tool to search for 45 interaction patterns for pairwise SNP interactions. Although only binary outcome models were discussed in this study, it can be extended to various outcome data types, such as numeric and time-to-event data. The promising interaction pairs identified by SIPI can be included in a risk prediction model with other significant individual SNPs, other known clinical risk factors, and biomarkers in order to increase prediction accuracy.

Figure 1. Power comparisons of the SNP Interaction Pattern Identifier (SIPI) and other four methods for Models 1-3



¹Proportion of the outcome event in the genotype combination of the 3x3 table; a lowercase letter denotes the minor allele, and an uppercase letter denotes the major allele. ²MDR (Multifactor dimensionality reduction), AA_Full, Geno_Full (full interaction model and each SNP is treated as an additive or genotypic mode), and SNPAssoc R package


Figure 2. Power comparisons of the SNP Interaction Pattern Identifier (SIPI) and other four methods for Models 4-6



¹Proportion of the outcome event in the genotype combination of the 3x3 table; a lowercase letter denotes the minor allele, and an uppercase letter denotes the major allele. ²MDR (Multifactor dimensionality reduction), AA_Full, Geno_Full (full interaction model and each SNP is treated as an additive or genotypic mode), and SNPAssoc R package

SNP Interaction Pattern Identifier (SIPI)

Figure 3. Comparisons of Type I errors of the SNP Interaction Pattern Identifier (SIPI) and other four methods



¹Proportion of the outcome event in the genotype combination of the 3x3 table; a lowercase letter denotes the minor allele, and an uppercase letter denotes the major allele. ²MDR (Multifactor dimensionality reduction), AA_Full, Geno_Full (full interaction model and each SNP is treated as an additive or genotypic mode), and SNPAssoc R package

Figure 4. Proportions of prostate cancer aggressiveness by genotype combinations for the top four SNP-SNP interaction pairs associated with prostate cancer aggressiveness by datasets

Pair	Discovery: Pr(aggr) ¹			Validation: Pr(aggr) ¹			Combined: Pr(aggr) ¹		
	model	p-value		model	p-value		model	p-value	
1	EGFR: rs10488141	MMP16: rs6994019		EGFR: rs10488141	MMP16: rs6994019		EGFR: rs10488141	MMP16: rs6994019	
	CC	AC	AA	CC	AC	AA	CC	AC	AA
	AA	0.19	0.18	0.17	0.17	0.20	0.18	0.18	0.18
	AT	0.19	0.18	0.14	0.18	0.16	0.19	0.17	0.15
	TT	0.14	0.28	0.34	0.13	0.25	0.22	0.14	0.27
	RD_int_oo	2.8x10 ⁻⁴		RD_int_oo	0.005		RD_int_oo	4.5x10 ⁻⁶	
	AA_Full	0.047		AA_Full	0.842		AA_Full	0.192	
2	EGFR: rs2058502	EGFR: rs4947972		EGFR: rs2058502	EGFR: rs4947972		EGFR: rs2058502	EGFR: rs4947972	
	GG	CG	CC	GG	CG	CC	GG	CG	CC
	GG	0.16	0.20	0.23	0.14	0.18	0.20	0.15	0.19
	AG	0.19	0.20	0.21	0.18	0.17	0.18	0.18	0.19
	AA	0.20	0.18	0.17	0.18	0.15	0.21	0.19	0.19
	DD_int_rr	8.9x10 ⁻⁴		RD_int_or ²	0.002		DD_int_rr	5.8x10 ⁻⁶	
	AA_full	2.3x10 ⁻⁴		AA_full	0.023		AA_full	2.4x10 ⁻⁵	
3	EGFR: rs723527	EGFR: rs845555		EGFR: rs723527	EGFR: rs845555		EGFR: rs723527	EGFR: rs845555	
	GG	AG	AA	GG	AG	AA	GG	AG	AA
	AA	0.19	0.18	0.18	0.17	0.17	0.18	0.18	0.18
	AG	0.17	0.19	0.18	0.16	0.16	0.18	0.17	0.18
	GG	0.17	0.22	0.22	0.19	0.18	0.20	0.18	0.21
	RD_int_oo	4.5x10 ⁻⁴		RR_int_rr	0.009		RD_int_oo	3.1x10 ⁻⁴	
	AA_full	0.034		AA_full	0.776		AA_Full	0.186	
4	EGFR: rs2075110	CSF1: rs7538029		EGFR: rs2075110	CSF1: rs7538029		EGFR: rs2075110	CSF1: rs7538029	
	CC	CA	AA	CC	CA	AA	CC	CA	AA
	AA	0.17	0.20	0.18	0.17	0.18	0.20	0.19	0.19
	AG	0.17	0.20	0.22	0.16	0.18	0.23	0.17	0.23
	GG	0.21	0.17	0.17	0.17	0.19	0.17	0.19	0.17
	RD_int_rr	7.7x10 ⁻⁴		DD_int_oo	0.007		RD_int_rr	2.6x10 ⁻⁵	
	AA_full	0.023		AA_full	0.834		AA_full	0.138	

¹ Pr(aggr): Values in the 3x3 table are proportions of aggressive prostate cancer (=number of aggressive PCa patients/ number of PCa patients)
² RD_int_or in the validation set indicated the same pattern as DD_int_rr in the discovery set. The different pattern name is due to the revise issue of the major and minor allele in the validation set. rs2058502 (minor < major allele): (A<G) in discovery set, and (G<A) in validation set.

Table 4. SNP-SNP interaction models associated with prostate cancer aggressiveness

	Univariate model		Multivariable model ²	
	Unadjusted	p-value	adjusted	p-value
	OR (95% CI) ¹		OR (95% CI) ¹	
rs10488141+ rs6994019, TT+ AC/AA vs. others	1.7 (1.4-2.1)	4.5x10 ⁻⁶	1.8 (1.4-2.6)	6.3x10 ⁻⁷
rs2058502+ rs4947972, GG+ GG vs. others	0.8 (0.7-0.9)	5.8x10 ⁻⁶	0.8 (0.7-0.9)	5.2x10 ⁻⁵
rs723527+ rs845555, GG+ AG/AA vs. others	1.2 (1.1-1.3)	3.1x10 ⁻⁴	1.1 (1.0-1.3)	1.6x10 ⁻²
rs2075110+ rs7538029, AA/AG+ CC vs. others	0.9 (0.8-0.9)	2.6x10 ⁻⁵	0.9 (0.8-0.9)	6.9x10 ⁻⁴

¹Odds ratio (95% confidence interval)

² all four SNP pairs and the first five principal components of European ancestry were included in the multivariable model.

Acknowledgements

H. Lin et al.

We thank our anonymous reviewers for their valuable comments, which have led to many improvements to this article. This study was supported by the National Cancer Institute (R01CA128813, PI: Park, JY and R21CA202417, PI: Lin, HY).

Conflict of Interest: none declared.

References

- Al Olama, A.A., et al. A meta-analysis of 87,040 individuals identifies 23 new susceptibility loci for prostate cancer. *Nat Genet* 2014;46(10):1103-1109.
- Amin Al Olama, A., et al. Risk Analysis of Prostate Cancer in PRACTICAL, a Multinational Consortium, Using 25 Known Prostate Cancer Susceptibility Loci. *Cancer Epidemiol Biomarkers Prev* 2015;24(7):1121-1129.
- Bonaccorsi, L., et al. Altered endocytosis of epidermal growth factor receptor in androgen receptor positive prostate cancer cell lines. *J Mol Endocrinol* 2007;38(1-2):51-66.
- Boulesteix, A.L., et al. Multiple testing for SNP-SNP interactions. *Stat. Appl. Genet. Mol. Biol.* 2007;6:Article37.
- Cordell, H.J. Detecting gene-gene interactions that underlie human diseases. *Nat. Rev. Genet.* 2009;10(6):392-404.
- Davidson, B., et al. High levels of MMP-2, MMP-9, MT1-MMP and TIMP-2 mRNA correlate with poor survival in ovarian carcinoma. *Clin Exp Metastasis* 1999;17(10):799-808.
- Eeles, R.A., et al. Identification of 23 new prostate cancer susceptibility loci using the iCOGS custom genotyping array. *Nat Genet* 2013;45(4):385-391, 391e381-382.
- Espinosa, I., et al. CSF1 expression in nongynecological leiomyosarcoma is associated with increased tumor angiogenesis. *Am J Pathol* 2011;179(4):2100-2107.
- Gayan, J., et al. A method for detecting epistasis in genome-wide studies using case-control multi-locus association analysis. *BMC Genomics* 2008;9:360.
- Gonzalez, J.R., et al. SNPassoc: an R package to perform whole genome association studies. *Bioinformatics* 2007;23(5):644-645.
- Hadchouel, A., et al. Matrix metalloproteinase gene polymorphisms and bronchopulmonary dysplasia: identification of MMP16 as a new player in lung development. *PLoS One* 2008;3(9):e3188.
- Ioannidis, J.P., Castaldi, P. and Evangelou, E. A compendium of genome-wide associations for cancer: critical synopsis and reappraisal. *J. Natl. Cancer Inst.* 2010;102(12):846-858.
- Jung, M., et al. mRNA expression of the five membrane-type matrix metalloproteinases MT1-MT5 in human prostatic cell lines and their down-regulation in human malignant prostatic tissue. *Prostate* 2003;55(2):89-98.
- Kacinski, B.M. CSF-1 and its receptor in breast carcinomas and neoplasms of the female reproductive tract. *Mol Reprod Dev* 1997;46(1):71-74.
- Leotoing, L., et al. Crosstalk between androgen receptor and epidermal growth factor receptor-signalling pathways: a molecular switch for epithelial cell differentiation. *J Mol Endocrinol* 2007;39(2):151-162.
- Lin, E.Y., et al. Colony-stimulating factor 1 promotes progression of mammary tumors to malignancy. *J Exp Med* 2001;193(6):727-740.
- Lin, F., et al. Inhibitory effects of miR-146b-5p on cell migration and invasion of pancreatic cancer by targeting MMP16. *J Huazhong Univ Sci Technol Med Sci* 2011;31(4):509-514.
- Lin, H.Y., et al. SNP-SNP Interaction Network in Angiogenesis Genes Associated with Prostate Cancer Aggressiveness. *PLoS ONE* 2013;8(4):e59688.
- Lin, H.Y., et al. TRM: a powerful two-stage machine learning approach for identifying SNP-SNP interactions. *Annals of human genetics* 2012;76(1):53-62.
- Lin, H.Y., et al. Comparison of multivariate adaptive regression splines and logistic regression in detecting SNP-SNP interactions and their application in prostate cancer. *Journal of Human Genetics* 2008;53(9):802-811.
- Migliaccio, A., et al. Crosstalk between EGFR and extranuclear steroid receptors. *Ann N Y Acad Sci* 2006;1089:194-200.
- Moore, J.H. The ubiquitous nature of epistasis in determining susceptibility to common human diseases. *Hum. Hered.* 2003;56(1-3):73-82.
- Moore, J.H. and Williams, S.M. New strategies for identifying gene-gene interactions in hypertension. *Annals of medicine* 2002;34(2):88-95.
- Motsinger-Reif, A.A. The effect of alternative permutation testing strategies on the performance of multifactor dimensionality reduction. *BMC Res. Notes* 2008;1:139.
- Onay, V.U., et al. SNP-SNP interactions in breast cancer susceptibility. *BMC Cancer* 2006;6:114.
- Pei, B., et al. [Expression of colony-stimulating factor 1 in lung adenocarcinoma and its prognostic implication]. *Zhonghua Zhong Liu Za Zhi* 2015;37(2):113-118.
- Piegorsch, W.W., Weinberg, C.R. and Taylor, J.A. Non-hierarchical logistic models and case-only designs for assessing susceptibility in population-based case-control studies. *Stat. Med.* 1994;13(2):153-162.
- Purcell, S., et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am. J. Hum. Genet.* 2007;81(3):559-575.
- Pyonteck, S.M., et al. Deficiency of the macrophage growth factor CSF-1 disrupts pancreatic neuroendocrine tumor development. *Oncogene* 2012;31(11):1459-1467.
- Ritchie, M.D., Hahn, L.W. and Moore, J.H. Power of multifactor dimensionality reduction for detecting gene-gene interactions in the presence of genotyping error, missing data, phenocopy, and genetic heterogeneity. *Genet. Epidemiol.* 2003;24(2):150-157.
- Ritchie, M.D., et al. Multifactor-dimensionality reduction reveals high-order interactions among estrogen-metabolism genes in sporadic breast cancer. *Am J Hum Genet* 2001;69(1):138-147.
- Rundhaug, J.E. Matrix metalloproteinases and angiogenesis. *J Cell Mol Med* 2005;9(2):267-285.
- Schwarz, G. Estimating the Dimension of a Model. *Annals of Statistics* 1978;6(2):461-464.
- Su, W.H., et al. How genome-wide SNP-SNP interactions relate to nasopharyngeal carcinoma susceptibility. *PLoS ONE* 2013;8(12):e83034.
- Van den Broeck, T., et al. The role of single nucleotide polymorphisms in predicting prostate cancer risk and therapeutic decision making. *Biomed Res Int* 2014;2014:627510.
- Van Meter, T.E., et al. Induction of membrane-type-1 matrix metalloproteinase by epidermal growth factor-mediated signaling in gliomas. *Neuro Oncol* 2004;6(3):188-199.
- Wan, X., et al. BOOST: A fast approach to detecting gene-gene interactions in genome-wide case-control studies. *Am. J. Hum. Genet.* 2010;87(3):325-340.
- Wang, Y., et al. Identifying novel prostate cancer associated pathways based on integrative microarray data analysis. *Comput Biol Chem* 2011;35(3):151-158.
- Yang, Y. Can the strengths of AIC and BIC be shared? A conflict between model identification and regression estimation. *Biometrika* 2005;92(4):937-950.