

ORIGINAL ARTICLE

The functional *ALDH2* polymorphism is associated with breast cancer risk: A pooled analysis from the Breast Cancer Association Consortium

Tomotaka Ugai¹ | Roger L. Milne^{2,3} | Hidemi Ito^{4,5}  | Kristan J. Aronson⁶ |
 Manjeet K. Bolla⁷ | Tsun Chan^{8,9} | Ching W. Chan¹⁰ | Ji-Yeob Choi^{11,12} |
 Don M. Conroy¹³ | Joe Dennis⁷ | Alison M. Dunning¹³ | Douglas F. Easton^{7,13} |
 Valerie Gaborieau¹⁴ | Anna Gonzalez-Neira¹⁵ | Mikael Hartman^{10,16} |
 Catherine S. Healey¹³ | Motoki Iwasaki¹⁷ | Esther M. John¹⁸ | Daehee Kang^{11,12,19} |
 Sung-Won Kim²⁰ | Ava Kwong^{8,21,22} | Artitaya Lophatananon^{23,24} |
 Kyriaki Michailidou^{7,25} | Nur Aishah Mohd Taib²⁶ | Kenneth Muir^{23,24} | Sue K. Park²⁷ |
 Paul D. P. Pharoah^{7,11} | Suleeporn Sangrajrang²⁸ | Chen-Yang Shen^{29,30} | Xiao-Ou Shu³¹ |
 John J. Spinelli^{32,33} | Soo H. Teo^{26,34} | Daniel C. Tessier³⁵ | Chiu-Chen Tseng³⁶ |
 Shoichiro Tsugane¹⁷ | Daniel Vincent³⁵ | Qin Wang⁷ | Anna H. Wu³⁶ | Pei-Ei Wu²⁹ |
 Wei Zheng³¹ | Keitaro Matsuo^{1,5} 

¹Division of Cancer Epidemiology and Prevention, Department of Preventive Medicine, Aichi Cancer Center Research Institute, Nagoya, Japan

²Cancer Epidemiology & Intelligence Division, Melbourne, VIC, Australia

³Centre for Epidemiology and Biostatistics, School of Population and Global Health, The University of Melbourne, Melbourne, VIC, Australia

⁴Division of Cancer Information and Control, Department of Preventive Medicine, Aichi Cancer Center Research Institute, Nagoya, Japan

⁵Department of Epidemiology, Nagoya University Graduate School of Medicine, Nagoya, Japan

⁶Department of Public Health Sciences, Queen's Cancer Institute, Queen's University, Kingston, Ontario, Canada

⁷Centre for Cancer Genetic Epidemiology, Department of Public Health and Primary Care, University of Cambridge, Cambridge, UK

⁸Hong Kong Hereditary Breast Cancer Family Registry, Happy Valley, Hong Kong

⁹Department of Pathology, Hong Kong Sanatorium and Hospital, Happy Valley, Hong Kong

¹⁰Department of Surgery, National University Health System, Singapore

¹¹Department of Biomedical Sciences, Seoul National University College of Medicine, Seoul, Korea

¹²Cancer Research Institute, Seoul National University College of Medicine, Seoul, Korea

¹³Centre for Cancer Genetic Epidemiology, Department of Oncology, University of Cambridge, Cambridge, UK

¹⁴Genetic Epidemiology Group, International Agency for Research on Cancer, Lyon, France

¹⁵Human Cancer Genetics Program, Spanish National Cancer Research Centre, Madrid, Spain

¹⁶Saw Swee Hock School of Public Health, National University of Singapore, Singapore

¹⁷Epidemiology and Prevention Group, Center for Public Health Sciences, National Cancer Center, Tokyo, Japan

¹⁸Department of Medicine and Stanford Cancer Institute, Stanford University School of Medicine, Stanford, California, USA

¹⁹Department of Preventive Medicine, Seoul National University College of Medicine, Seoul National University, Seoul, Korea

²⁰Department of Surgery, Daerim Saint Mary's Hospital, Seoul, Korea

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2019 The Authors. *Molecular Genetics & Genomic Medicine* published by Wiley Periodicals, Inc.

²¹Department of Surgery, Queen Mary Hospital, The University of Hong Kong, Happy Valley, Hong Kong

²²Department of Surgery, Hong Kong Sanatorium and Hospital, Happy Valley, Hong Kong

²³Division of Health Sciences, Warwick Medical School, Warwick University, Coventry, UK

²⁴Division of Population Sciences, Warwick Medical School, Warwick University, Coventry, UK

²⁵Department of Electron Microscopy/Molecular Pathology, The Cyprus Institute of Neurology and Genetics, Nicosia, Cyprus

²⁶Breast Cancer Research Unit, University Malaya Cancer Research Institute, University Malaya Medical Centre, Kuala Lumpur, Malaysia

²⁷Department of Preventive Medicine, Seoul National University College of Medicine, Seoul, Korea

²⁸National Cancer Institute, Bangkok, Thailand

²⁹Taiwan Biobank, Institute of Biomedical Sciences, Academia Sinica, Taipei, Taiwan

³⁰College of Public Health, China Medical University, Taichong, Taiwan

³¹Division of Epidemiology, Department of Medicine, Vanderbilt Epidemiology Center, Vanderbilt-Ingram Cancer Center, Vanderbilt University School of Medicine, Nashville, Tennessee, USA

³²School of Population & Public Health, University of British Columbia, Vancouver, British Columbia, Canada

³³Cancer Control Research, BC Cancer Agency, Vancouver, British Columbia, Canada

³⁴Cancer Research Initiatives Foundation, Sime Darby Medical Centre, Subang Jaya, Malaysia

³⁵McGill University and Génome Québec Innovation Centre, Montreal, Quebec, Canada

³⁶Department of Preventive Medicine, Keck School of Medicine, University of Southern California, Los Angeles, California, USA

Correspondence

Keitaro Matsuo, Division of Cancer Epidemiology and Prevention, Department of Preventive Medicine, Aichi Cancer Center Research Institute, 1-1 Kanokoden, Chikusa-ku, Nagoya 464-8681, Japan.

Email: kmatsuo@aichi-cc.jp

Funding information

BCAC is funded by Cancer Research UK [C1287/A16563, C1287/A10118], the European Union's Horizon 2020 Research and Innovation Programme (grant numbers 634935 and 633784 for BRIDGES and B-CAST respectively), and by the European Community's Seventh Framework Programme under grant agreement number 223175 (grant number HEALTH-F2-2009-223175) (COGS). The EU Horizon 2020 Research and Innovation Programme funding source had no role in study design, data collection, data analysis, data interpretation, or writing of the report. Genotyping of the OncoArray was funded by the NIH Grant U19 CA148065, and Cancer UK Grant C1287/A16563 and the PERSPECTIVE project supported by the Government of Canada through Genome Canada and the Canadian Institutes of Health Research (grant GPH-129344) and, the Ministère de l'Économie, Science et Innovation du Québec through Genome Québec and the PERSIIRI-701 grant, and the Quebec Breast Cancer Foundation. Funding for the iCOGS infrastructure came from: the European Community's Seventh Framework Programme under grant agreement n° 223175 (HEALTH-F2-2009-223175) (COGS), Cancer Research UK (C1287/A10118, C1287/A10710, C12292/A11174, C1281/A12014, C5047/A8384, C5047/A15007, C5047/A10692, C8197/A16565), the National Institutes of Health (CA128978) and Post-Cancer GWAS initiative (U19 CA148537, U19 CA148065 and U19 CA148112—the GAME-ON initiative), the Department of Defence (W81XWH-10-1-0341), the Canadian Institutes of Health Research (CIHR) for the CIHR Team in Familial Risks of Breast Cancer, and Komen Foundation for the Cure, the Breast Cancer Research Foundation, and the Ovarian Cancer Research Fund. The ACP study is funded by the Breast Cancer Research Trust, UK. CBCS is funded by the Canadian Cancer Society (grant # 313404) and the Canadian Institutes of Health Research. The HERPACC was supported by MEXT Kakenhi (No. 170150181 and 26253041) from the Ministry of Education, Science, Sports, Culture, and Technology of Japan, by a Grant-in-Aid for the Third Term Comprehensive 10-Year Strategy for Cancer Control from Ministry Health, Labour, and Welfare of Japan, by Health and Labour Sciences Research Grants for Research on Applying Health Technology from Ministry Health, Labour, and Welfare of Japan, by National Cancer Center Research and Development Fund, and "Practical Research for Innovative Cancer Control (15ck0106177h0001)" from Japan Agency for Medical Research and development, AMED, and Cancer Bio Bank Aichi. The KOHBRA study was partially supported by a grant from the Korea Health Technology R&D Project through the Korea Health Industry Development Institute (KHIDI), and the National R&D Program for Cancer Control, Ministry of Health & Welfare, Republic of Korea (HI16C1127; 1020350; 1420190). LAABC is supported by grants (1RB-0287, 3PB-0102, 5PB-0018, 10PB-0098) from the California Breast Cancer Research Program. MYBRCA is funded by research grants from the Malaysian Ministry of Higher Education (UM.C/HIR/MOHE/06) and Cancer Research Malaysia. MYMAMMO is supported by research grants from Yayasan Sime Darby LPGA Tournament and Malaysian Ministry of Higher Education (RP046B-15HTM). The Northern California Breast Cancer Family Registry (NC-BCFR) was supported by grant UMI CA164920 from the National Cancer Institute (USA). The content of this manuscript does not necessarily reflect the views or policies of the National Cancer Institute or any of the collaborating centers in the Breast Cancer Family Registry (BCFR), nor does mention of trade names, commercial products, or organizations imply endorsement by the USA Government or the BCFR. The NCOBCS was supported by National Cancer Center Research and Development Fund. The SBCGS was supported primarily by NIH grants R01CA64277, R01CA148667, UMCA182910, and R37CA70867. Biological sample preparation was conducted the Survey and Biospecimen Shared Resource, which is supported by P30 CA68485. The scientific development and funding of this project were, in part, supported by the Genetic Associations and Mechanisms in Oncology (GAME-ON) Network U19 CA148065. SEBCS was supported by the BRL (Basic Research Laboratory) program through the National Research Foundation of Korea funded by the Ministry of Education, Science and Technology (2012-0000347). SGBCC is funded by the NUS start-up Grant, National University Cancer Institute Singapore (NCIS) Centre Grant and the NMRC Clinician Scientist Award. Additional controls were recruited by the Singapore Consortium of Cohort Studies-Multi-ethnic cohort (SCCS-MEC), which was funded by the Biomedical Research Council, grant number: 05/1/21/19/425. The TBCS was funded by The National Cancer Institute Thailand. The TWBCS is supported by the Institute of Biomedical Sciences, Academia Sinica, Taiwan.

Abstract

Background: Epidemiological studies consistently indicate that alcohol consumption is an independent risk factor for female breast cancer (BC). Although the aldehyde dehydrogenase 2 (*ALDH2*) polymorphism (rs671: Glu>Lys) has a strong effect on acetaldehyde metabolism, the association of rs671 with BC risk and its interaction with alcohol intake have not been fully elucidated. We conducted a pooled analysis of 14 case-control studies, with individual data on Asian ancestry women participating in the Breast Cancer Association Consortium.

Methods: We included 12,595 invasive BC cases and 12,884 controls for the analysis of rs671 and BC risk, and 2,849 invasive BC cases and 3,680 controls for the analysis of the gene-environment interaction between rs671 and alcohol intake for BC risk. The pooled odds ratios (OR) with 95% confidence intervals (CI) associated with rs671 and its interaction with alcohol intake for BC risk were estimated using logistic regression models.

Results: The Lys/Lys genotype of rs671 was associated with increased BC risk (OR = 1.16, 95% CI 1.03–1.30, $p = 0.014$). According to tumor characteristics, the Lys/Lys genotype was associated with estrogen receptor (ER)-positive BC (OR = 1.19, 95% CI 1.05–1.36, $p = 0.008$), progesterone receptor (PR)-positive BC (OR = 1.19, 95% CI 1.03–1.36, $p = 0.015$), and human epidermal growth factor receptor 2 (HER2)-negative BC (OR = 1.25, 95% CI 1.05–1.48, $p = 0.012$). No evidence of a gene-environment interaction was observed between rs671 and alcohol intake ($p = 0.537$).

Conclusion: This study suggests that the Lys/Lys genotype confers susceptibility to BC risk among women of Asian ancestry, particularly for ER-positive, PR-positive, and HER2-negative tumor types.

KEYWORDS

acetaldehyde, alcohol drinking, aldehyde dehydrogenase-2, breast cancer, single nucleotide polymorphism

1 | INTRODUCTION

Epidemiological studies consistently indicate that alcohol is an independent risk factor for female breast cancer (BC) (Singletary & Gapstur, 2001). The International Agency for Research on Cancer concluded that there is sufficient evidence to classify alcohol as a carcinogen for female BC (IARC Working Group on the Evaluation of Carcinogenic Risks to Humans, 2010). One hypothesized mechanism behind alcohol-related breast carcinogenesis is the involvement of acetaldehyde, a metabolite of ethanol. An impact of acetaldehyde on carcinogenesis for several types of alcohol-induced cancers has been shown in experimental models (Brooks & Theruvathu, 2005). Molecular epidemiological studies demonstrated a gene-environment interaction between a functional aldehyde dehydrogenase 2 (*ALDH2*) polymorphism (rs671: Glu>Lys, OMIM: 100650) and alcohol intake for esophageal and upper digestive tract cancers in East Asian countries (Matsuo et al., 2001; Oze et al., 2010), where rs671 is prevalent (Li et al., 2009). These studies support

the hypothesis that acetaldehyde is a carcinogen. The Glu/Lys heterozygotes of rs671 have far less than half of *ALDH2* activity of Glu/Glu homozygotes, and the Lys/Lys homozygotes have no detectable *ALDH2* activity, which leads to high acetaldehyde concentrations upon alcohol intake in individuals harboring the Lys allele (Crabb, Edenberg, Bosron, & Li, 1989). Therefore, exploring the association of rs671 with BC risk and its interaction with alcohol intake is one approach to elucidate whether acetaldehyde is a causative agent for breast carcinogenesis. To date, evidence of an association of rs671 with BC risk is scarce; statistically significant associations have not been observed in case-control studies in Japan (456 cases and 912 controls) (Kawase et al., 2009) Korea (346 cases and 377 controls) (Choi et al., 2003) or Thailand (561 cases and 486 controls) (Sangrajrang et al., 2010). We conducted a pooled analysis of individual genetic and alcohol consumption data for women of Asian ancestry participating in studies in the Breast Cancer Association Consortium (BCAC) with at least 18 times larger sample size than previous studies.

2 | METHODS

2.1 | Study population

We used data from 14 case-control studies in the BCAC. Table 1 shows participating studies contributing to this pooled analysis. All study participants were of Asian ancestry and recruited from studies conducted in Asian countries, Canada, and the USA. Eight studies were hospital-based, five were population-based, and one included hospital-based cases and population-based controls. We included 12,595 BC cases and 12,884 controls for the analysis of rs671 and BC risk. For the analysis of the gene-environment interaction between rs671 and alcohol intake for BC risk, we included 2,849 BC cases and 3,680 controls after excluding participants with missing values for alcohol intake from seven studies. All studies were approved by their local ethics review boards, and all participants provided informed consent. This investigation was approved by a human research investigations committee at Aichi Cancer Center.

2.2 | Genotyping methods

Genotyping was carried out using the iCOGS array (<http://ccge.medschl.cam.ac.uk/research/consortia/icogs/>), or the OncoArray (<https://support.illumina.com/downloads/infinium-oncoarray-500k-v1-0-product-files.html>). Details of array design, genotyping, postgenotyping quality control, and imputation have been provided elsewhere (Michailidou et al., 2013, 2017). The rs671 SNP on *ALDH2* was a candidate SNP selected on the basis of specific hypotheses described above.

To adjust for potential population stratification, principal components analyses (PCA) were carried out separately for Asian subgroups. Briefly, PCA was performed based on a subset of 37,000 uncorrelated SNPs for the iCOGS data and based on 33,661 uncorrelated SNPs for the OncoArray data. For the present analyses, we used two Asian principal components for the iCOGS dataset and 10 Asian principal components for the OncoArray dataset as covariates. Further details have been provided in previous articles (Michailidou et al., 2013, 2017).

2.3 | Alcohol assessment

Each study ascertained alcohol intake via self-reported questionnaire. Daily alcohol intake in grams was determined by summing the product of frequency of consumption of specified alcoholic beverages (beer, wine, and other alcoholic beverages) by the alcohol content of each beverage using national estimates of alcohol content for that country. The exposure period was the year preceding recruitment. A multistep harmonization procedure was used to reconcile differences in individual study questionnaires.

2.4 | Statistical analysis

To assess the associations of rs671 with BC risk, we estimated odds ratios (ORs) with 95% confidence intervals (CIs) by unconditional logistic regression models using the Glu/Glu genotype as reference. This was done separately for iCOGS and OncoArray datasets, and results were combined by a fixed-effects meta-analysis. The ORs were adjusted for age, Asian principal components, and study. We also evaluated the associations by tumor characteristics (estrogen receptor, ER; progesterone receptor, PR; human epidermal growth factor receptor 2, HER2) and tumor subtypes (luminal [either ER or PR positive, HER2 negative], triple positive [ER, PR, HER2 positive], HER2 enrich [ER, PR negative, HER2 positive], triple negative [ER, PR, HER2 negative]) using cases with these specific characteristics. Heterogeneity by tumor characteristics and between studies was assessed using Cochran's Q test. We assessed the gene-environment interaction between rs671 and alcohol intake by including an interaction term. Alcohol intake was classified in three ways: 1) two categories (none, any alcohol intake); 2) three categories (none, <15 g ethanol/day, ≥15 g ethanol/day); and 3) four categories (none, <15 g ethanol/day, 15–30 g ethanol/day, ≥30 g ethanol/day). We also performed stratified analyses by menopausal status: women with missing menopausal status were considered premenopausal if they were ≤50 years or postmenopausal if >50 years. All statistical analyses were performed using Stata version 15.1 (Stata Corp., College Station, TX, USA), with a P value <0.05 considered to be statistically significant.

3 | RESULTS

Demographic characteristics of participants are shown in Table 2. The median age was 50 years for both cases and controls, with a higher proportion of women in the oldest age groups for cases. The proportion of nondrinkers and heavy drinkers (≥15 g ethanol/day) was higher among controls than cases, possibly due to the smaller number of unknown category in controls (71.4%) than in cases (77.4%). The distributions of tumor characteristics among cases were 7,648 ER positive (60.7%), 6,308 PR positive (50.1%), and 3,054 HER2 positive (24.3%) for participants included in the analysis of rs671 alone and, 1,871 ER positive (65.7%), 1,620 PR positive (56.9%), and 552 HER2 positive (19.4%) for those in the analysis of gene-environment interaction, respectively.

Table 3 presents the associations of rs671 with BC risk. Overall, the Lys/Lys genotype was associated with increased BC risk, with OR of 1.16 (95% CI = 1.03–1.30, $p = 0.014$) relative to Glu/Glu genotype. According to tumor characteristics, we observed an association of the Lys/Lys genotype with ER-positive BC (OR = 1.19, 95% CI 1.05–1.36, $p = 0.008$),

TABLE 1 List of participating studies and number of participants

Study acronym	Study name	Study design	Country	Subjects of analysis for rs671				Subjects of analysis for GE interaction			
				Case	Control	Lys allele frequency among cases (%)	Lys allele frequency among controls (%)	Case	Control	Lys allele frequency among cases (%)	Lys allele frequency among controls (%)
ACP	Asia Cancer Program	Hospital based case-control study	Thailand	830	1,060	8.9	8.0	—	—	—	—
CBCS	Canadian Breast Cancer Study	Population-based case-control study	Canada	252	170	28.6	20.0	—	—	—	—
HERPACC	Hospital-based Epidemiologic Research Program at Aichi Cancer	Hospital-based case-control study	Japan	792	1,659	29.9	28.3	783	1,632	30.1	28.6
HKBCS	Hong Kong Breast Cancer Study	Hospital-based case-control study	China	466	451	32.1	28.4	—	—	—	—
KOHBRA	Korean Hereditary Breast Cancer Study	Population-based case-control study	Korea	1,251	665	17.1	15.9	413	601	6.8	15.4
LAABC	Los Angeles County Asian-American Breast Cancer Case-Control Study	Population-based case-control study	USA	808	990	24.9	27.5	808	990	24.9	27.5
MYBRCA	Malaysian Breast Cancer Genetic Study	Hospital-based case-control study	Malaysia	1,408	1,866	24.5	22.6	—	—	—	—
NC-BCFR	Northern California Breast Cancer Family Registry	Population-based case-control study	USA	446	52	21.4	21.2	400	46	22.4	23.9

(Continues)

TABLE 1 (Continued)

Study acronym	Study name	Study design	Country	Subjects of analysis for rs671				Subjects of analysis for GE interaction			
				Case	Control	Lys allele frequency among cases (%)	Lys allele frequency among controls (%)	Case	Control	Lys allele frequency among cases (%)	Lys allele frequency among controls (%)
NGOBCS	Nagano Breast Cancer Study	Hospital-based case-control study	Japan	366	366	25.4	23.6	366	365	25.4	23.7
SBCGS	Shanghai Breast Cancer Genetic Study	Population-based case-control study, cohort study	China	1,644	1,827	24.9	23.9	5	46	10.0	7.6
SEBCS	Seoul Breast Cancer Study	Hospital-based case-control study	Korea	2,129	2,236	16.9	15.1	74	—	2.7	—
SGBCC	Singapore Breast Cancer Cohort	Hospital based breast cancer cohort and population-based controls	Singapore	775	798	20.8	23.6	—	—	—	—
TBCS	IARC-Thai Breast Cancer Study	Hospital-based case-control study	Thailand	138	253	6.9	11.7	—	—	—	—
TWBCS	Taiwanese Breast Cancer Study	Hospital-based case-control study	Taiwan	1,290	491	27.8	31.5	—	—	—	—
Total				12,595	12,884	22.1	21.4	2,849	3,680	22.8	23.3

Abbreviation: GE interaction, gene-environment interaction.

TABLE 2 Characteristics of cases and controls

	Subjects of analysis for rs671		Subjects of analysis for GE interaction					
	Cases (N = 12,595)	(%)	Control (N = 12,884)	(%)	Cases (N = 2,849)	(%)	Control (N = 3,680)	(%)
Age (years)								
Median (range)	50 (20–91)		50 (15–92)		50 (20–81)		50 (19–86)	
≤29	205	1.6	300	2.3	60	2.1	68	1.9
30–39	1,641	13.0	1,254	9.7	421	14.8	408	11.1
40–49	4,255	33.8	4,547	35.3	845	29.7	1,234	33.5
50–59	3,847	30.5	4,179	32.4	830	29.1	1,117	30.4
60–69	1,911	15.2	2,138	16.6	498	17.5	638	17.3
≥70	736	5.8	466	3.6	195	6.8	215	5.8
Alcohol consumption [†]								
g/day (mean ± SD)	31.2 ± 91.2		30.5 ± 83.0		31.2 ± 91.2		30.5 ± 83.0	
Nondrinker	1,746	13.9	2,348	18.2	1,746	61.3	2,348	63.8
<15 g ethanol/day	895	7.1	1,052	8.2	895	31.4	1,052	28.6
≥15 g ethanol/day	208	1.7	280	2.2	208	7.3	280	7.6
Unknown	9,746	77.4	9,204	71.4				
ALDH2 Glu/Glu genotype								
g/day (mean ± SD)	42.4 ± 103.9		44.8 ± 102.0		42.4 ± 103.9		44.8 ± 102.0	
Nondrinker	828	10.6	1,040	12.9	828	48.0	1,040	50.7
<15 g ethanol/day	719	9.2	774	9.6	719	41.7	774	37.8
≥15 g ethanol/day	178	2.3	236	2.9	178	10.0	236	11.5
Unknown	6,056	77.8	5,988	74.5				
ALDH2 Glu/Lys genotype								
g/day (mean ± SD)	16.1 ± 68.5		13.3 ± 43.0		16.1 ± 68.5		13.3 ± 43.0	
Nondrinker	745	18.3	1,076	25.8	745	78.6	1,076	77.1
<15 g ethanol/day	173	4.3	276	6.6	173	18.3	276	19.8
≥15 g ethanol/day	30	0.7	44	1.1	30	3.2	44	3.2
Unknown	3,122	76.7	2,779	66.6				
ALDH2 Lys/Lys genotype								

(Continues)

TABLE 2 (Continued)

	Subjects of analysis for rs671		Subjects of analysis for GE interaction					
	Cases (N = 12,595)	(%)	Control (N = 12,884)	(%)	Cases (N = 2,849)	(%)	Control (N = 3,680)	(%)
g/day (mean ± SD)	0.4 ± 3.4		0.5 ± 5.1		0.4 ± 3.4		0.5 ± 5.1	
Nondrinker	232	99.2	173	98.3	232	99.2	173	98.3
<15 g ethanol/day	2	0.8	3	1.7	2	0.8	3	1.7
≥15 g ethanol/day	0	0	0	0	0	0	0	0
Unknown	0	0	0	0	0	0	0	0
Menopausal status								
Premenopausal	3,690	29.3	5,234	40.6	836	29.3	1,393	37.9
Postmenopausal	4,287	34.0	4,830	37.5	879	30.9	1,246	33.9
Unknown	4,618	36.7	2,820	21.9	1,134	39.8	1,041	28.3
ER status								
Positive	7,648	60.7			1,871	65.7		
Negative	3,701	29.4			658	23.1		
Unknown	1,246	9.9			320	11.2		
PR status								
Positive	6,308	50.1			1,620	56.9		
Negative	3,776	30.0			865	30.4		
Unknown	2,511	19.9			364	12.8		
HER2 status								
Positive	3,054	24.3			552	19.4		
Negative	4,054	32.2			557	19.6		
Unknown	5,487	43.6			1,740	61.1		

Abbreviations: ALDH2, aldehyde dehydrogenase 2; ER, estrogen receptor; GE interaction, gene-environment interaction; HER2, human epidermal growth factor receptor 2; PR, progesterone receptor.
[†]Exposure period was the year preceding recruitment.

PR-positive BC (OR = 1.19, 95% CI 1.03–1.36, $p = 0.015$), and HER2-negative BC (OR = 1.25, 95% CI 1.05–1.48, $p = 0.012$), but not with ER-negative BC (OR = 1.07, 95% CI 0.90–1.27, $p = 0.453$), PR-negative BC (OR = 1.13, 95% CI 0.95–1.34, $p = 0.176$), or HER2-positive BC (OR = 1.19, 95% CI 0.97–1.48, $p = 0.102$), although no statistically significant heterogeneity was observed by tumor characteristics. According to tumor subtypes, the Lys/Lys genotype was only associated with luminal BC (OR = 1.30, 95% CI 1.09–1.55, $p = 0.004$), and not with other subtypes (Table 4). No evidence of heterogeneity was also observed by menopausal status (Table S1).

Figure S1 and Figure S2 show the forest plots of study-specific ORs for the association between rs671 and BC risk. With regard to the association between the Glu/Lys genotype and BC risk, there was no evidence of between-study heterogeneity (p for heterogeneity = 0.380). In contrast, significant between-study heterogeneity was observed for the association of the Lys/Lys genotype with BC risk (p for heterogeneity = 0.003), which was mainly attributable to a strong positive association for CBCS and a strong inverse association for ACP and TWBCS. However, exclusion of these studies did not alter the significant association of the Lys/Lys genotype with BC risk (OR = 1.18, 95% CI 1.05–1.33, $p = 0.008$).

TABLE 3 Association between ALDH2 genotype and breast cancer risk

	ALDH2 genotype			p for heterogeneity between tumor characteristics	
	Glu/Glu	Glu/Lys	Lys/Lys	For Glu/Lys	For Lys/Lys
Overall					
Cases/controls	7,781/8,038	4,070/4,175	744/671		
OR (95% CI) [†]	1 (ref.)	1.03 (0.97–1.08, $p = 0.350$)	1.16 (1.03–1.30, $p = 0.014$)		
ER status					
Positive					
Cases/controls	4,636/8,038	2,531/4,175	481/671		
OR (95% CI) [†]	1 (ref.)	1.01 (0.95–1.08, $p = 0.669$)	1.19 (1.05–1.36, $p = 0.008$)	0.447	0.329
Negative					
Cases/control	2,321/8,038	1,187/4,175	193/671		
OR (95% CI) [†]	1 (ref.)	1.05 (0.97–1.14, $p = 0.257$)	1.07 (0.90–1.27, $p = 0.453$)		
PR status					
Positive					
Cases/controls	3,842/8,038	2,066/4,175	400/671		
OR (95% CI) [†]	1 (ref.)	0.98 (0.92–1.05, $p = 0.591$)	1.19 (1.03–1.36, $p = 0.015$)	0.410	0.653
Negative					
Cases/control	2,333/8,038	1,238/4,175	205/671		
OR (95% CI) [†]	1 (ref.)	1.02 (0.95–1.11, $p = 0.545$)	1.13 (0.95–1.34, $p = 0.176$)		
HER2 status					
Positive					
Cases/control	1,961/8,038	940/4,175	153/671		
OR (95% CI) [†]	1 (ref.)	1.02 (0.92–1.14, $p = 0.674$)	1.19 (0.97–1.48, $p = 0.102$)	1.000	0.720
Negative					
Cases/control	2,521/7,841	1,287/4,175	246/671		
OR (95% CI) [†]	1 (ref.)	1.02 (0.93–1.11, $p = 0.722$)	1.25 (1.05–1.48, $p = 0.012$)		

Abbreviations: ALDH2, aldehyde dehydrogenase 2; CI, confidence intervals; ER, estrogen receptor; HER2, human epidermal growth factor receptor 2; OR, odds ratios; PR, progesterone receptor.

[†]ORs were adjusted for age (continuous), Asian principal components and study site.

TABLE 4 Association between ALDH2 genotype and breast cancer risk by tumor subtypes

	ALDH2 genotype			<i>p</i> for heterogeneity between tumor characteristics	
	Glu/Glu	Glu/Lys	Lys/Lys	For Glu/Lys	For Lys/Lys
Luminal					
Cases/controls	1,950/8,038	979/4,175	198/671		
OR (95% CI) [†]	1 (ref.)	1.00 (0.92–1.10, <i>p</i> = 0.916)	1.30 (1.09–1.55, <i>p</i> = 0.004)	0.452	0.755
Triple positive					
Cases/controls	1,202/8,038	583/4,175	93/671		
OR (95% CI) [†]	1 (ref.)	1.03 (0.91–1.16, <i>p</i> = 0.640)	1.19 (0.93–1.53, <i>p</i> = 0.164)		
HER2 enrich					
Cases/controls	694/8,038	322/4,175	55/671		
OR (95% CI) [†]	1 (ref.)	0.96 (0.83–1.11, <i>p</i> = 0.557)	1.12 (0.83–1.51, <i>p</i> = 0.453)		
Triple negative					
Cases/control	546/8,038	310/4,175	46/671		
OR (95% CI) [†]	1 (ref.)	1.13 (0.97–1.32, <i>p</i> = 0.108)	1.11 (0.81–1.53, <i>p</i> = 0.519)		

Abbreviations: ALDH2, aldehyde dehydrogenase 2; CI, confidence intervals; HER2, human epidermal growth factor receptor 2; OR, odds ratios.

[†]ORs were adjusted for age (continuous), Asian principal components and study site.

and there was no longer evidence of between-study heterogeneity (*p* for heterogeneity = 0.133). Furthermore, when we repeated analyses using random effects meta-analyses to calculate summary study-specific estimates, the results did not change substantially (Table S2).

Stratified analyses by alcohol intake categories assessing a gene-environment interaction between rs671 and alcohol intake showed no evidence of interaction, although the sample size is small compared to the analysis of rs671 and BC risk (Table S3, *p* for interaction = 0.537).

4 | DISCUSSION

In this study, we found that the Lys/Lys genotype of rs671 was associated with increased BC risk among women of Asian ancestry. No evidence of interaction was observed between rs671 and alcohol intake. This is the largest study to date to perform this evaluation quantitatively using high-quality individual-level data for Asian women.

Several epidemiological studies have reported a gene-environment interaction between rs671 and alcohol intake for several types of cancer (Hiraki et al., 2007; Ishioka et al., 2018; Masaoka et al., 2016; Matsuo et al., 2001, 2013; Oze et al., 2010). Our findings are not consistent with our hypothesis of gene-environment interaction between rs671 and alcohol intake. Considering the established impact of rs671 on cancer risk, this lack of interaction suggests that acetaldehyde may

be less influential in breast carcinogenesis. Other biological mechanisms for alcohol-related breast carcinogenesis have been hypothesized, including increased circulating estrogens and androgens, enhancement of mammary gland susceptibility to carcinogenesis, increased mammary carcinogen DNA damage, interference of folate metabolism by alcohol, and greater potential for invasiveness into BC cells (Bernstein & Ross, 1993; Singletary & Gapstur, 2001; Singletary & McNary, 1994; Stolzenberg-Solomon et al., 2006). To better understand the etiologic nature of the effect of alcohol on breast carcinogenesis, further investigations are needed.

We observed an association of the Lys/Lys genotype with increased BC risk. Because individuals with the Lys/Lys genotype have no detectable ALDH2 activity and almost completely refrain from drinking due to severe adverse reactions caused by acetaldehyde (e.g., facial flushing, nausea and headache) (Matsuo et al., 2006), the observed genetic association suggests that the Lys/Lys genotype confers susceptibility to BC risk independently of alcohol intake. ALDH2 plays a key role in removal of not only ethanol-derived acetaldehyde, but also other toxic endogenous aldehydes such as 4-hydroxy-2-nonenal (4-HNE) and malondialdehyde (Chen, Ferreira, Gross, & Mochly-Rosen, 2014). These endogenous aldehydes have been reported to cause DNA damage and might be related to breast carcinogenesis (Chen et al., 2014; Garaycochea et al., 2018). In addition, we did not find an association of the Glu/Lys genotype with BC risk. This suggest that

ALDH2 activity of the Glu/Lys homozygotes may be sufficient for detoxifying toxic endogenous aldehydes related to breast carcinogenesis. In contrast, the Lys/Lys homozygotes have no detectable ALDH2 activity, thus may not tolerate these endogenous aldehydes. Furthermore, the Lys/Lys genotype was associated with increased risk only in hormone receptor positive BC, and not in hormone receptor negative BC. These results suggest that the biological mechanism could be through a hormonal receptor mediated pathway (Zhang, Man, Zhao, Dong, & Ma, 2014). The evidence of an association of rs671 with BC risk is scarce and may warrant additional evaluation in future studies.

The strengths of this investigation include the analysis of individual-level data from a large sample of Asian women, allowing us to obtain stable, and precise summary estimates of the association of rs671 with BC risk. Other strengths are the uniform genotyping procedures and quality-control measures undertaken for the iCOGS and the OncoArray, respectively. We were also able to control for population stratification by including Asian principal components as a covariate to control for residual genetic heterogeneity. Furthermore, the Lys allele of rs671 is only prevalent in East Asia, and has not been found in Caucasians or Africans (Li et al., 2009). Thus, this analysis is unique and can be performed only among Asian women. Several limitations also warrant consideration. First, we could not evaluate the association between alcohol intake and BC risk because there were a lot of missing data on potential confounding factors (e.g., smoking, estrogen-related factors) and we were not able to control for them. However, genotypes are fixed at birth and these factors cannot influence genotypes; therefore, our results about rs671 and BC risk may be unbiased even though we did not adjust for these factors. Second, even though all study participants were of Asian ancestry, the heterogeneity across study populations, designs, and methods are potential limitations. Third, careful interpretation of results from the analysis of gene-environment interaction and stratified analyses is necessary because we had a limited number of participants in some sub-groups and did not adjust for multiple comparisons.

In conclusion, we observed an association between the Lys/Lys genotype of rs671 and increased BC risk. Among women of Asian ancestry, this study suggests that the Lys/Lys genotype confers susceptibility to BC risk, particularly for ER-positive, PR-positive, and HER2-negative tumor types. These findings warrant further investigation in future studies.

ACKNOWLEDGMENTS

We thank all the individuals who took part in these studies and all the researchers, clinicians, technicians, and administrative staff who have enabled this work to be carried out. The COGS study would not have been possible without the contributions of the following: Per Hall (COGS); Douglas F. Easton, Paul

Pharoah, Kyriaki Michailidou, Manjeet K. Bolla, Qin Wang (BCAC), Joe Dennis, Alison M. Dunning, Andrew Lee, and Ed Dicks, Craig Luccarini and the staff of the Centre for Genetic Epidemiology Laboratory, Javier Benitez, Anna Gonzalez-Neira and the staff of the CNIO genotyping unit, Jacques Simard and Daniel C. Tessier, Francois Bacot, Daniel Vincent, Sylvie LaBoissière and Frederic Robidoux and the staff of the McGill University and Génome Québec Innovation Centre, Stig E. Bojesen, Sune F. Nielsen, Borge G. Nordestgaard, and the staff of the Copenhagen DNA laboratory, and Julie M. Cunningham, Sharon A. Windebank, Christopher A. Hilker, Jeffrey Meyer and the staff of Mayo Clinic Genotyping Core Facility. The ACP study wishes to thank the participants in the Thai Breast Cancer study. Special Thanks also go to the Thai Ministry of Public Health (MOPH), doctors and nurses who helped with the data collection process. Finally, the study would like to thank Dr. Prat Boonyawongviroj, the former Permanent Secretary of MOPH and Dr. Pornthep Siriwanarungsan, the former Department Director-General of Disease Control who has supported the study throughout. CBCS thanks study participants, co-investigators, collaborators, and staff of the Canadian Breast Cancer Study, and project coordinators Agnes Lai and Celine Morissette. HKBCS thanks Hong Kong Sanatorium and Hospital, Dr. Ellen Li Charitable Foundation, The Kerry Group Kuok Foundation, National Institute of Health 1R03CA130065, and the North California Cancer Center for support. We thank all investigators of the KOHBRA (Korean Hereditary Breast Cancer) Study. LAABC thanks all the study participants and the entire data collection team, especially Annie Fung and June Yashiki. MYBRCA thanks study participants and research staff (particularly Patsy Ng, Nurhidayu Hassan, Yoon Sook-Yee, Daphne Lee, Lee Sheau Yee, Phuah Sze Yee and Norhashimah Hassan) for their contributions and commitment to this study. SGBCC thanks the participants and research coordinator Ms Tan Siew Li.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ORCID

Hidemi Ito  <https://orcid.org/0000-0002-8023-4581>

Keitaro Matsuo  <https://orcid.org/0000-0003-1761-6314>

REFERENCES

- Bernstein, L., & Ross, R. K. (1993). Endogenous hormones and breast cancer risk. *Epidemiologic Reviews*, 15(1), 48–65. <https://doi.org/10.1093/oxfordjournals.epirev.a036116>
- Brooks, P. J., & Theruvathu, J. A. (2005). DNA adducts from acetaldehyde: Implications for alcohol-related carcinogenesis. *Alcohol*, 35(3), 187–193. <https://doi.org/10.1016/j.alcohol.2005.03.009>

- Chen, C. H., Ferreira, J. C., Gross, E. R., & Mochly-Rosen, D. (2014). Targeting aldehyde dehydrogenase 2: New therapeutic opportunities. *Physiological Reviews*, *94*(1), 1–34. <https://doi.org/10.1152/physrev.00017.2013>
- Choi, J.-Y., Abel, J., Neuhaus, T., Ko, Y., Harth, V., Hamajima, N., ... Kang, D. (2003). Role of alcohol and genetic polymorphisms of CYP2E1 and ALDH2 in breast cancer development. *Pharmacogenetics*, *13*(2), 67–72. <https://doi.org/10.1097/00008571-200302000-00002>
- Crabb, D. W., Edenberg, H. J., Bosron, W. F., & Li, T. K. (1989). Genotypes for aldehyde dehydrogenase deficiency and alcohol sensitivity. The inactive ALDH2(2) allele is dominant. *Journal of Clinical Investigation*, *83*(1), 314–316. <https://doi.org/10.1172/JCI113875>
- Garaycochea, J. I., Crossan, G. P., Langevin, F., Mulderigg, L., Louzada, S., Yang, F., ... Patel, K. J. (2018). Alcohol and endogenous aldehydes damage chromosomes and mutate stem cells. *Nature*, *553*(7687), 171–177.
- Hiraki, A., Matsuo, K., Wakai, K., Suzuki, T., Hasegawa, Y., & Tajima, K. (2007). Gene-gene and gene-environment interactions between alcohol drinking habit and polymorphisms in alcohol-metabolizing enzyme genes and the risk of head and neck cancer in Japan. *Cancer Science*, *98*(7), 1087–1091. <https://doi.org/10.1111/j.1349-7006.2007.00505.x>
- IARC Working Group on the Evaluation of Carcinogenic Risks to Humans. (2010). Alcohol consumption and ethyl carbamate. *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans*, *96*, 3–1383.
- Ishioka, K., Masaoka, H., Ito, H., Oze, I., Ito, S., Tajika, M., ... Matsuo, K. (2018). Association between ALDH2 and ADH1B polymorphisms, alcohol drinking and gastric cancer: A replication and mediation analysis. *Gastric Cancer*, *21*(6), 936–945. <https://doi.org/10.1007/s10120-018-0823-0>
- Kawase, T., Matsuo, K., Hiraki, A., Suzuki, T., Watanabe, M., Iwata, H., ... Tajima, K. (2009). Interaction of the effects of alcohol drinking and polymorphisms in alcohol-metabolizing enzymes on the risk of female breast cancer in Japan. *Journal of Epidemiology*, *19*(5), 244–250. <https://doi.org/10.2188/jea.JE20081035>
- Li, H., Borinskaya, S., Yoshimura, K., Kal'ina, N., Marusin, A., Stepanov, V. A., ... Kidd, K. K. (2009). Refined geographic distribution of the oriental ALDH2*504Lys (nee 487Lys) variant. *Annals of Human Genetics*, *73*(Pt 3), 335–345. <https://doi.org/10.1111/j.1469-1809.2009.00517.x>
- Masaoka, H., Ito, H., Soga, N., Hosono, S., Oze, I., Watanabe, M., ... Matsuo, K. (2016). Aldehyde dehydrogenase 2 (ALDH2) and alcohol dehydrogenase 1B (ADH1B) polymorphisms exacerbate bladder cancer risk associated with alcohol drinking: Gene-environment interaction. *Carcinogenesis*, *37*(6), 583–588. <https://doi.org/10.1093/carcin/bgw033>
- Matsuo, K., Hamajima, N., Shinoda, M., Hatooka, S., Inoue, M., Takezaki, T., & Tajima, K. (2001). Gene-environment interaction between an aldehyde dehydrogenase-2 (ALDH2) polymorphism and alcohol consumption for the risk of esophageal cancer. *Carcinogenesis*, *22*(6), 913–916. <https://doi.org/10.1093/carcin/22.6.913>
- Matsuo, K., Oze, I., Hosono, S., Ito, H., Watanabe, M., Ishioka, K., ... Tanaka, H. (2013). The aldehyde dehydrogenase 2 (ALDH2) Glu504Lys polymorphism interacts with alcohol drinking in the risk of stomach cancer. *Carcinogenesis*, *34*(7), 1510–1515. <https://doi.org/10.1093/carcin/bgt080>
- Matsuo, K., Wakai, K., Hirose, K., Ito, H., Saito, T., & Tajima, K. (2006). Alcohol dehydrogenase 2 His47Arg polymorphism influences drinking habit independently of aldehyde dehydrogenase 2 Glu487Lys polymorphism: Analysis of 2,299 Japanese subjects. *Cancer Epidemiology, Biomarkers & Prevention*, *15*(5), 1009–1013. <https://doi.org/10.1158/1055-9965.EPI-05-0911>
- Michailidou, K., Hall, P., Gonzalez-Neira, A., Ghoussaini, M., Dennis, J., Milne, R. L., ... Easton, D. F. (2013). Large-scale genotyping identifies 41 new loci associated with breast cancer risk. *Nature Genetics*, *45*(4), 353–361. <https://doi.org/10.1038/ng.2563>
- Michailidou, K., Lindstrom, S., Dennis, J., Beesley, J., Hui, S., Kar, S., ... Easton, D. F. (2017). Association analysis identifies 65 new breast cancer risk loci. *Nature*, *551*(7678), 92–94. <https://doi.org/10.1038/nature24284>
- Oze, I., Matsuo, K., Hosono, S., Ito, H., Kawase, T., Watanabe, M., ... Tanaka, H. (2010). Comparison between self-reported facial flushing after alcohol consumption and ALDH2 Glu504Lys polymorphism for risk of upper aerodigestive tract cancer in a Japanese population. *Cancer Science*, *101*(8), 1875–1880. <https://doi.org/10.1111/j.1349-7006.2010.01599.x>
- Sangrajang, S., Sato, Y., Sakamoto, H., Ohnami, S., Khuhaprema, T., & Yoshida, T. (2010). Genetic polymorphisms in folate and alcohol metabolism and breast cancer risk: A case-control study in Thai women. *Breast Cancer Research and Treatment*, *123*(3), 885–893. <https://doi.org/10.1007/s10549-010-0804-4>
- Singletary, K. W., & Gapstur, S. M. (2001). Alcohol and breast cancer: Review of epidemiologic and experimental evidence and potential mechanisms. *JAMA*, *286*(17), 2143–2151. <https://doi.org/10.1001/jama.286.17.2143>
- Singletary, K. W., & McNary, M. Q. (1994). Influence of ethanol intake on mammary gland morphology and cell proliferation in normal and carcinogen-treated rats. *Alcoholism, Clinical and Experimental Research*, *18*(5), 1261–1266. <https://doi.org/10.1111/j.1530-0277.1994.tb00116.x>
- Stolzenberg-Solomon, R. Z., Chang, S.-C., Leitzmann, M. F., Johnson, K. A., Johnson, C., Buys, S. S., ... Ziegler, R. G. (2006). Folate intake, alcohol use, and postmenopausal breast cancer risk in the Prostate, Lung, Colorectal, and Ovarian Cancer Screening Trial. *American Journal of Clinical Nutrition*, *83*(4), 895–904. <https://doi.org/10.1093/ajcn/83.4.895>
- Zhang, M. H., Man, H. T., Zhao, X. D., Dong, N., & Ma, S. L. (2014). Estrogen receptor-positive breast cancer molecular signatures and therapeutic potentials (Review). *Biomed Rep*, *2*(1), 41–52. <https://doi.org/10.3892/br.2013.187>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Ugai T, Milne RL, Ito H, et al. The functional *ALDH2* polymorphism is associated with breast cancer risk: A pooled analysis from the Breast Cancer Association Consortium. *Mol Genet Genomic Med*. 2019;7:e707. <https://doi.org/10.1002/mgg3.707>