

Body mass index and subsequent fracture risk: a meta-analysis to update FRAX

Nicholas C. Harvey^{1,2,*} , Helena Johansson³, Eugene V. McCloskey^{3,4} , Enwu Liu⁵ , Kristina E. Åkesson^{6,7} , Fred A. Anderson⁸, Rafael Azagra-Ledesma^{9,10,11,12}, Cecilie L. Bager¹³, Charlotte Beaudart¹⁴, Heike A. Bischoff-Ferrari^{15,16}, Emmanuel Biver¹⁷ , Olivier Bruyère¹⁸, Jane A. Cauley¹⁹, Jacqueline R. Center^{20,21} , Roland Chapurlat²², Claus Christiansen¹³, Cyrus Cooper^{1,2,23} , Carolyn J. Crandall²⁴ , Steven R. Cummings²⁵, José A.P. da Silva^{26,27}, Bess Dawson-Hughes²⁸ , Adolfo Diez-Perez²⁹, Alyssa B. Dufour^{30,31} , John A. Eisman^{21,32,33} , Petra J.M. Elders³⁴, Serge Ferrari¹⁷ , Yuki Fujita³⁵, Saeko Fujiwara³⁶, Claus-Christian Glüer³⁷, Inbal Goldshtein^{38,39}, David Goltzman⁴⁰, Vilmundur Gudnason^{41,42}, Jill Hall⁴³, Didier Hans⁴⁴ , Mari Hoff^{45,46}, Rosemary J. Hollick⁴⁷, Martijn Huisman^{48,49}, Masayuki Iki⁵⁰ , Sophia Ish-Shalom⁵¹, Graeme Jones⁵², Magnus K. Karlsson^{6,53} , Sundeep Khosla⁵⁴ , Douglas P. Kiel^{31,55} , Woon-Puay Koh^{56,57}, Fjorda Koromani^{58,59} , Mark A. Kotowicz^{60,61,62}, Heikki Kröger^{63,64}, Timothy Kwok^{65,66} , Olivier Lamy^{67,68} , Arnulf Langhammer^{69,70}, Bagher Larijani⁷¹ , Kurt Lippuner⁷², Fiona E.A. McGuigan⁶ , Dan Mellström^{73,74}, Thomas Merlijn³⁴ , Tuan V. Nguyen^{75,76,77}, Anna Nordström^{78,79,80}, Peter Nordström⁸¹, Terence W. O'Neill^{82,83}, Barbara Obermayer-Pietsch⁸⁴, Claes Ohlsson^{85,86} , Eric S. Orwoll⁸⁷ , Julie A. Pasco^{88,89,90,91}, Fernando Rivadeneira⁵⁸, Berit Schei^{92,93}, Anne-Marie Schott⁹⁴, Eric J. Shiroma⁹⁵, Kristin Siggeirsdottir^{41,96}, Eleanor M. Simonsick⁹⁷ , Elisabeth Sornay-Rendu⁹⁸, Reijo Sund⁶⁴ , Karin M.A. Swart^{99,100}, Pawel Szulc⁹⁸ , Junko Tamaki¹⁰¹ , David J. Torgerson¹⁰², Natasja M. van Schoor^{103,104}, Tjeerd P. van Staa¹⁰⁵, Joan Vila¹⁰⁶, Nicholas J. Wareham¹⁰⁷, Nicole C. Wright¹⁰⁸ , Noriko Yoshimura¹⁰⁹, M. Carola Zillikens⁵⁸ , Marta Zwart^{12,110,111,112}, Liesbeth Vandenput⁸⁵, Mattias Lorentzon^{113,114} , William D. Leslie¹¹⁵ , John A. Kanis³

¹MRC Lifecourse Epidemiology Centre, University of Southampton, Southampton, SO16 6YD, United Kingdom

²NIHR Southampton Biomedical Research Centre, University of Southampton and University Hospital Southampton NHS Foundation Trust, Southampton, SO16 6YD, United Kingdom

³Centre for Metabolic Bone Diseases, University of Sheffield, Sheffield, United Kingdom

⁴MRC Versus Arthritis Centre for Integrated Research in Musculoskeletal Ageing, Mellanby Centre for Musculoskeletal Research, University of Sheffield, Sheffield, United Kingdom

⁵South Australian Health and Medical Research Institute (SAHMRI), Adelaide, SA, Australia

⁶Clinical and Molecular Osteoporosis Research Unit, Department of Clinical Sciences, Lund University, Lund, Sweden

⁷Department of Orthopedics, Skåne University Hospital, Malmö, Sweden

⁸GLOW Coordinating Center, Center for Outcomes Research, University of Massachusetts Medical School, Worcester, MA, United States

⁹Department of Medicine, Autonomous University of Barcelona, Barcelona, Spain

¹⁰Docencia Atenció Primària, Metropolitana Nord, Institut Català de la Salut, Barcelona, Spain

¹¹GROIMAP/GROICAP (research groups), Unitat de Suport a la Recerca de Girona, Institut Universitari d'Investigació en Atenció Primària Jordi Gol, Girona, Spain

¹²PRECIOSA-Fundación para la investigación, Barberà del Vallés, Barcelona, Spain

¹³Nordic Bioscience A/S, Herlev, Denmark

¹⁴Public Health Aging Research and Epidemiology (PHARE) Group, Research Unit in Clinical Pharmacology and Toxicology (URPC), NAMur Research Institute for Life Sciences (NARILIS), Faculty of Medicine, University of Namur, 5000 Namur, Belgium

¹⁵Department of Aging Medicine, Felix Platter, University of Basel, 4055 Basel, Switzerland

¹⁶Centre on Aging and Mobility, University of Zurich and City Hospital, Zurich, Switzerland

¹⁷Division of Bone Diseases, Department of Medicine, Geneva University Hospitals and Faculty of Medicine, University of Geneva, Geneva, Switzerland

¹⁸Research Unit in Public Health, Epidemiology and Health Economics, University of Liège, Liège, Belgium

Received: May 1, 2025. Revised: June 11, 2025. Accepted: June 20, 2025

© The Author(s) 2025. Published by Oxford University Press on behalf of the American Society for Bone and Mineral Research.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

- ¹⁹Department of Epidemiology, School of Public Health, University of Pittsburgh, Pittsburgh, PA, United States
- ²⁰Garvan Institute of Medical Research, Sydney, NSW, Australia
- ²¹St Vincent's Clinical School, School of Medicine and Health, University of New South Wales Sydney, Sydney, NSW, Australia
- ²²INSERM UMR 1033, Université Claude Bernard-Lyon1, Hôpital Edouard Herriot, Lyon, France
- ²³NIHR Oxford Biomedical Research Centre, University of Oxford, Oxford, United Kingdom
- ²⁴Division of General Internal Medicine and Health Services Research, David Geffen School of Medicine, University of California, Los Angeles, CA, United States
- ²⁵San Francisco Coordinating Center, California Pacific Medical Center Research Institute, San Francisco, CA, United States
- ²⁶Centre for Innovative Biomedicine and Biotechnology, Faculty of Medicine, University of Coimbra, Coimbra, Portugal
- ²⁷Rheumatology Department, University Hospital and University of Coimbra, Coimbra, Portugal
- ²⁸Bone Metabolism Laboratory, Jean Mayer US Department of Agriculture Human Nutrition Research Center on Aging, Tufts University, Boston, MA, United States
- ²⁹Department of Internal Medicine, Hospital del Mar and CIBERFES, Autonomous University of Barcelona, Barcelona, Spain
- ³⁰Marcus Institute for Aging Research, Hebrew SeniorLife, Boston, MA, United States
- ³¹Department of Medicine, Beth Israel Deaconess Medical Center and Harvard Medical School, Boston, MA, United States
- ³²Skeletal Diseases Program, Garvan Institute of Medical Research, Sydney, NSW, Australia
- ³³School of Medicine Sydney, University of Notre Dame Australia, Sydney, NSW, Australia
- ³⁴Department of General Practice, Amsterdam UMC, location AMC, Amsterdam Public Health Research Institute, Amsterdam, The Netherlands
- ³⁵Department of Hygiene and Public Health, Faculty of Medicine, Kansai Medical University, Osaka, Japan
- ³⁶Department of Pharmacy, Yasuda Women's University, Hiroshima, Japan
- ³⁷Section Biomedical Imaging, Molecular Imaging North Competence Center, Department of Radiology and Neuroradiology, University Medical Center Schleswig-Holstein Kiel, Kiel University, Kiel, Germany
- ³⁸Maccabitech Institute of Research and Innovation, Maccabi Healthcare Services, Tel Aviv, Israel
- ³⁹Department of Epidemiology and Preventive Medicine, Sackler Faculty of Medicine, School of Public Health, Tel Aviv University, Tel Aviv, Israel
- ⁴⁰Department of Medicine, McGill University and McGill University Health Centre, Montreal, QC, Canada
- ⁴¹Icelandic Heart Association, Kopavogur, Iceland
- ⁴²University of Iceland, Reykjavik, Iceland
- ⁴³MRC Centre for Reproductive Health, University of Edinburgh, Edinburgh, United Kingdom
- ⁴⁴Interdisciplinary Centre of Bone Diseases, Bone and Joint Department, Lausanne University Hospital (CHUV) and University of Lausanne, Lausanne, Switzerland
- ⁴⁵Department of Neuromedicine and Movement Science, Norwegian University of Science and Technology, Trondheim, Norway
- ⁴⁶Department of Rheumatology, St. Olavs Hospital, Trondheim, Norway
- ⁴⁷Aberdeen Centre for Arthritis and Musculoskeletal Health, Epidemiology Group, University of Aberdeen, Aberdeen, United Kingdom
- ⁴⁸Department of Epidemiology and Data Science, Amsterdam Public Health Research Institute, VU University Medical Center, Amsterdam, The Netherlands
- ⁴⁹Department of Sociology, VU University, Amsterdam, The Netherlands
- ⁵⁰Department of Public Health, Kindai University Faculty of Medicine, Osaka, Japan
- ⁵¹Endocrine Clinic, Elisha Hospital, Haifa, Israel
- ⁵²Menzies Institute for Medical Research, University of Tasmania, Hobart, TAS, Australia
- ⁵³Department of Orthopaedics, Skåne University Hospital, Malmö, Sweden
- ⁵⁴Robert and Arlene Kogod Center on Aging and Division of Endocrinology, Mayo Clinic College of Medicine, Mayo Clinic, Rochester, MN, United States
- ⁵⁵Marcus Institute for Aging Research, Hebrew Senior Life, Boston, MA, United States
- ⁵⁶Healthy Longevity Translational Research Programme, Yong Loo Lin School of Medicine, National University of Singapore, Singapore
- ⁵⁷Singapore Institute for Clinical Sciences, Agency for Science Technology and Research (A*STAR), Singapore
- ⁵⁸Department of Internal Medicine, Erasmus University Medical Center, Rotterdam, The Netherlands
- ⁵⁹Department of Radiology and Nuclear Medicine, Erasmus University Medical Center, Rotterdam, The Netherlands
- ⁶⁰Institute for Mental and Physical Health and Clinical Translation (IMPACT), Deakin University, Geelong, VIC, Australia
- ⁶¹Barwon Health, Geelong, VIC, Australia
- ⁶²Department of Medicine-Western Health, The University of Melbourne, St Albans, VIC, Australia
- ⁶³Department of Orthopedics and Traumatology, Kuopio University Hospital, Kuopio, Finland
- ⁶⁴Kuopio Musculoskeletal Research Unit, University of Eastern Finland, Kuopio, Finland
- ⁶⁵Department of Medicine and Therapeutics, Faculty of Medicine, The Chinese University of Hong Kong, Hong Kong
- ⁶⁶Jockey Club Centre for Osteoporosis Care and Control, Faculty of Medicine, The Chinese University of Hong Kong, Hong Kong
- ⁶⁷Centre of Bone Diseases, Lausanne University Hospital, Lausanne, Switzerland
- ⁶⁸Service of Internal Medicine, Lausanne University Hospital, Lausanne, Switzerland
- ⁶⁹HUNT Research Centre, Department of Public Health and Nursing, Faculty of Medicine and Health Sciences, Norwegian University of Science and Technology, Trondheim, Norway
- ⁷⁰Levanger Hospital, Nord-Trøndelag Hospital Trust, Levanger, Norway
- ⁷¹Endocrinology and Metabolism Research Center, Endocrinology and Metabolism Clinical Sciences Institute, Tehran University of Medical Sciences, Tehran, Iran
- ⁷²ARTORG Center for Biomedical Engineering Research, Faculty of Medicine, University of Bern, Bern, Switzerland
- ⁷³Geriatric Medicine, Department of Internal Medicine and Clinical Nutrition, Institute of Medicine, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden
- ⁷⁴Geriatric Medicine, Sahlgrenska University Hospital Mölndal, Mölndal, Sweden
- ⁷⁵School of Biomedical Engineering, University of Technology Sydney, Sydney, NSW, Australia
- ⁷⁶School of Population Health, UNSW Medicine, UNSW Sydney, Kensington, NSW, Australia

- ⁷⁷Tam Anh Research Institute, Tam Anh Hospital, Ho Chi Minh City, Vietnam
- ⁷⁸School of Sport Sciences, UiT The Arctic University of Norway, Tromsø, Norway
- ⁷⁹Department of Health Sciences, Swedish Winter Sports Research Centre, Mid Sweden University, Östersund, Sweden
- ⁸⁰Department of Medical Sciences, Uppsala University, Uppsala, Sweden
- ⁸¹Department of Public Health and Caring Sciences, Uppsala University, Uppsala, Sweden
- ⁸²National Institute for Health Research Manchester Biomedical Research Centre, Manchester University NHS Foundation Trust, Manchester Academic Health Science Centre, Manchester, United Kingdom
- ⁸³Centre for Epidemiology Versus Arthritis, University of Manchester, Manchester, United Kingdom
- ⁸⁴Division of Endocrinology and Diabetology, Department of Internal Medicine, Medical University Graz, Graz, Austria
- ⁸⁵Sahlgrenska Osteoporosis Centre, Department of Internal Medicine and Clinical Nutrition, Institute of Medicine, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden
- ⁸⁶Department of Drug Treatment, Sahlgrenska University Hospital, Gothenburg, Sweden
- ⁸⁷Department of Medicine, Oregon Health and Science University, Portland, OR, United States
- ⁸⁸Institute for Physical and Mental Health and Clinical Translation (IMPACT), Deakin University, Geelong, VIC, Australia
- ⁸⁹Department of Medicine-Western Health, The University of Melbourne, St Albans, VIC, Australia
- ⁹⁰Barwon Health, Geelong, VIC, Australia
- ⁹¹Department of Epidemiology and Preventive Medicine, Monash University, Melbourne, VIC, Australia
- ⁹²Department of Public Health and Nursing, Norwegian University of Science and Technology, Trondheim, Norway
- ⁹³Department of Obstetrics and Gynaecology, St Olavs Hospital, Trondheim, Norway
- ⁹⁴Université Claude Bernard Lyon 1, RESHAPE U INSERM U-1290, Lyon, France
- ⁹⁵Laboratory of Epidemiology and Population Sciences, National Institute on Aging, Baltimore, MD, United States
- ⁹⁶Janus Rehabilitation, Reykjavik, Iceland
- ⁹⁷Translational Gerontology Branch, National Institute on Aging Intramural Research Program, Baltimore, MD, United States
- ⁹⁸INSERM UMR 1033, University of Lyon, Hôpital Edouard Herriot, Lyon, France
- ⁹⁹Department of General Practice, Amsterdam UMC, location VUmc, Amsterdam Public Health Research Institute, Amsterdam, The Netherlands
- ¹⁰⁰PHARMO Institute for Drug Outcomes Research, Utrecht, The Netherlands
- ¹⁰¹Department of Hygiene and Public Health, Faculty of Medicine, Educational Foundation of Osaka Medical and Pharmaceutical University, Osaka, Japan
- ¹⁰²York Trials Unit, Department of Health Sciences, University of York, York, United Kingdom
- ¹⁰³Department of Epidemiology and Data Science, Amsterdam UMC location Vrije Universiteit, Amsterdam, The Netherlands
- ¹⁰⁴Amsterdam Public Health Research Institute, Aging and Later Life Program, Amsterdam, The Netherlands
- ¹⁰⁵Centre for Health Informatics, Faculty of Biology, Medicine and Health, School of Health Sciences, University of Manchester, Manchester, United Kingdom
- ¹⁰⁶Statistics Support Unit, Hospital del Mar Medical Research Institute, CIBER Epidemiology and Public Health (CIBERESP), Barcelona, Spain
- ¹⁰⁷MRC Epidemiology Unit, University of Cambridge, Cambridge, United Kingdom
- ¹⁰⁸Center for Health Outcomes, Implementation, and Community-Engaged Sciences, Tulane University, New Orleans, LA, United States
- ¹⁰⁹Department of Preventive Medicine for Locomotive Organ Disorders, The University of Tokyo Hospital, Tokyo, Japan
- ¹¹⁰Health Center Can Gibert del Pla, Catalan Institute of Health, Girona, Spain
- ¹¹¹Department of Medical Sciences, University of Girona, Girona, Spain
- ¹¹²GROICAP (Research Group), Institut Universitari d'Investigació en Atenció Primària Jordi Gol, Girona, Spain
- ¹¹³Sahlgrenska Osteoporosis Centre, Institute of Medicine, University of Gothenburg, Gothenburg, Sweden
- ¹¹⁴Region Västra Götaland, Geriatric Medicine, Sahlgrenska University Hospital, Mölndal, Sweden
- ¹¹⁵Department of Medicine, University of Manitoba, Winnipeg, MB, Canada
- *Corresponding author: Nicholas C. Harvey, MRC Lifecourse Epidemiology Centre, University of Southampton, Southampton General Hospital, Southampton SO16 6YD, United Kingdom (nch@mrc.soton.ac.uk).

Abstract

The aim of this international meta-analysis was to quantify the predictive value of BMI for incident fracture and relationship of this risk with age, sex, follow-up time, and BMD. A total of 1 667 922 men and women from 32 countries (63 cohorts), followed for a total of 16.0 million person-years were studied. 293 325 had FN BMD measured (2.2 million person-years follow-up). An extended Poisson model in each cohort was used to investigate relationships between WHO-defined BMI categories (Underweight: <18.5 kg/m²; Normal: 18.5-24.9 kg/m²; Overweight: 25.0-29.9 kg/m²; Obese I: 30.0-34.9 kg/m²; Obese II: ≥35.0 kg/m²) and risk of incident osteoporotic, major osteoporotic and hip fracture (HF). Inverse-variance weighted β -coefficients were used to merge the cohort-specific results. For the subset with BMD available, in models adjusted for age and follow-up time, the hazard ratio (95% CI) for HF comparing underweight with normal weight was 2.35 (2.10-2.60) in women and for men was 2.45 (1.90-3.17). Hip fracture risk was lower in overweight and obese categories compared to normal weight [obese II vs normal: women 0.66 (0.55-0.80); men 0.91 (0.66-1.26)]. Further adjustment for FN BMD T-score attenuated the increased risk associated with underweight [underweight vs normal: women 1.69 (1.47-1.96); men 1.46 (1.00-2.13)]. In these models, the protective effects of overweight and obesity were attenuated, and in both sexes, the direction of association reversed to higher fracture risk in Obese II category [Obese II vs Normal: women 1.24 (0.97-1.58); men 1.70 (1.06-2.75)]. Results were similar for other fracture outcomes. Underweight is a risk factor for fracture in both men and women regardless of adjustment for BMD. However, while overweight/obesity appeared protective in base models, they became risk factors after additional adjustment for FN BMD, particularly in the Obese II category. This effect in the highest BMI categories was of greater magnitude in men than women. These results will inform the second iteration of FRAX®.

Keywords: BMI, meta-analysis, hip fracture, osteoporosis, epidemiology, major osteoporotic fracture, FRAX

Introduction

Body mass index (BMI), calculated as weight divided by height squared, is an accepted surrogate for adiposity in population studies and as a clinical risk factor in assessment for outcomes, such as myocardial infarction and fracture.^{1,2} Multiple studies have documented the complex relationships between fat mass, lean mass, and bone mineral density (BMD) in the determination of fracture risk.^{3,4} Obesity is associated with an increased risk of falling.⁵ Both low and high BMI have been associated with greater risk of fracture, but importantly, with different fracture sites at either end of the BMI spectrum: Findings from the Global Longitudinal study of Osteoporosis in Women (GLOW) suggested high risk of ankle and upper leg fractures with obesity,⁶ with similar results in the Spanish SIDAP dataset.⁷ More recently, a Mendelian Randomization study demonstrated causal associations between low BMI and risk of forearm fracture.⁸ Consistent with these observations, in our previous meta-analysis of 398 610 women, mean age 63 years, low BMI was a risk factor for hip and all osteoporotic fractures, but appeared to be protective for lower leg/ankle fracture.² In contrast, high BMI was associated with increased risk for humerus and elbow fractures. This relationship was dependent partly on BMD. Thus, after adjustment for FN BMD, high BMI remained a risk factor for upper arm fracture but also predicted increased risk of all osteoporotic fractures. BMI therefore conveys risk information for incident fracture, but with important considerations regarding the relationships with BMD and fracture site. However, the evidence base thus far principally focuses on women, with relationships in men, and indeed across ethnicities in either sex, still requiring detailed elucidation.

FRAX, currently available in 86 territories, is the most widely used fracture risk assessment tool and is incorporated into a large number of assessment guidelines,⁹ recommended by the Committee for Medicinal Products for Human Use (CHMP),¹⁰ and approved by the National Institute for Health and Care Excellence (NICE).¹¹ The incorporation of BMI as an input variable for risk prediction was based on an earlier meta-analysis.¹² Since then, many more prospectively studied cohorts have become available that have the potential to improve the accuracy of FRAX and understand better the potential influences of BMD, sex, and ethnicity in these relationships.¹³

The aim of the present study was to quantify magnitude of the risk relationship between BMI and incident fracture in an international setting, and to explore the dependence of this risk on age, sex, ethnicity, time since baseline assessment, and BMD.

Materials and methods

Details of our initial systematic review undertaken to identify appropriate cohorts, and of the cohorts included, have been documented previously¹³ and are summarized in Table S1. This was registered in the PROSPERO international prospective register of systematic reviews (CRD42021227266), and followed the Preferred Reporting Items for Systematic Reviews (PRISMA) guidelines. In brief, we studied 1 667 922 men and women from 63 prospectively studied cohorts. A total of 57 cohorts included women ($n = 1\,127\,206$) and 40 cohorts included men ($n = 540\,716$). Of these 293 325 men and women had femoral neck (FN) BMD measured, from 53

cohorts in 20 countries, with a total follow-up time of 2.2 million person-years.

Baseline and outcome variables

BMI (kg/m^2) was calculated from height and weight assessed at baseline. We considered incident fractures in the following categories: all, osteoporotic, major osteoporotic (MOF: clinical spine, distal forearm, hip, or humerus), and hip fracture (HF). For the purpose of this analysis, and related meta-analyses informing the next iteration of FRAX,^{14–16} we defined “osteoporotic fractures” on the basis of previously identified sites at which fractures were more common with increasing age and decreasing BMD, thus excluding fractures of the skull, face, hands, feet, ankle, and patella, as well as tibia and fibula fractures in men.^{17,18} This should not be taken to mean that anyone experiencing such a fracture has osteoporosis, merely that they appear more frequent in those with low BMD and/or higher age. No distinction was made according to trauma since both high- and low-trauma fractures show similar relationships with low BMD and future fracture risk.^{19,20} Details of fracture ascertainment and validation have been published previously.¹³

Statistical methods

The risk of incident fracture, expressed as hazard ratio (HR) per 1 kg/m^2 greater baseline BMI, was estimated using an extended Poisson model applied separately to each cohort (and also separately by sex for those cohorts with both men and women).²¹ Because of an embargo on data transfer, Cox regression was used on the Manitoba cohort. Covariates included current time since start of follow-up, current age, prior history of fracture, and BMD T-score at the FN. BMD was adjusted for manufacturer and T-scores were calculated from the NHANES III White female reference values.¹³ We categorized BMI according to WHO recommendations (Underweight: <18.5 ; Normal: $18.5\text{--}24.9$ (referent); Overweight: $25.0\text{--}29.9$; Obese I: $30.0\text{--}34.9$; Obese II: ≥ 35.0 kg/m^2)²² for analyses by category. We illustrated non-linear associations using spline models with nodes at 21, 25, and 33 kg/m^2 . Results of each cohort and the 2 sexes were weighted according to the variance and merged to determine the weighted means and standard deviations. The HR was equal to $e^{\text{weighted mean of } \beta}$. A priori, a random effects model was used in the meta-analysis. Assessment of the effects of ethnicity was confined to those cohorts recording more than one ethnic group [Asian, Black, Hispanic, White and the Black, Asian and Minority Ethnic category (BAME)], comprising Health ABC, CAMOS, MrOS USA, LASA, WHI, SOF, Manitoba, and UK Biobank. Ascertainment of these data was via self-report. On the basis of these 8 cohorts there was sufficient data to undertake pairwise comparisons. Finally, we investigated associations between BMI and risk of death during follow-up.

Results

Of 1 667 922 men and women studied in 32 countries, the average BMI was 27.2 kg/m^2 . At follow up, 151 473 men and women were identified as having a subsequent fracture of any kind; 125 135 were characterized as osteoporotic in men and women, 95 748 men and women sustained a MOF; 31 383

Table 1. Hazard ratio (HR) and 95% CI for fracture at the sites indicated per unit greater BMI (linear models). HRs are adjusted for age and time since baseline.

Outcome fracture	Adjusted for age and time since baseline			Adjusted for age and time since baseline—for those with BMD			Adjusted for age, time since baseline and BMD		
	Number of cohorts	I ²	HR (95% CI)	Number of cohorts	I ²	HR (95% CI)	Number of cohorts	I ²	HR (95% CI)
Female									
Any	55	73	0.99 (0.98, 0.99)	46	65	0.98 (0.98, 0.99)	46	42	1.01 (1.01, 1.01)
Hip	57	57	0.95 (0.94, 0.95)	46	41	0.95 (0.94, 0.96)	46	33	0.99 (0.98, 1.00) ^{0.060}
MOF	52	83	0.98 (0.97, 0.98)	45	77	0.98 (0.97, 0.98)	45	57	1.01 (1.00, 1.02) ^{0.0018}
Ost	51	83	0.98 (0.97, 0.98)	44	73	0.98 (0.97, 0.98)	44	55	1.01 (1.00, 1.01) ^{0.0051}
Male									
Any	38	25	0.99 (0.98, 0.99)	32	0	0.99 (0.98, 0.99)	32	0	1.02 (1.01, 1.02)
Hip	36	48	0.95 (0.93, 0.96)	29	32	0.95 (0.93, 0.97)	28	0	1.03 (1.01, 1.04)
MOF	35	60	0.98 (0.97, 0.98)	30	0	0.98 (0.97, 0.98)	30	1	1.02 (1.01, 1.03)
Ost	34	56	0.98 (0.97, 0.99)	30	18	0.98 (0.97, 0.99)	30	0	1.02 (1.01, 1.03)

Except where indicated, $p < .001$. Abbreviations: MOF, major osteoporotic fracture; Ost, osteoporotic fracture.

were HFs. The total follow-up time was 16.0 million-person years in men and women. BMD measurements were available in 17.6% (293325) of individuals. Table S1 summarizes key characteristics of the cohorts.

Association between BMI and risk of incident fracture

In continuous models, summarized in Table 1, after adjustment for age and time since baseline, greater BMI was associated with lower risk of incident fracture in both men and women, for example, HR (95% CI) for major osteoporotic fracture per unit greater BMI in women was 0.98 (0.97, 0.98); in men 0.98 (0.97, 0.98). Figure 1 documents these associations for HF and major osteoporotic fracture in men and women combined as Forest plots. Associations appeared similar by sex and were not materially different in the subset of cohorts in which BMD was measured. In this subset, after further adjustment for BMD, there was a reversal in the relationships such that greater BMI was weakly associated with higher fracture risk. For example, among women HR for major osteoporotic fracture per unit greater BMI was 1.01 (1.00, 1.02) and for men 1.02 (1.01, 1.03) (Table 1). Overall, associations were similar between men and women (Table S2).

Categorical and non-linear associations between BMI and risk of incident fracture

On average, those who were underweight tended to be somewhat older than those in other categories and had lower FN BMD T-score (Table S3). Indeed, there was modest but positive gradient of mean FN BMD T-score going from underweight to normal to overweight to obese categories. Tables 2 and 3 and Figure 2 document the associations between BMI category relative to the normal BMI for men and women, in the subset with BMD measurements, unadjusted and adjusted for FN BMD. As with the initial models, associations were not materially different in the subset with BMD, without BMD adjustment, but the increased risk associated with low BMI and the more modest protective effect of higher BMI category was attenuated by adjustment for FN BMD. Indeed, after BMD adjustment, obesity became a risk factor for fracture in both sexes, with evidence of a greater effect size in men than women (Tables 2 and 3, Table S4). Figure S1 presents

the continuous relationships between BMI (HR for fracture for a given BMI vs a BMI of 25 kg/m²) and risk of HF in men and women separately, in the subset with BMD, without and then with adjustment for FN BMD.

Ethnicity

Table S5 documents the fracture outcome models with continuous BMI as the exposure, by ethnicity (HR and 95%CI fracture risk for each one-unit greater BMI for White, Asian, Black, and Hispanic ethnicities, together with p value for a BMI•ethnicity interaction). There was no statistical evidence for a difference in the association between BMI and HF by ethnicity. In contrast to White ethnicity, for Asian and Hispanic ethnicities, higher BMI was associated with a greater risk of any fracture [Asian: HR (95% CI) 1.02 (1.00, 1.04); Hispanic 1.02 (1.00, 1.05)].

Interaction with age and follow-up time

For all, osteoporotic and MOF fracture outcomes, the HR for fracture per one-unit greater BMI tended to decrease with increasing age. Conversely, for the outcome of HF, there was no obvious relationship with age. (Table S6). There was a modest increase in HR with greater follow-up time for the HF and MOF in the whole cohort after adjusting for age (Table S7).

Association between BMI and mortality

Compared to normal weight, both underweight [HR (95%CI): 1.60 (1.44, 1.78) in women and 1.71 (1.50, 1.94) in men] and obese I and II categories [e.g., obese II: 1.31 (1.21, 1.43) in women and 1.27 (1.07, 1.51) in men] were associated with greater risk of death during follow-up, in both sexes. Overweight was not associated with increased mortality risk and indeed was protective in women [0.92 (0.88, 0.95)] and men [0.96 (0.92, 1.00)]. Overall associations were similar after adjustment for FN BMD T-score. Table S8A and B summarizes these associations.

Discussion

Summary of findings

The present study represents the largest meta-analysis to date on the association between BMI and subsequent fracture risk.




Figure 1. Forest plot showing effect size on hip fracture (HF) risk (left panel, A) and major osteoporotic fracture (right panel, B) per one unit greater BMI in men and women combined adjusted for age and time since baseline.

Table 2. Hazard ratio (HR) for fracture by BMI category (reference normal) in women and men, adjusted for age and time since baseline, in subset with FN BMD measure available (but not adjusted for BMD). (BMI categories: underweight BMI < 18.5, normal BMI 18.5-24.9, overweight BMI 25.0-29.9, obese I BMI 30.0-34.9, obese II BMI ≥ 35.0 kg/m²). HR in bold are <0.001.

Outcome fracture	N, cohorts	1 Underweight vs Normal HR (95% CI)	3 Overweight vs Normal HR (95% CI)	4 Obese I vs Normal HR (95% CI)	5 Obese II vs Normal HR (95% CI)
Women					
Any	38, 47, 45, 40	1.31 [1.20, 1.43]	0.94 [0.90, 0.98]	0.88 [0.83, 0.93]	0.84 [0.77, 0.91]
Hip	27, 41, 33, 22	2.35 [2.13, 2.60]	0.75 [0.72, 0.78]	0.64 [0.59, 0.69]	0.66 [0.55, 0.80]
MOF	33, 45, 42, 36	1.43 [1.30, 1.59]	0.90 [0.85, 0.95]	0.82 [0.76, 0.89]	0.78 [0.70, 0.88]
Ost	34, 44, 42, 36	1.38 [1.27, 1.50]	0.90 [0.86, 0.94]	0.84 [0.79, 0.90]	0.77 [0.70, 0.85]
Men					
Any	21, 30, 27, 23	1.77 [1.54, 2.04]	0.87 [0.84-0.90]	0.88 [0.82, 0.94]	1.04 [1.04, 1.05]
Hip	14, 23, 16, 11	2.45 [1.90, 3.17]	0.69 [0.65-0.74]	0.63 [0.53, 0.74]	0.91 [0.66, 1.26]
MOF	16, 28, 23, 17	1.87 [1.60, 2.20]	0.85 [0.80-0.90]	0.80 [0.72, 0.89]	0.96 [0.96, 0.97]
Ost	19, 28, 25, 21	1.83 [1.58, 2.12]	0.85 [0.82-0.89]	0.82 [0.76, 0.89]	0.96 [0.96, 0.97]
p women vs men					
Any		<0.001	0.0067	>0.30	<0.001
Hip		>0.30	0.024	>0.30	0.088
MOF		0.0040	0.18	>0.30	<0.001
Ost		0.0011	0.053	>0.30	<0.001

Abbreviations: MOF, major osteoporotic fracture; Ost, osteoporotic fracture.

It confirms our previous observation in women that low BMI is associated with increased fracture risk, persisting after adjustment for FN BMD.² Furthermore, it confirms in women

that overall higher BMI becomes a risk factor for fracture after adjustment for BMD, despite being protective when used unadjusted. In addition to these confirmatory findings, our

Table 3. Hazard ratio (HR) for fracture by BMI category (reference normal) in women and men, adjusted for age and time since baseline, in subset with FN BMD measure available, additionally adjusted for BMD. (BMI categories: underweight BMI < 18.5, normal BMI 18.5-24.9, overweight BMI 25.0-29.9, obese I BMI 30.0-34.9, obese II BMI \geq 35.0 kg/m²). HR in bold are <0.001.

Outcome fracture	N, cohorts	1 Underweight vs Normal HR (95% CI)	3 Overweight vs Normal HR (95% CI)	4 Obese I vs Normal HR (95% CI)	5 Obese II vs Normal HR (95% CI)
Women					
Any	38, 47, 45, 40	1.09 [1.03, 1.17]	1.08 [1.04, 1.12]	1.13 [1.07, 1.20]	1.19 [1.10, 1.29]
Hip	27, 41, 33, 22	1.69 [1.47, 1.96]	0.99 [0.93, 1.05]	0.97 [0.89, 1.06]	1.24 [0.97, 1.58]
MOF	33, 45, 42, 36	1.16 [1.08, 1.25]	1.07 [1.01, 1.13]	1.12 [1.04, 1.21]	1.21 [1.09, 1.35]
Ost	34, 44, 42, 36	1.13 [1.06, 1.21]	1.06 [1.01, 1.11]	1.13 [1.05, 1.21]	1.17 [1.05, 1.29]
Men					
Any	21, 30, 27, 23	1.23 [1.00, 1.51]	1.03 [0.97, 1.09]	1.14 [1.05, 1.24]	1.43 [1.24, 1.65]
Hip	14, 23, 16, 11	1.46 [1.00, 2.13]	1.06 [0.95, 1.18]	1.10 [0.87, 1.38]	1.70 [1.06, 2.75]
MOF	16, 28, 23, 17	1.25 [0.98, 1.60]	1.06 [0.98, 1.15]	1.14 [1.01, 1.29]	1.54 [1.26, 1.88]
Ost	19, 28, 25, 21	1.23 [1.00, 1.52]	1.03 [0.96, 1.09]	1.11 [1.02, 1.22]	1.53 [1.27, 1.83]
p women vs men					
Any		0.27	0.19	>0.30	0.027
Hip		>0.30	0.29	>0.30	0.25
MOF		>0.30	>0.30	>0.30	0.037
Ost		>0.30	>0.30	>0.30	0.045

Abbreviations: MOF, major osteoporotic fracture; Ost, osteoporotic fracture.

new analysis documents important novel associations in men: although we observed a similar pattern overall to that in women, the increased risk of fracture associated with obesity after adjustment for BMD was of greater magnitude in men compared with women.

Comparison with previous studies

This analysis provides an updated comparison with our previous meta-analysis, which examined relationships between BMI and fracture risk only in women, based on a much smaller cohort database with substantially less time at risk available.² There, we observed that low BMI was a risk factor for hip and all osteoporotic fractures; when adjusted for BMD, it remained a risk factor for HF but was weakly protective for osteoporotic fracture. In contrast, high BMI, when adjusted for BMD, was a risk factor for osteoporotic fractures. Our initial 2005 meta-analysis focused on both men and women but with a smaller number of participants and, while the results were broadly comparable, was not able to elucidate any differences by sex.¹² Similar adverse effect of high BMI on fracture risk, after adjustment for BMD, were observed in the US MrOS cohort.²³ Compston et al. examined the relationship between BMI and fracture outcomes in the GLOW observing, in 52 939 postmenopausal women, a protective association between greater BMI and risk of hip, clinical spine, and wrist fractures.²⁴ In contrast, higher BMI was associated weakly with greater risk of ankle fracture. While this study has sometimes been interpreted by others as demonstrating that obesity is not protective for osteoporotic fracture, there is clearly a question of whether ankle fractures are a consequence of osteoporosis per se, given that the age and sex relationship differs from that with classical osteoporotic fractures.^{17,18,25-27} Thus rather than being protective for ankle fractures, obesity appears to be a risk factor for these events. The findings are overall consistent with our current meta-analysis in suggesting that associations between BMI and fracture risk are site specific, with potential further mechanistic implications.

While factors, such as chronic inflammation, type 2 diabetes, and propensity to falls, may contribute to fracture risk

in obesity,^{5,28} other considerations may underpin apparent sex differences observed. For example, in the present analysis, the greater risk of HF in men compared with women at high BMI after consideration of FN BMD may reflect sex dependent distribution of adipose tissue, with the female distribution more classically subcutaneous around the hips than abdomen, compared with the visceral abdominal deposition in men.^{29,30} Biomechanically, the additional adipose tissue overlying the proximal femur would favor a protective effect for HF in women compared with men. Additionally, or alternatively, a further mechanism might relate to sex-dependent differential hormonal patterns in obesity. Excess adipose tissue is associated with increased estrogen activity, with likely positive effects on bone and muscle.³¹ However, in men, obesity may lead to a concomitant reduction in male sex steroids and thus potentially effective hypogonadism,^{32,33} with increased fracture risk possibly additionally mediated via impaired muscle health in addition to reduced BMD.³⁴

Strengths and limitations

We undertook the largest meta-analysis to date of mostly population-based cohorts to investigate and quantify associations between BMI and risk of incident fractures. We were able to investigate BMI as a continuous exponential, nonlinear and categorical exposure, and investigate possible interactions by age, sex, ethnicity, and follow-up time. The use of primary data decreases the risk of publication bias and the general consistency of the BMI-fracture association between cohorts provides strong support for the international validity of this risk variable. However, there are limitations which need to be considered in the interpretation of our findings. First, as with nearly all population-based studies, non-response biases may have occurred, which we were unable to document for all cohorts. This is likely to result in a cohort which is healthier than the background population and may thus lead to underestimation of the absolute risk of fracture. In particular, there may have been reduced responses from those at the lowest and highest BMI values. It is therefore likely, if anything, to have had a conservative effect on our risk estimates. Second, it is possible that there were differences in




Figure 2. Hazard ratio (HR) (95% CI) for fracture according to categories of BMI (referent: normal BMI) for men and women. (A) Hip fracture (HF) and (B) major osteoporotic fracture (MOF). Associations are adjusted for age and follow-up time +/- FN BMD T-score, in subset with FN BMD measure available.

protocols and, for example, execution of height and weight measures between cohorts. This is unavoidable in such a study, and importantly, exposure-outcome analyses were undertaken within each cohort and then the beta coefficients merged

through random effects meta-analysis. Third, we were not able to elucidate whether the association between BMI and incident fracture was causal. However, this was not the purpose of the study and for the purpose of risk assessment, a causal

relationship is not required. Fourth, BMI is affected by both fat and lean mass and does not account for distribution of fat between subcutaneous and visceral compartments. This is irrelevant in terms risk assessment, but of course limits mechanistic influence. Finally, we were not able to account for levels of physical activity, which might be related to both BMI and fracture risk. Although this might be mechanistically interesting, it would not reflect the way in which BMI is considered in the FRAX algorithm.

Clinical and risk assessment implications

The role of BMI in the current FRAX model was informed by an earlier meta-analysis by De Laet et al.,¹² which was broadly consistent with the Johansson et al., 2014 study,² the conclusion at that time being that the new findings did not necessitate modification of the FRAX engine itself. Here we present novel associations relating to men as well as women, with interactions by sex, follow-up time, ethnicity, and age, which will directly inform the revised FRAX engine, as a continuous exposure, in the second version of this globally established tool. Importantly our current findings also confirm the non-linear association between BMI and fracture. The variably modifying effects of age, ethnicity, and follow-up time, together with differences by sex, support the accommodation of such nuances in the FRAX risk engine, but conversely mitigate against using BMI thresholds in risk assessment, since these would not necessarily convey consistent risk information across these parameters or fracture outcomes. The associations between increased risk of death during follow-up and underweight, and to a lesser extent obesity, are consistent with a large recent analyses,³⁵ and will also be important to consider in the derivation of the revised FRAX tool.

While there are likely to be causal implications, underpinned by our understanding of biology and biomechanics, for the purposes of risk assessment, the principal requirement is quantification of the exposure-outcome association and of interactions between this relationship and with other potential input variables. Notwithstanding, our findings directly inform clinical practice, indicating that greater BMI is not necessarily protective for all fracture types. While this may play into strategies to tackle obesity, there is evidence that weight loss usually leads to concomitant loss of muscle and bone and so approaches addressing obesity in the context of fracture risk must be carefully thought out.³⁶ Adequate calcium and protein nutrition, together with a mix of weight-bearing and resistance exercise training, alongside dietary interventions to reduce calorie intake are likely to be important,³⁷ given the clear demonstration of increasing risk of HF with lower BMI, even after adjustment for BMD.

Conclusion

In the largest meta-analysis to date of prospective cohorts worldwide, we have confirmed inverse associations between fracture risk and BMI, with the apparent protective effect of higher BMI reversed after adjustment for FN BMD. The magnitude of this effect appeared greater in men than women. The increased accuracy of the elucidated effect sizes, and their specificity by age, sex, follow-up time, and ethnicity will inform improved risk assessment in the second iteration of the FRAX tool.

Human and animal rights

This review does not contain any original studies with human participants or animals performed by any of the authors.

Ethics

All individual cohorts with candidate risk factors available have been approved by their local ethics committees and informed consent has been obtained from all study participants. General ethics approval for the use of these cohorts is also given by the University of Sheffield. Participant data are stored in coded, de-identified form. Only summary statistics and aggregate data are published, not allowing for identification of individual study participants.

Acknowledgments

We are grateful to Dr. Östen Ljunggren for contributing the MrOS Sweden cohort. UK Biobank data are included under approved access agreement 3593. The authors acknowledge the Manitoba Centre for Health Policy for use of Manitoba data contained in the Population Health Research Data Repository (HIPC 2016/2017-29). The results and conclusions are those of the authors and no official endorsement by the Manitoba Centre for Health Policy, Manitoba Health, Seniors and Active Living, or other data providers is intended or should be inferred. N.C.H. acknowledges funding from the UK Medical Research Council [MC_PC_21003; MC_PC_21001] and the NIHR Southampton Biomedical Research Centre, University of Southampton and University Hospital Southampton NHS Foundation Trust, UK. Terence O'Neill is supported by the National Institute for Health and Care Research (NIHR) Manchester Biomedical Research Centre (BRC) (NIHR203308). The WHI program is funded by the National Heart, Lung, and Blood Institute, National Institutes of Health, U.S. Department of Health and Human Services through 75N92021D00001, 75N92021D00002, 75N92021D00003, 75N92021D00004, 75N92021D00005. The Osteoporotic Fractures in Men (MrOS) Study is supported by National Institutes of Health funding. The following institutes provide support: the National Institute on Aging (NIA), the National Institute of Arthritis and Musculoskeletal and Skin Diseases (NIAMS), the National Center for Advancing Translational Sciences (NCATS), and NIH Roadmap for Medical Research under the following grant numbers: U01 AG027810, U01 AG042124, U01 AG042139, U01 AG042140, U01 AG042143, U01 AG042145, U01 AG042168, U01 AR066160, R01 AG066671, and UL1 TR002369. Funding for the SOF study comes from the National Institute on Aging (NIA), and the National Institute of Arthritis and Musculoskeletal and Skin Diseases (NIAMS), supported by grants (AG05407, AR35582, AG05394, AR35584, and AR35583). Funding for the Health ABC study was from the Intramural research program at the National Institute on Aging under the following contract numbers: NO1-AG-6-2101, NO1-AG-6-2103 and NO1-AG-6-2106. The Longitudinal Aging Study Amsterdam is supported by a grant from the Netherlands Ministry of Health, Welfare and Sport, Directorate of Long-Term Care. Funding for the Framingham Study comes from the National Institute of Arthritis and Musculoskeletal and Skin Diseases (NIAMS) R01 AR041398 and AR061445 and National Heart, Lung, and Blood Institute Framingham Heart Study (N01-HC-25195, HHSN268201500001I). Funding for the GOS was from the Victorian Health Promotion Foundation: ID 91-0095. For the purpose of Open Access, the author has applied a Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising from this submission.

Author contributions

Nicholas C. Harvey and Helena Johansson contributed equally to this work.

Nicholas C. Harvey (Conceptualization, Investigation, Project administration, Supervision, Writing—original draft, Writing—review & editing), Helena Johansson (Conceptualization, Data curation, Formal analysis, Writing—original draft, Writing—review & editing), Eugene V. McCloskey (Conceptualization, Data curation, Funding acquisition, Supervision, Writing—original draft, Writing—review & editing), Enwu Liu (Data curation, Formal analysis, Writing—review & editing), Kristina E. Åkesson (Data curation, Funding acquisition, Writing—review & editing), Fred A. Anderson (Data curation, Funding acquisition, Writing—review & editing), Rafael Azagra-Ledesma (Data curation, Funding acquisition, Writing—review & editing), Cecilie L. Bager (Data curation, Funding acquisition, Writing—review & editing), Charlotte Beaudart (Data curation, Funding acquisition, Writing—review & editing), Heike A. Bischoff-Ferrari (Data curation, Funding acquisition, Writing—review & editing), Emmanuel Biver (Data curation, Funding acquisition, Writing—review & editing), Olivier Bruyère (Data curation, Funding acquisition, Writing—review & editing), Jane A. Cauley (Data curation, Funding acquisition, Writing—review & editing), Jacqueline R. Center (Data curation, Funding acquisition, Writing—review & editing), Roland Chapurlat (Data curation, Funding acquisition, Writing—review & editing), Claus Christiansen (Data curation, Funding acquisition, Writing—review & editing), Cyrus Cooper (Data curation, Funding acquisition, Writing—review & editing), Carolyn J. Crandall (Data curation, Funding acquisition, Writing—review & editing), Steven R. Cummings (Data curation, Funding acquisition, Writing—review & editing), José A.P. da Silva (Data curation, Funding acquisition, Writing—review & editing), Bess Dawson-Hughes (Data curation, Funding acquisition, Writing—review & editing), Adolfo Diez-Perez (Data curation, Funding acquisition, Writing—review & editing), Alyssa B. Dufour (Data curation, Funding acquisition, Writing—review & editing), John A. Eisman (Data curation, Funding acquisition, Writing—review & editing), Petra J.M. Elders (Data curation, Funding acquisition, Writing—review & editing), Serge Ferrari (Data curation, Funding acquisition, Writing—review & editing), Yuki Fujita (Data curation, Funding acquisition, Writing—review & editing), Saeko Fujiwara (Data curation, Funding acquisition, Writing—review & editing), Claus-Christian Glüer (Data curation, Funding acquisition, Writing—review & editing), Inbal Goldshtein (Data curation, Funding acquisition, Writing—review & editing), David Goltzman (Data curation, Funding acquisition, Writing—review & editing), Vilmundur Gudnason (Data curation, Funding acquisition, Writing—review & editing), Jill Hall (Data curation, Funding acquisition, Writing—review & editing), Didier Hans (Data curation, Funding acquisition, Writing—review & editing), Mari Hoff (Data curation, Funding acquisition, Writing—review & editing), Rosemary J. Hollick (Data curation, Funding acquisition, Writing—review & editing), Martijn Huisman (Data curation, Funding acquisition, Writing—review & editing), Masayuki Iki (Data curation, Funding acquisition, Writing—review & editing), Sophia Ish-Shalom (Data curation, Funding acquisition, Writing—review & editing), Graeme Jones (Data curation, Funding acquisition, Writing—review & editing), Magnus K. Karlsson (Data curation, Funding acquisition, Writing—review & editing), Sundeep Khosla (Data curation, Funding acquisition, Writing—review & editing), Douglas P. Kiel (Data curation, Funding acquisition, Writing—review & editing), Woon-Puay Koh (Data curation, Funding acquisition, Writing—review & editing), Fjorda Koromani (Data curation, Funding acquisition, Writing—review & editing), Mark A. Kotowicz (Data curation, Funding acquisition, Writing—review & editing), Heikki Kröger (Data curation, Funding acquisition, Writing—review & editing), Timothy Kwok (Data curation, Funding acquisition, Writing—review & editing), Olivier Lamy (Data curation, Funding acquisition, Writing—review & editing), Arnulf Langhammer (Data curation, Funding acquisition, Writing—review & editing), Bagher Larijani (Data curation, Funding acquisition, Writing—review & editing), Kurt Lippuner (Data curation, Funding acquisition, Writing—review & editing), Fiona E.A. McGuigan (Data curation, Funding acquisition, Writing—review & editing), Dan Mellström (Data curation, Funding acquisition, Writing—review & editing),

Thomas Merlijn (Data curation, Funding acquisition, Writing—review & editing), Tuan V. Nguyen (Data curation, Funding acquisition, Writing—review & editing), Anna Nordström (Data curation, Funding acquisition, Writing—review & editing), Peter Nordström (Data curation, Funding acquisition, Writing—review & editing), Terence W. O'Neill (Data curation, Funding acquisition, Writing—review & editing), Barbara Obermayer-Pietsch (Data curation, Funding acquisition, Writing—review & editing), Claes Ohlsson (Data curation, Funding acquisition, Writing—review & editing), Eric S. Orwoll (Data curation, Funding acquisition, Writing—review & editing), Julie A. Pasco (Data curation, Funding acquisition, Writing—review & editing), Fernando Rivadeneira (Data curation, Funding acquisition, Writing—review & editing), Berit Schei (Data curation, Funding acquisition, Writing—review & editing), Anne-Marie Schott (Data curation, Funding acquisition, Writing—review & editing), Eric J. Shiroma (Data curation, Funding acquisition, Writing—review & editing), Kristin Siggeirsdottir (Data curation, Funding acquisition, Writing—review & editing), Eleanor M. Simonsick (Data curation, Funding acquisition, Writing—review & editing), Elisabeth Sornay-Rendu (Data curation, Funding acquisition, Writing—review & editing), Reijo Sund (Data curation, Funding acquisition, Writing—review & editing), Karin M.A. Swart (Data curation, Funding acquisition, Writing—review & editing), Pawel Szulc (Data curation, Funding acquisition, Writing—review & editing), Junko Tamaki (Data curation, Funding acquisition, Writing—review & editing), David J. Torgerson (Data curation, Funding acquisition, Writing—review & editing), Natasja M. van Schoor (Data curation, Funding acquisition, Writing—review & editing), Tjeerd P. van Staa (Data curation, Funding acquisition, Writing—review & editing), Joan Vila (Data curation, Funding acquisition, Writing—review & editing), Nicholas J. Wareham (Data curation, Funding acquisition, Writing—review & editing), Nicole C. Wright (Data curation, Funding acquisition, Writing—review & editing), Noriko Yoshimura (Data curation, Funding acquisition, Writing—review & editing), M Carola Zillikens (Data curation, Funding acquisition, Writing—review & editing), Marta Zwart (Data curation, Funding acquisition, Writing—review & editing), Liesbeth Vandendput (Data curation, Funding acquisition, Writing—review & editing), Mattias Lorentzon (Conceptualization, Investigation, Writing—original draft, Writing—review & editing), William D. Leslie (Conceptualization, Data curation, Supervision, Writing—original draft, Writing—review & editing), and John A. Kanis (Conceptualization, Data curation, Formal analysis, Funding acquisition, Methodology, Project administration, Resources, Supervision, Writing—original draft, Writing—review & editing)

Supplementary material

Supplementary material is available at *Journal of Bone and Mineral Research* online.

Funding

None declared.

Conflicts of interest

N.C.H. has received consultancy/lecture fees/honoraria/grant funding from Alliance for Better Bone Health, Amgen, MSD, Eli Lilly, Radius Health, Servier, Shire, UCB, Consilient Healthcare, Kyowa Kirin, Theramex, and Internis Pharma.

J.A.K. led the team that developed FRAX as director of the WHO Collaborating Centre for Metabolic Bone Diseases. E.V.M., W.D.L., M.L., N.C.H., E.L., L.V., and H.J. are members of the FRAX team. J.A.K., N.C.H., and E.V.M. are members of the advisory body to the National Osteoporosis Guideline Group. J.A.K. reports no additional competing interests.

K.E.Å. has no financial interest related to FRAX; chaired the National SALAR Group for Person-Centered Care Pathway Osteoporosis.

F.A.A. led the team that developed GLOW, while director of the Center for Outcomes Research at the University of Massachusetts Medical School; he has no financial interest in FRAX.

R.A. has received funding for research from Instituto Carlos III of Spanish Ministry of Health, IDIAP Jordi Gol of Catalan Government and from Scientific Societies SEMFYC and SEIOMM.

C.L.B. is employed at Nordic Bioscience and owns stock in Nordic Bioscience. She declares no competing interests in relation to this work.

H.A.B.-F. has no financial interest in FRAX. For the DO-HEALTH trial cohort, Prof. H.A.B.-F. reports independent and investigator-initiated grants from European Commission Framework 7 Research Program, from the University of Zurich, from NESTEC, from Pfizer Consumer Healthcare, from Streuli Pharma, plus non-financial support from DNP. For the study cohort extension, she reports independent and investigator-initiated grants from Pfizer and from Vifor. Further, Prof. H.A.B.-F. reports non-financial support from Roche Diagnostics and personal fees from Wild, Sandoz, Pfizer, Vifor, Mylan, Roche, Meda Pharma, outside the submitted work with regard to speaker fees and travel fees.

J.R.C. has received honoraria for speaking at educational meetings and for advisory boards from Amgen and honoraria for an advisory board from Bayer, all unrelated to this work. R.C. has no financial interest in FRAX. He has received grant funding from Amgen, UCB, Chugai, MSD, Mylan, and Medac. He has received honoraria from Amgen, UCB, Chugai, Galapagos, Biocon, Abbvie, Haoma Medica, Pfizer, Amolyt, MSD, Lilly, BMS, Novartis, Arrow, PKMed, Kyowa-Kirin, and Sanofi.

C.C. owns stock in Nordic Bioscience. He declares no competing interests in relation to this work.

C.C. reports personal fees from Alliance for Better Bone Health, Amgen, Eli Lilly, GSK, Medtronic, Merck, Novartis, Pfizer, Roche, Servier, Takeda, and UCB.

A.D.-P. reports personal fees from Amgen, Lilly, Theramex, and grants from Instituto Carlos III and owns shares of Active Life Scientific, all outside the submitted work.

J.A.E. declares consulting and research support from Actavis, Amgen, Aspen, Lilly, Merck Sharp and Dohme, Novartis, Sanofi-Aventis, Servier, and Theramex.

P.J.M.E. has no financial interest in FRAX. P.J.M.E. reports support for the SOS study by Stichting Achmea Gezondheidszorg, Achmea, and VGZ zorgverzekeraar. Additional support was given by the stichting Artsenlaboratorium en Trombosedienst. Outside the submitted work, she did receive independent investigator driven grants by Zonmw, The Netherlands, de Hartstichting, The Netherlands, The European foundation for the study of Diabetes, Amgen, The Netherlands, TEVA, The Netherlands, and Takeda, The Netherlands.

C.-C.G. reports honoraria and research support from AgNovos, Amgen, osteolabs, and UCB unrelated to this work.

D.P.K. has no financial interest in FRAX but has received support for his work in the Framingham Study over the past 30 yr by the National Institutes of Health, Astra Zeneca, Merck, Amgen, and Radius Health.

M.A.K. has received funding from the National Health and Medical Research Council (NHMRC) Australia, and the Medical Research Future Fund (MRFF) Australia. He has served on advisory boards for Amgen Australia, Novartis, and Eli Lilly—all unrelated to this work and was the Director of the Geelong Bone Densitometry Service until 2022.

A.L. is PI of the HUNT Lung and Osteoporosis project, the project got some funding from AstraZeneca for data collection. He has received lecture or consultant fees from AstraZeneca, GSK, Boehringer Ingelheim, and Diagnostica. No funding related to this study.

M.L. has received lecture fees from Amgen, Lilly, Meda, Gedeon Richter, Medison Pharma, and UCB Pharma and consulting fees from Amgen, Radius Health, UCB Pharma, Medac AB, and Parexel International, all outside the presented work.

E.V.M. has received consultancy/lecture fees/grant funding/honoraria from AgNovos, Amgen, AstraZeneca, Consilient Healthcare, Fresenius Kabi, Gilead, GSK, Hologic, Internis, Lilly, Merck, Novartis, Pfizer, Radius Health, Redx Oncology, Roche, Sanofi Aventis, UCB, ViiV, Warner Chilcott, and I3 Innovus.

C.O. is listed as a coinventor on two patent applications regarding probiotics in osteoporosis treatment.

T.W. O'N. reports honoraria from UCB unrelated to this work.

E.S.O. reports consulting fees from Angios, Biocon, Radius, and Bayer.

J.A.P. has received funding from the National Health and Medical Research Council (NHMRC) Australia, and the Medical Research Future Fund (MRFF) Australia, and Amgen, all unrelated to this work.

K.M.A.S. is an employee of the PHARMO Institute for Drug Outcomes Research. This independent research institute performs financially supported studies for government and related healthcare authorities and several pharmaceutical companies.

N.C.W. sits on the Board of Trustee of the US Bone Health and Osteoporosis Foundation, and has received consulting fees from Radius and ArgencX.

M.C.Z. reports honoraria for lectures and consulting from Amgen and Kyowa Kirin and research support from Kyowa Kirin to her institution, unrelated to this work.

M.Z. has received research funding from national societies (SEMFYC and SEIOMM).

Data availability

Data availability is cohort specific and individual cohort Principal Investigators should be contacted for enquiries about data access.

References

1. Wolk R, Berger P, Lennon RJ, Brilakis ES, Somers VK. Body mass index: a risk factor for unstable angina and myocardial infarction in patients with angiographically confirmed coronary artery disease. *Circulation*. 2003;108(18):2206-2211.
2. Johansson H, Kanis JA, Oden A, et al. A meta-analysis of the association of fracture risk and body mass index in women. *J Bone Miner Res*. 2014;29(1):223-233.
3. Nielson CM, Srikanth P, Orwoll ES. Obesity and fracture in men and women: an epidemiologic perspective. *J Bone Miner Res*. 2012;27(1):1-10.
4. Harvey NC, Kanis JA, Liu E, Johansson H, Lorentzon M, McCloskey E. Appendicular lean mass and fracture risk assessment: implications for FRAX(R) and sarcopenia. *Osteoporos Int*. 2019;30(3):537-539.
5. Neri SGR, Oliveira JS, Dario AB, Lima RM, Tiedemann A. Does obesity increase the risk and severity of falls in people aged 60 years and older? A systematic review and meta-analysis of observational studies. *J Gerontol A Biol Sci Med Sci*. 2020;75(5):952-960.
6. Compston JE, Watts NB, Chapurlat R, et al. Obesity is not protective against fracture in postmenopausal women: GLOW. *Am J Med*. 2011;124(11):1043-1050.
7. Prieto-Alhambra D, Premaor MO, Fina Aviles F, et al. The association between fracture and obesity is site-dependent: a population-based study in postmenopausal women. *J Bone Miner Res*. 2012;27(2):294-300.
8. Nethander M, Movérare-Skrttic S, Kämpe A, et al. An atlas of genetic determinants of forearm fracture. *Nat Genet*. 2023;55(11):1820-1830.
9. Schini M, Johansson H, Harvey NC, Lorentzon M, Kanis JA, McCloskey EV. An overview of the use of the fracture risk assessment tool (FRAX) in osteoporosis. *J Endocrinol Invest*. 2024;47(3):501-511.
10. CHMP. *Guideline on the Evaluation of Medicinal Products in the Treatment of Primary Osteoporosis*. European Medicines Agency. Ref CPMP/EWP/552/95Rev.2; 2006.

11. NICE. *CG146: Osteoporosis: Fragility Fracture Risk. Short Clinical Guideline- Evidence and Recommendation*. National Clinical Guideline Centre; 2012.
12. De Laet C, Kanis JA, Odén A, et al. Body mass index as a predictor of fracture risk: a meta-analysis. *Osteoporos Int*. 2005;16(11):1330-1338.
13. Vandenput L, Johansson H, McCloskey EV, et al. Update of the fracture risk prediction tool FRAX: a systematic review of potential cohorts and analysis plan. *Osteoporos Int*. 2022;33(10):2103-2136.
14. Kanis JA, Johansson H, McCloskey EV, et al. Previous fracture and subsequent fracture risk: a meta-analysis to update FRAX. *Osteoporos Int*. 2023;34(12):2027.
15. Vandenput L, Johansson H, McCloskey EV, et al. A meta-analysis of previous falls and subsequent fracture risk in cohort studies. *Osteoporos Int*. 2024;35(3):469-494.
16. Kanis JA, Johansson H, McCloskey EV, et al. Rheumatoid arthritis and subsequent fracture risk: an individual person meta-analysis to update FRAX. *Osteoporos Int*. 2025;36(4):653-671.
17. Kanis JA, Johnell O, Oden A, Dawson A, De Laet C, Jonsson B. Ten year probabilities of osteoporotic fractures according to BMD and diagnostic thresholds. *Osteoporos Int*. 2001;12(12):989-995.
18. Kanis JA, Oden A, Johnell O, Jonsson B, de Laet C, Dawson A. The burden of osteoporotic fractures: a method for setting intervention thresholds. *Osteoporos Int*. 2001;12(5):417-427.
19. Leslie WD, Schousboe JT, Morin SN, et al. Fracture risk following high-trauma versus low-trauma fracture: a registry-based cohort study. *Osteoporos Int*. 2020;31(6):1059-1067.
20. Crandall CJ, Larson JC, LaCroix AZ, et al. Risk of subsequent fractures in postmenopausal women after nontraumatic vs traumatic fractures. *JAMA Intern Med*. 2021;181(8):1055-1063.
21. Breslow NE, Day NE. *Statistical Methods in Cancer Research*. IARC Scientific Publications No 32. IARC, Lyon, France, Vol. 1987:131-135.
22. Obesity: preventing and managing the global epidemic. Report of a WHO consultation. *World Health Organ Tech Rep Ser*. 2000;894(i-xii):1-253.
23. Nielson CM, Marshall LM, Adams AL, et al. BMI and fracture risk in older men: the osteoporotic fractures in men study (MrOS). *J Bone Miner Res*. 2011;26(3):496-502.
24. Compston JE, Flahive J, Hosmer DW, et al. Relationship of weight, height, and body mass index with fracture risk at different sites in postmenopausal women: the Global Longitudinal study of Osteoporosis in Women (GLOW). *J Bone Miner Res*. 2014;29(2):487-493.
25. Stone KL, Seeley DG, Lui LY, et al. BMD at multiple sites and risk of fracture of multiple types: long-term results from the study of osteoporotic fractures. *J Bone Miner Res*. 2003;18(11):1947-1954.
26. Chalhoub D, Orwoll ES, Cawthon PM, et al. Areal and volumetric bone mineral density and risk of multiple types of fracture in older men. *Bone*. 2016;92:100-106.
27. Charles A, Mugisha A, Iconaru L, et al. Distribution of fracture sites in postmenopausal overweight and obese women: the FRIS-BEE study. *Calcif Tissue Int*. 2022;111(1):29-34.
28. Khosla S, Samakkarnthai P, Monroe DG, Farr JN. Update on the pathogenesis and treatment of skeletal fragility in type 2 diabetes mellitus. *Nat Rev Endocrinol*. 2021;17(11):685-697.
29. Karastergiou K, Smith SR, Greenberg AS, Fried SK. Sex differences in human adipose tissues - the biology of pear shape. *Biol Sex Differ*. 2012;3(1):13.
30. Gavin KM, Bessesen DH. Sex differences in adipose tissue function. *Endocrinol Metab Clin N Am*. 2020;49(2):215-228.
31. Kopelman PG, Pilkington TR, White N, Jeffcoate SL. Abnormal sex steroid secretion and binding in massively obese women. *Clin Endocrinol*. 1980;12(4):363-369.
32. Ahmed F, Hetty S, Laterveer R, et al. Altered expression of aromatase and estrogen receptors in adipose tissue from men with obesity or type 2 diabetes. *J Clin Endocrinol Metab*. 2025:dga038.
33. Trabert B, Graubard BI, Nyante SJ, et al. Relationship of sex steroid hormones with body size and with body composition measured by dual-energy X-ray absorptiometry in US men. *Cancer Causes Control*. 2012;23(12):1881-1891.
34. Mouser JG, Loprinzi PD, Loenneke JP. The association between physiologic testosterone levels, lean mass, and fat mass in a nationally representative sample of men in the United States. *Steroids*. 2016;115:62-66.
35. Xia X, Chen S, Tian X, et al. Association of body mass index with risk of cardiometabolic disease, multimorbidity and mortality: a multi-state analysis based on the Kailuan cohort. *Endocrine*. 2024;84(2):355-364.
36. Weiss EP, Jordan RC, Frese EM, Albert SG, Villareal DT. Effects of weight loss on lean mass, strength, bone, and aerobic capacity. *Med Sci Sports Exerc*. 2017;49(1):206-217.
37. *Clinical Guideline [CG189]: Obesity: Identification, Assessment and Management*. National Institute for Health and Care Excellence, London, UK; 2014.