

# On the Role of Salt, Pressure and Water Activity on Homogeneous Ice Nucleation

Jorge R. Espinosa <sup>†</sup>, Guiomar D. Soria <sup>†</sup>, Jorge Ramirez <sup>§</sup>, Chantal Valeriani <sup>‡</sup>,  
Carlos Vega <sup>†</sup>, and Eduardo Sanz <sup>†\*</sup>

<sup>†</sup>*Departamento de Química Física I, Facultad de Ciencias Químicas, Universidad  
Complutense de Madrid, 28040 Madrid, Spain.*

<sup>‡</sup>*Departamento de Física Aplicada I, Facultad de Ciencias Físicas, Universidad  
Complutense de Madrid, 28040 Madrid, Spain.*

<sup>§</sup> *Departamento de Ingeniería Química Industrial y Medio Ambiente, Escuela Técnica  
Superior de Ingenieros Industriales, Universidad Politécnica de Madrid, 28006 Madrid,  
Spain*

E-mail: esa01@quim.ucm.es

## Abstract

Pure water can be substantially supercooled below the melting temperature without transforming into ice. The achievable supercooling can be enhanced by adding solutes or by applying hydrostatic pressure. Avoiding ice formation is of great importance in the cryopreservation of food or biological samples. In this paper we investigate the similarity between the effects of pressure and salt on ice formation using a combination of state-of-the-art simulation techniques. We find that both hinder ice formation by increasing the energetic cost of creating the ice-fluid interface. Moreover, we examine

---

\*To whom correspondence should be addressed

the widely-accepted proposal that the ice nucleation rate for different pressures and solute concentrations can be mapped through the activity of water [Koop, Luo, Tsias, Peter, 406, 611, Nature, 2000]. We show that such proposal is not consistent with the nucleation rates predicted in our simulations because it does not include all parameters affecting ice nucleation. Therefore, even though salt and pressure have a qualitatively similar effect on ice formation, they cannot be quantitatively mapped onto one another.

The formation of ice from supercooled water is the most important freezing transition on Earth. Despite its relevance and ubiquity there are still many uncertainties about this phase transition. For instance, important issues such as the rate<sup>1-3</sup> and the mechanism<sup>4,5</sup> by which ice nucleates in supercooled water are still currently under debate.<sup>6</sup>

The urge for understanding in detail water freezing comes from the fact that it has a central role in key industrial and environmental processes. For example, the content of ice in clouds has a strong impact on the Earth's albedo and, therefore, on climate change.<sup>7</sup> On the other hand, a successful cryopreservation crucially depends on avoiding water freezing, that can be deleterious for the cells.<sup>8</sup>

A way to delay the formation of ice in cryopreservation protocols is to put the sample under hydrostatic pressure<sup>9</sup> to slow down ice nucleation.<sup>10</sup> Salt is also known to have a decelerating effect on ice nucleation.<sup>11</sup>

In a seminal work Koop et al. compiled experimental data of the freezing of salty aqueous solutions at different concentrations and pressures. Quite remarkably, they were able to map the ice nucleation rate of all systems into a single curve that solely depends on the activity of water in solution.<sup>12</sup> This spectacular result, based on the parallelism between the effects of salt and pressure on ice nucleation, has had a great acceptance in the scientific community studying water freezing.

The goal of the present work is to investigate the analogy between the effects of pressure and salt on water freezing by means of computer simulations. By comparing the effects of salt and pressure we are able to assess the validity of the proposal by Koop et al. To achieve

this we compare three systems: pure water at 1 bar, pure water at 2000 bar and an NaCl 1.85 m aqueous solution at 1 bar. State-of-the art simulations techniques are required to compute the ice nucleation rate and the interfacial free energy for these systems. We use the Seeding<sup>13–15</sup> and the Mold Integration<sup>16</sup> methods for that purpose. These techniques enable us to gain understanding on the parameters that affect the ice nucleation rate.

We use the TIP4P/2005 water model<sup>17</sup> combined with the Joung Cheetham/SPC/E model for NaCl.<sup>18,19</sup> We perform Molecular Dynamics simulations at constant temperature and pressure with the GROMACS package.<sup>20</sup> We refer the reader to our previous work for further simulation details.<sup>3,21</sup>

The Seeding Technique consists in simulating ice nuclei embedded in the supercooled fluid (either pure or salty water). We insert pure spherical ice Ih seeds as that shown in the graphical abstract. The embedded ice cluster configuration is equilibrated as described in Ref.<sup>22</sup> We make sure the cluster interface is well equilibrated by looking at the ionic density profile<sup>23</sup>. Once we get an equilibrated configuration of an ice cluster embedded in the solution we monitor the evolution of the nucleus at different temperatures. For temperatures higher than that at which the inserted cluster is critical, the cluster melts, and viceversa.

In this way we obtain an estimate for the temperature at which the inserted cluster with  $N_c$  molecules is critical. We determine  $N_c$  using local-bond order parameters<sup>24</sup> as done in our previous work.<sup>14,15,25</sup> We use large system sizes to ensure that concentration changes are negligible as the cluster grows or melts in the brine solution.<sup>23</sup> In Fig. 1(a) we show  $N_c$  versus the supercooling,  $\Delta T$ , which is the melting temperature,  $T_m$ , minus the temperature of interest (the model melting temperatures for pure ice in coexistence with pure water at 1 and 2000 bar<sup>26</sup> and with a 1.85 m NaCl aqueous solution at 1 bar<sup>23</sup> are 250, 227 and 240.5 K, respectively). Clearly, for a given  $\Delta T$ , the number of molecules required to reach the critical size is larger in salty or compressed water than in pure water at normal pressure. This is consistent with the experimental observation that salt and pressure hinder ice nucleation.<sup>11</sup>

To quantify the decelerating effect of salt and pressure on homogeneous ice nucleation




Figure 1: Plotted for pure (black), salty (red) and compressed water (blue) as a function of the supercooling: (a) Number of particles in the critical cluster; (b) Water chemical potential difference between the liquid and the solid phases; (c) decimal logarithm of the nucleation rate; (d) ice-water interfacial free energy. Symbols in (d) for  $\Delta T = 0$  and  $\Delta T > 0$  correspond to our calculations of  $\gamma$  with MI and Seeding, respectively. Shaded regions in (c) and (d) indicate the error bar.

we compute the nucleation rate,  $J$ . In the Seeding method, simulations are used to obtain the parameters on which the  $J$  expression given by Classical Nucleation Theory (CNT)<sup>27–29</sup> depends:<sup>14,15,30</sup>

$$J = \sqrt{\frac{-\Delta\mu_w}{6\pi k_B T N_c}} \rho_f f^+ \exp\left(\frac{N_c \Delta\mu_w}{2k_B T}\right). \quad (1)$$

Such parameters are the fluid number density,  $\rho_f$ , the number of particles in the critical cluster  $N_c$  (Fig. 1(a)), the chemical potential difference of water in ice and in the liquid,  $\Delta\mu_w = \mu_w^i - \mu_w$  (Fig. 1(b)), and the frequency with which particles attach to the critical cluster (the attachment rate). Further details on the calculation of these factors will be given in a forthcoming publication.<sup>23</sup> The nucleation rate as a function of the supercooling

is plotted in Fig. 1(c) for the three systems under comparison. For a given supercooling, pure water at normal pressure (in black) has the highest nucleation rate. In other words, the supercooling required to freeze water increases when adding salt (in red) or applying pressure (in blue), in agreement with the experimental trend.<sup>10,11</sup>

Since we have access to all variables required for the computation of  $J$  we can rationalise which is the main factor behind the pressure/salt-induced deceleration of ice nucleation. The nucleation rate is given by  $J = Ae^{-\Delta G_c/(k_B T)}$  where  $A$  is a kinetic pre-factor and  $\Delta G_c/(k_B T)$  is the Gibbs free energy barrier required to form a critical ice nucleus in the supercooled fluid. We have checked that  $A$  does not change by more than one or two orders of magnitude between different systems for a given supercooling, which is insignificant as compared to the large differences between the  $J$  curves in Fig. 1(c). The differences in  $J$  must be then ascribed to changes in  $\Delta G_c/(k_B T)$ . Within CNT,  $\Delta G_c/(k_B T)$  is proportional to the third power of the ice-liquid interfacial free energy,  $\gamma$ , and inversely proportional to the second power of  $|\Delta\mu_w|$ . The latter does not significantly change from one system to another (see Fig. 1(b)). Therefore, the key to the decrease of  $J$  must be found in an increase of  $\gamma$ . This is in fact what we concluded in our recent study of ice nucleation under high pressure.<sup>31</sup> In Fig. 1(d) we represent  $\gamma$  versus the supercooling for the three studied systems (the points for the supercooled fluid have been obtained with the Seeding method whereas those at coexistence,  $\Delta T = 0$ , with the Mold Integration method<sup>16,21</sup>). Indeed, the interfacial free energy increases when adding salt or applying pressure for a given supercooling. Therefore, both pressure and salt decelerate ice nucleation by increasing the ice-liquid interfacial free energy, which is one of the main conclusions of our work. This is a valuable simulation prediction given that there is no experimental consensus for the value of the ice-liquid interfacial free energy (not even for the case of pure water at 1 bar<sup>32</sup>).

The fact that pressure and salt hinder ice nucleation for the same reason is in principle consistent with the proposition that applying a certain pressure should have the same effect on ice nucleation as adding a given amount of salt.<sup>12</sup> Such proposal is the basis of the so-called

water-activity-based ice nucleation conjecture (WAB-INC).<sup>12,33</sup> The WAB-INC is based on an analysis of the freezing of salty water drops at different pressures and proposes that the nucleation rate depends only on the activity of water,  $a_w$ . This idea is extremely appealing because it enables to obtain the nucleation rate for any solution at any pressure with a fit solely depending on a measurable thermodynamic parameter such as the activity. Although the WAB-INC apparently works<sup>12</sup> it has never been carefully checked for three reasons: (i)  $a_w$  cannot be experimentally measured for deeply supercooled solutions due to the formation of ice (in fact, in many cases it is assumed to be independent of temperature<sup>33</sup>); (ii) the general character of the theory has not been tested since the  $J$ -range that can be measured is narrow (it is limited by the sample's volume and the cooling rate); (iii) the physical basis of the WAB-INC is unclear since many of the parameters that affect ice nucleation, like  $\gamma$ ,  $N_c$  or  $\Delta\mu_w$  for  $T < T_m$ , cannot be measured experimentally. Our simulations do not have these shortcomings and enable us to test the validity of the WAB-INC in a wide range of nucleation rates.

To start with, we evaluate the temperature dependence of  $a_w$  in the brine solution at 1 bar and with salt concentration  $c = 1.85$  m via:

$$\mu_w(c, T, 1) = \mu_w(0, T, 1) + RT \ln[a_w(c, T, 1)] \quad (2)$$

In practise, the chemical potential difference with and without salt at 1 bar,  $\mu_w(c, T, 1) - \mu_w(0, T, 1)$ , is obtained as  $\Delta\mu_w(0, T, 1) - \Delta\mu_w(c, T, 1)$ , the difference between the black and the red curves in Fig. 1(b) (the chemical potential of water in ice cancels out when both  $\Delta\mu_w$ 's are subtracted). In Fig. 2 we plot  $a_w(c, T, 1)$  as a function of the supercooling (red curve). To our knowledge, this is the first time that the temperature dependence of water activity is reported up to such deep supercooling either in simulations or experiments (there are simulation works that report activities as a function of concentration at constant temperature<sup>34</sup>). As explained below, the cross with the orange curve (activity of water

coexisting with ice) corresponds to the melting temperature, above which the water activity remains almost constant, in accordance with experimental observations for most simple ionic solutions.<sup>33</sup> By contrast,  $a_w$  sharply increases below the melting temperature.

The activity of pure water under pressure is in principle 1. However, in the framework of the WAB-INC, pure water under pressure  $p$  is effectively assigned the activity of a water solution at normal pressure having the same chemical potential difference with ice,  $\Delta\mu_w$ :

$$\mu_w(0, T, p) - \mu_w^i(0, T, p) = \quad (3)$$

$$\mu_w(0, T, 1) + RT \ln a_w(c^{eff}, T, 1) - \mu_w^i(0, T, 1) \quad (4)$$

where  $c^{eff}$  is the effective concentration required to satisfy the equality above. Thus, we can compute  $a_w(c^{eff}, T, 1)$  as  $\exp[(\Delta\mu_w(0, T, 1) - \Delta\mu_w(0, T, p))/(RT)]$  (the exponent is obtained by subtracting the black and the blue curves in Fig. 1(b)). In Fig. 2 we show  $a_w(c^{eff}, T, 1)$  as a function of temperature (blue curve). Again, the activity sharply raises below the melting temperature of compressed water, given by the crossing with the orange curve. Therefore, according to our simulations, the assumption that  $a_w$  does not depend on temperature for most ionic solutions<sup>12,33</sup> cannot be safely made, especially at low temperatures where water behaves anomalously.<sup>35,36</sup>

As pointed out in Ref.,<sup>33</sup> the validity of the WAB-INC does not rely on whether or not  $a_w$  depends on temperature.<sup>12</sup> What the WAB-INC really proposes is that  $J(\Delta a_w)$  is a universal curve for any pressure or concentration.  $\Delta a_w$  is the difference, for a given temperature, between the activity of water in the solution (or in compressed water) and the activity of water at coexistence with ice,  $a_w^i$ . The latter can be obtained through:

$$\mu_w^i(0, T, 1) = \mu_w(0, T, 1) + RT \ln[a_w^i(c, T, 1)] \quad (5)$$

The chemical potential difference  $\mu_w^i(0, T, 1) - \mu_w(0, T, 1)$  is simply  $\Delta\mu_w$  for pure water at 1 bar (black curve in Fig. 1(b)).  $a_w^i$  is plotted in orange in Fig. 2 (at the melting point the




Figure 2: Water activity as a function of  $T - T_m^0$ , the difference between the temperature of interest and the melting temperature of pure water ( $T_m^0 = 250$  K for the model). In red we plot the activity of water in the studied 1.85 m solution and in blue the effective activity for pure water at 2000 bar as defined in the main text. In orange we plot the activity of water coexisting with ice.

orange and the red/blue curves cross because the activity of water becomes equal to  $a_w^i$ ). Thus, to obtain  $\Delta a_w$  for the studied solution we have to subtract, for a given temperature, the red and the orange curves in Fig. 2, whereas for pure water under pressure we compute the difference between the blue and the orange curves (see red and blue arrows in Fig. 2, respectively). Knowing  $\Delta a_w(T)$  and  $J(T)$  (Fig. 1(c)) we obtain  $J(\Delta a_w)$ . In Fig. 3(a) we plot  $J(\Delta a_w)$  for compressed and salty water. Both curves are not the same for every  $\Delta a_w$ . Therefore, the WAB-INC proposal of a universal  $J(\Delta a_w)$  curve is not consistent with our simulation predictions. However, the  $J(\Delta a_w)$  curves lie close to each other beyond the rate value measured in typical experiments,  $J \sim 10^{16} m^{-3} s^{-1}$ , given by the horizontal green line in Fig. 3(a). Thus, our results are compatible with the fact that the WAB-INC may seem to work when data of typical freezing experiments are analysed.<sup>12</sup> Why a theory that is not general may seemingly work at high values of  $\Delta a_w$ ?

On the one hand, it is perhaps worth noting that the WAB-INC is not as neat as it may appear. In the publication where the WAB-INC was proposed variations of up to 15-20



K around the universal freezing curve can be clearly noticed.<sup>12</sup> In that temperature range there may be many orders of magnitude difference in the nucleation rate. On the other hand, the WAB-INC may be a particular case of CNT in certain limits. Since in the WAB-INC framework the rate solely depends on the activity of the solvent (chemical potential), the WAB-INC will be compatible with CNT if, for a given  $\Delta a_w$ , all other parameters that affect ice nucleation are the same. We check if this is the case for the ice-fluid interfacial free energy, a parameter that we have shown to be central in understanding the effect of salt (this work) and pressure (Ref.<sup>31</sup>) on homogeneous ice nucleation. Combining  $\Delta a_w(T)$  and  $\gamma(T)$  (Fig. 1(d)) we obtain  $\gamma(\Delta a_w)$ , shown in Fig. 3(b). Clearly,  $\gamma$  is not the same for salty and compressed water at low  $\Delta a_w$  and becomes increasingly similar as  $\Delta a_w$  increases. When both  $\Delta a_w$  and  $\gamma$  are similar, the WAB-INC may seem to work, but this is just a particular case of CNT. That  $\gamma$  becomes similar between different systems at high  $\Delta a_w$  may be related with the nucleation of ice clusters in solute-depleted regions when the critical cluster size becomes small.<sup>37</sup> With our analysis we conclude that CNT is a much more comprehensive theoretical framework than the WAB-INC to understand ice nucleation and that the WAB-INC may only seemingly work as a particular case of CNT at high nucleation rates. The fact that the simulation approach employed in this work provides estimates of the nucleation rate in a much wider range than current experiments has clearly evidenced the deficiencies of the WAB-INC. We expect our work to inspire future attempts to derive phenomenological frameworks to describe ice nucleation that do not necessarily comply with the WAB-INC.<sup>38</sup>

Our conclusions are based on the results obtained with a water model. One may wonder if our observations can be safely applied to real water as well. We argue that if WAB-INC had a strong physical foundation it should be able to explain ice nucleation both in real water and in any family of realistic water models. Our model is quite realistic because it closely reproduces the behaviour of real water for many thermodynamic and dynamic properties,<sup>39</sup> including anomalies<sup>40</sup> and, most importantly, homogeneous ice nucleation in pure<sup>14,30</sup> and salty water.<sup>23</sup> Moreover, in this work we show that the model captures the experimentally

observed deceleration of ice nucleation with pressure and salt<sup>10,12</sup> (Fig. 1(c)), as well as the fact that the WAB-INC may seemingly work in the range of nucleation rates typically accessible to experiments (see Fig. 3(a)). Furthermore, we do not need a perfect model for the NaCl ions to test the WAB-INC, as the theory should in principle work for a wide variety of solutes. So even if the simulated solute does not behave exactly as real dissolved NaCl, it can be taken as any arbitrary solute with which testing the WAB-INC. Be it as it may, we hope our work will motivate future experiments to definitely confirm our prediction. As a matter of fact, there have been already experiments reporting inconsistencies with the WAB-INC.<sup>41</sup>




Figure 3: (a) Nucleation rate of salty (1.85 m NaCl) and compressed (2000 bar) water as a function of  $\Delta a_w$ . (b) Ice-fluid interfacial free energy as a function of  $\Delta a_w$ .

Our Seeding approach to ice nucleation relies on two assumptions: (i) the proposed structure of the critical cluster is the correct one, and (ii) the formation free energy of such cluster can be obtained via CNT. Our guess for the critical nucleus structure is that of a

spherical pure ice Ih cluster. We use ice Ih because this structure, or stacking mixtures of ice Ic and Ih, has shown to be that of critical ice clusters at 1 bar.<sup>4,14,25,31,42</sup> We insert pure ice seeds disregarding the possibility that NaCl ions enter the ice lattice. This approximation is inspired by the experimental phase diagram, where the brine coexists with pure ice.<sup>43</sup> To make sure this is a good approximation we have performed a long (260 ns) direct coexistence simulation below the melting temperature and computed the fraction of ions incorporated into the grown ice lattice. Such fraction was smaller than 0.2 per cent, a value in agreement with previous simulation work<sup>44,45</sup> and sufficiently small to justify our approximation. We use spherical clusters, consistently with previous studies showing that the cluster’s shape quickly equilibrates into a sphere<sup>25</sup> –with thermal fluctuations typical of a rough ice-fluid interface.<sup>46,47</sup> Assumption (ii), in turn, is expected to be satisfied for the large critical cluster sizes used in this work. Independent evidence of the validity of CNT to describe the free energy of crystal cluster formation can be found, e. g., in Ref.<sup>48</sup> As long as conditions (i) and (ii) are satisfied, one can use seeding to obtain nucleation rates regardless the pathway leading to the formation of the critical cluster is CNT-like (a one-by-one addition of particles to a growing crystal cluster) or a more sophisticated one like composition fluctuations followed by the crystal cluster growth.<sup>37,49–51</sup> Thus, although seeding does not provide any information on the way the critical cluster is formed, it can be used to obtain nucleation rates,<sup>15</sup> even in systems where the formation of the critical cluster has been reported to be two-step, like crystallization in hard spheres<sup>52,53</sup> or in mw-water.<sup>54</sup> In any case, since we are dealing with the crystallization of the solvent and with large critical clusters whose size exceeds that of typical composition fluctuations we do not expect the formation mechanism of the critical nucleus to be markedly two-step. It is also worth noting that seeding has been successful in predicting crystal nucleation rates in the Lennard Jones and the Tosi-Fumi NaCl systems<sup>15</sup> as well, which gives further confidence in the predictions made in this paper. Moreover, in this work we find consistent values between the  $\gamma$  obtained at coexistence with Mold Integration, a method that does not rely on CNT and that does not suffer from strong finite

size effects,<sup>16</sup> and that obtained below the melting temperature with seeding (see Fig. 1(d)). This consistency test, and the considerations above mentioned, strongly support the validity of the approach followed in this work.

In summary, we compare the effects of pressure and salt on homogeneous ice nucleation. We find that both, pressure and salt, decelerate ice nucleation by increasing the ice-liquid interfacial free energy. Despite this qualitative similarity, it is not possible to quantitatively map ice nucleation rates of salty water onto those of compressed water through the activity of water. Our results question the validity of the so-called water-activity-based ice nucleation theory.<sup>12</sup>

Understanding and quantifying the effect of salt on homogeneous ice nucleation can be useful to develop climate change models.<sup>7</sup> On the other hand, improving cryopreservation protocols, aimed at vitrifying biological samples by averting ice nucleation and growth upon cooling, requires deep understanding of the effect of freezing preventing factors like pressure, salt or combinations of both.<sup>8,9</sup> Our work may also have implications in the experimental search of the putative liquid-liquid transition of water,<sup>55-58</sup> that in some cases uses pressure and solutes as strategies to approach the so-called no-man's-land.<sup>36,59,60</sup>

## Acknowledgement

This work was funded by grants FIS2013/43209-P and FIS2016-78117-P FIS2016-78847-P of the MEC, and the UCM/Santander 910570 and PR26/16-10B-2. C.Valeriani and E. Sanz acknowledge financial support from a Ramon y Cajal Fellowship. J. R. Espinosa acknowledges financial support from the FPI grant BES-2014-067625. Calculations were carried out in the supercomputer facilities La Palma and Magerit from the Spanish Supercomputing Network (RES) (projects QCM-2015-3-0036 and QCM-2016-1-0039). The authors acknowledge the computer resources and technical assistance provided by the Centro de Supercomputacion y Visualizacion de Madrid (CeSViMa).

## References

- (1) Laksmono, H. et al. Anomalous Behavior of the Homogeneous Ice Nucleation Rate in No-Mans Land. *The Journal of Physical Chemistry Letters* **2015**, *6*, 2826–2832.
- (2) Koop, T.; Murray, B. J. A physically constrained classical description of the homogeneous nucleation of ice in water. *The Journal of Chemical Physics* **2016**, *145*, 211915.
- (3) Espinosa, J. R.; Navarro, C.; Sanz, E.; Valeriani, C.; Vega, C. On the time required to freeze water. *The Journal of Chemical Physics* **2016**, *145*, 211922.
- (4) Malkin, T. L.; Murray, B. J.; Brukhno, A. V.; Anwar, J.; Salzmann, C. G. Structure of ice crystallized from supercooled water. *Proceedings of the National Academy of Sciences* **2012**, *109*, 1041–1045.
- (5) Russo, J.; Romano, F.; Tanaka, H. New metastable form of ice and its role in the homogeneous crystallization of water. *Nature Materials* **2014**, *13*, 733.
- (6) Sosso, G. C.; Chen, J.; Cox, S. J.; Fitzner, M.; Pedevilla, P.; Zen, A.; Michaelides, A. Crystal Nucleation in Liquids: Open Questions and Future Challenges in Molecular Dynamics Simulations. *Chemical Reviews* **2016**, *116*, 7078–7116.
- (7) Cantrell, W.; Heymsfield, A. Production of ice in tropospheric clouds. *Bull. Amer. Meteor. Soc.* **2005**, *86*, 795.
- (8) Morris, G. J.; Acton, E. Controlled ice nucleation in cryopreservation—a review. *Cryobiology* **2013**, *66*, 85.
- (9) Studer, D. High-pressure freezing system. 2001; US Patent 6,269,649.
- (10) Kanno, H.; Speedy, R. J.; Angell, C. A. Supercooling of Water to  $-92^{\circ}\text{C}$  Under Pressure. *Science* **1975**, *189*, 880–881.

- (11) Kanno, H.; Angell, C. A. Homogeneous nucleation and glass formation in aqueous alkali halide solutions at high pressures. *The Journal of Physical Chemistry* **1977**, *81*, 2639–2643.
- (12) Koop, T.; Luo, B.; Tsias, A.; Peter, T. Water activity as the determinant for homogeneous ice nucleation in aqueous solutions. *Nature* **2000**, *406*, 611–614.
- (13) Knott, B. C.; Molinero, V.; Doherty, M. F.; Peters, B. Homogeneous Nucleation of Methane Hydrates: Unrealistic under Realistic Conditions. *J. Am. Chem. Soc.* **2012**, *134*, 19544–19547.
- (14) Sanz, E.; Vega, C.; Espinosa, J. R.; Caballero-Bernal, R.; Abascal, J. L. F.; Valeriani, C. Homogeneous Ice Nucleation at Moderate Supercooling from Molecular Simulation. *Journal of the American Chemical Society* **2013**, *135*, 15008–15017.
- (15) Espinosa, J. R.; Vega, C.; Valeriani, C.; Sanz, E. Seeding approach to crystal nucleation. *J. Chem. Phys.* **2016**, *144*, 034501.
- (16) Espinosa, J. R.; Vega, C.; Sanz, E. The mold integration method for the calculation of the crystal-fluid interfacial free energy from simulations. *J Chem. Phys.* **2014**, *141*, 134709.
- (17) Abascal, J. L. F.; Vega, C. A general purpose model for the condensed phases of water: TIP4P/2005. *J. Chem. Phys.* **2005**, *123*, 234505.
- (18) Joung, I. S.; Cheatham, T. E. Determination of Alkali and Halide Monovalent Ion Parameters for Use in Explicitly Solvated Biomolecular Simulations. *The Journal of Physical Chemistry B* **2008**, *112*, 9020–9041.
- (19) Benavides, A. L.; Aragonés, J. L.; Vega, C. Consensus on the solubility of NaCl in water from computer simulations using the chemical potential route. *The Journal of Chemical Physics* **2016**, *144*, 124504.

- (20) Lindahl, E.; Hess, B.; van der Spoel, D. GROMACS 3.0: a package for molecular simulation and trajectory analysis. *J. Mol. Model.* **2001**, *7*, 306.
- (21) Espinosa, J. R.; Vega, C.; Sanz, E. Ice-Water interfacial free energy for the TIP4P, TIP4P/2005, TIP4P/ICE and mW models as obtained from the mold integration technique. *The Journal of Physical Chemistry C* **2016**, *120*, 8068–8075.
- (22) Espinosa, J. R.; Vega, C.; Valeriani, C.; Sanz, E. The crystal-fluid interfacial free energy and nucleation rate of NaCl from different simulation methods. *J. Chem. Phys.* **2015**, *142*, 194709.
- (23) Soria, G. D.; Espinosa, J. R.; Ramirez, J.; Valeriani, C.; Vega, C.; Sanz, E. A simulation study of homogeneous ice nucleation in supercooled salty water. *To be published* **2017**, -, -.
- (24) Lechner, W.; Dellago, C. Accurate determination of crystal structures based on averaged local bond order parameters. *J. Chem. Phys.* **2008**, *129*, 114707.
- (25) Zaragoza, A.; Conde, M. M.; Espinosa, J. R.; Valeriani, C.; Vega, C.; Sanz, E. Competition between ices Ih and Ic in homogeneous water freezing. *The Journal of Chemical Physics* **2015**, *143*, 134504.
- (26) Conde, M. M.; Gonzalez, M. A.; Abascal, J. L. F.; Vega, C. Determining the phase diagram of water from direct coexistence simulations: The phase diagram of the TIP4P/2005 model revisited. *The Journal of Chemical Physics* **2013**, *139*, 154505.
- (27) Kelton, K. F. *Crystal Nucleation in Liquids and Glasses*; Academic: Boston, 1991.
- (28) Volmer, M.; Weber, A. Keimbildung in übersättigten Gebilden. *Z. Phys. Chem.* **1926**, *119*, 277.
- (29) Becker, R.; Döring, W. Kinetische behandlung der keimbildung in übersättigten dampfen. *Ann. Phys.* **1935**, *416*, 719–752.

- (30) Espinosa, J. R.; Sanz, E.; Valeriani, C.; Vega, C. Homogeneous ice nucleation evaluated for several water models. *J. Chem. Phys.* **2014**, *141*, 18C529.
- (31) Espinosa, J. R.; Zaragoza, A.; Rosales-Pelaez, P.; Navarro, C.; Valeriani, C.; Vega, C.; Sanz, E. Interfacial Free Energy as the Key to the Pressure-Induced Deceleration of Ice Nucleation. *Phys. Rev. Lett.* **2016**, *117*, 135702.
- (32) Ickes, L.; Welti, A.; Hoose, C.; Lohmann, U. Classical nucleation theory of homogeneous freezing of water: thermodynamic and kinetic parameters. *Phys. Chem. Chem. Phys.* **2015**, *17*, 5514–5537.
- (33) Koop, T. Homogeneous Ice Nucleation in Water and Aqueous Solutions. *Zeitschrift für Physikalische Chemie* **2004**, *218*, 1231.
- (34) Mester, Z.; Panagiotopoulos, A. Z. Mean ionic activity coefficients in aqueous NaCl solutions from molecular dynamics simulations. *The Journal of Chemical Physics* **2015**, *142*, 044507.
- (35) Debenedetti, P. G. Supercooled and glassy water. *J. Phys.:Condens. Matter* **2003**, *15*, R1669–R1726.
- (36) Gallo, P. et al. Water: A Tale of Two Liquids. *Chemical Reviews* **2016**, *116*, 7463–7500.
- (37) Bullock, G.; Molinero, V. Low-density liquid water is the mother of ice: on the relation between mesostructure, thermodynamics and ice crystallization in solutions. *Faraday Discuss.* **2013**, *167*, 371–388.
- (38) Barahona, D. Analysis of the effect of water activity on ice formation using a new thermodynamic framework. *Atmos. Chem. Phys.* **2014**, *14*, 7665–7680.
- (39) Vega, C.; Abascal, J. L. F. Simulating water with rigid non-polarizable models: a general perspective. *Phys. Chem. Chem. Phys.* **2011**, *13*, 19663–19688.



- (40) Pi, H. L.; Aragoñes, J. L.; Vega, C.; Noya, E. G.; Abascal, J. L.; Gonzalez, M. A.; McBride, C. Anomalies in water as obtained from computer simulations of the TIP4P/2005 model: density maxima, and density, isothermal compressibility and heat capacity minima. *Molecular Physics* **2009**, *107*, 365–374.
- (41) Swanson, B. D. How Well Does Water Activity Determine Homogeneous Ice Nucleation Temperature in Aqueous Sulfuric Acid and Ammonium Sulfate Droplets? *Journal of the Atmospheric Sciences* **2009**, *66*, 741–754.
- (42) Malkin, T. L.; Murray, B. J.; Salzmänn, C. G.; Molinero, V.; Pickering, S. J.; Whale, T. F. Stacking disorder in ice I. *Phys. Chem. Chem. Phys.* **2015**, *17*, 60–76.
- (43) Koop, T.; Kapilashrami, A.; Molina, L. T.; Molina, M. J. Phase transitions of sea-salt/water mixtures at low temperatures: Implications for ozone chemistry in the polar marine boundary layer. *Journal of Geophysical Research: Atmospheres* **2000**, *105*, 26393–26402.
- (44) Conde, M. M.; Rovere, M.; Gallo, P. Spontaneous NaCl-doped ice at seawater conditions: focus on the mechanisms of ion inclusion. *Phys. Chem. Chem. Phys.* **2017**, *19*, 9566–9574.
- (45) Bauerecker, S.; Ulbig, P.; Buch, V.; Vrbka, L.; Jungwirth, P. Monitoring Ice Nucleation in Pure and Salty Water via High-Speed Imaging and Computer Simulations. *The Journal of Physical Chemistry C* **2008**, *112*, 7631–7636.
- (46) Benet, J.; MacDowell, L. G.; Sanz, E. A study of the ice-water interface using the TIP4P/2005 water model. *Phys. Chem. Chem. Phys.* **2014**, *16*, 22159–22166.
- (47) Benet, J.; Llombart, P.; Sanz, E.; MacDowell, L. G. Premelting-Induced Smoothing of the Ice-Vapor Interface. *Phys. Rev. Lett.* **2016**, *117*, 096101.

- (48) Koss, P.; Statt, A.; Virnau, P.; Binder, K. Free Energy Barriers for Crystal Nucleation from Fluid Phases. *arXiv* **2017**, -, 1705.08216.
- (49) Agarwal, V.; Peters, B. *Advances in Chemical Physics: Volume 155*; John Wiley & Sons, Inc., 2014; pp 97–160.
- (50) Vekilov, P. G. The two-step mechanism of nucleation of crystals in solution. *Nanoscale* **2010**, *2*, 2346–2357.
- (51) Black, S. Simulating nucleation of molecular solids. *Proc. R. Soc. A* **2007**, *463*, 2799–2811.
- (52) Schilling, T.; Schöpe, H. J.; Oettel, M.; Opletal, G.; Snook, I. Precursor-Mediated Crystallization Process in Suspensions of Hard Spheres. *Phys. Rev. Lett.* **2010**, *105*, 025701.
- (53) Russo, J.; Tanaka, H. The microscopic pathway to crystallization in supercooled liquids. *Scientific Reports* **2012**, *2*, 505.
- (54) Moore, E. B.; Molinero, V. Structural transformation in supercooled water controls the crystallization rate of ice. *Nature* **2011**, *479*, 506–508.
- (55) Poole, P. H.; Sciortino, F.; Essmann, U.; Stanley, H. E. Phase Behavior of Metastable Water. *Nature* **1992**, *360*, 324.
- (56) Mishima, O.; Stanley, H. E. The relationship between liquid, supercooled and glassy water. *Nature* **1998**, *396*, 329–335.
- (57) Manka, A.; Pathak, H.; Tanimura, S.; Wolk, J.; Strey, R.; Wyslouzil, B. E. Freezing water in no man’s land. *Phys. Chem. Chem. Phys.* **2012**, *14*, 4505–4516.
- (58) Palmer, J. C.; Martelli, F.; Liu, Y.; Car, R.; Panagiotopoulos, A. Z.; Debenedetti, P. G. Metastable liquid-liquid transition in a molecular model of water. *Nature* **2014**, *510*, 385–388.

- (59) Corradini, D.; Rovere, M.; Gallo, P. A route to explain water anomalies from results on an aqueous solution of salt. *The Journal of Chemical Physics* **2010**, *132*, 134508.
- (60) Caupin, F. Escaping the no man's land: Recent experiments on metastable liquid water. *Journal of Non-Crystalline Solids* **2015**, *407*, 441–448.

# Graphical TOC Entry

